ORIGINAL RESEARCH

D.A. Lewis
Y.H. Ding
D. Dai
R. Kadirvel
M.A. Danielson
H.J. Cloft
D.F. Kallmes

Morbidity and Mortality Associated with Creation of Elastase-Induced Saccular Aneurysms in a Rabbit Model

BACKGROUND AND PURPOSE: Elastase-induced aneurysms in rabbits have been proposed as a useful preclinical tool for device development. The object of this study was to report rates of morbidity and mortality associated with the creation and embolization of elastase-induced rabbit aneurysms and to assess the impact of operator experience on these rates.

MATERIALS AND METHODS: Elastase-induced model aneurysms were created in New Zealand white rabbits ($n = 700$). One neuroradiologist/investigator, naive to the aneurysm-creation procedure at the outset of the experiments, performed all surgeries. All morbidity and deaths related to aneurysm creation ($n = 700$) and embolization procedures ($n = 529$) were categorized into acute and chronic deaths. Data were analyzed with single-regression analysis and analysis of variance. To assess the impact of increasing operator experience, we broke the number of animals into 50-animal increments.

RESULTS: There were 121 (17%) deaths among 700 subjects. Among 700 aneurysm-creation procedures, 59 deaths (8.4%) were noted. Among 529 aneurysm-embolization procedures, 43 deaths (8.1%) were noted. Nineteen additional deaths (2.7% of 700 subjects) were unrelated to the procedures. Simple regression-indicated mortality associated with procedures diminished with increasing operator experience ($R^2 = 0.38$, $P = .0180$), and that for each 50-rabbit increment mortality was reduced, on average, by 0.6%.

CONCLUSIONS: Mortality rates of approximately 8% are associated with both experimental aneurysm creation and with embolization in the rabbit elastase-induced aneurysm model. Increasing operator experience is inversely correlated with mortality, and the age of the rabbit is positively associated with morbidity.

The development of endovascular occlusion devices has been facilitated through the use of various animal models of saccular aneurysms. Detachable platinum coils have been tested by using surgically created aneurysms in swine, canines, rabbits, and primates.¹⁻⁷ More recently, elastase-induced aneurysms in rabbits have been proposed as a useful preclinical tool for device development.⁸⁻¹⁴

Previous studies^{9,10,15} have shown that elastase-induced aneurysms in rabbits are similar to human aneurysms in size and shape. A recent study of the histopathology of the rabbit model¹⁶ has shown that it mimics well the healing seen in human aneurysms, especially when compared with the swine model. Furthermore, because these experimental aneurysms are in close proximity to the aorta and are situated on a curved vessel, they also are exposed to substantial hemodynamic stress.

We report rates of morbidity and mortality associated with model aneurysm creation and correlate these rates with operator experience and type of procedure.

Materials and Methods

All animals used in this study underwent surgeries that were approved by the Animal Care and Use Committee of our institution. Seven hundred female white New Zealand rabbits were used in this retrospective study. The single investigator who performed all the surgeries was a trained neuroradiologist who was naive to the aneurysm-creation procedure at the outset of the experiments. All the rabbits used in this study were originally used as part of another investigation. The original investigation was entirely unrelated to this project.

Aneurysm Creation

The surgical procedures used to create an aneurysm have been described elsewhere.^{9,17} Briefly, female New Zealand white rabbits (2–4 kg) were anesthetized with ketamine (74 mg/kg), xylazine (5 mg/kg), and acepromazine (1 mg/kg). Using a sterile technique through a vertical midline incision in the neck, the trachea and the right carotid artery were identified. The vagus nerve and jugular vein were gently dissected free from the right common carotid artery (CCA). A 5F sheath (Cordis Endovascular, Miami Lakes, Fla) was advanced retrograde in the CCA to a point approximately 3 cm cephalad to the CCA origin. Through this indwelling sheath, a 3F Fogarty balloon (Baxter Healthcare, Irvine, Calif) was advanced to the origin of the right CCA at its junction with the subclavian artery. The balloon was inflated with iodinated contrast with just enough pressure to achieve flow arrest in the CCA. Porcine elastase (5.23 U/mgP, 40.1 mgP/mL, approximately 200 U/mL; Worthington Biochemical, Lakewood, NJ) mixed with saline and iodinated contrast was incubated in the dead space of the CCA above the inflated balloon for approximately 20 minutes. Following incubation of the elastase solution, the balloon and sheath were removed, the CCA was ligated below the sheath entry site, and the incision was closed in 2 layers. The fascial layers

Received May 14, 2008; accepted after revision July 2.

From the Neuroradiology Research Laboratory, Department of Radiology, Mayo Clinic, Rochester, Minn.

This work was funded in part by National Institutes of Health Grants NS42646 and HL72247.

Please address correspondence to Debra A. Lewis, PhD, Mayo Clinic, 200 First St SW, Rochester, MN 55905; e-mail: lewis.debra@mayo.edu

Indicates open access to non-subscribers at www.ajnr.org

DOI 10.3174/ajnr.A1369

92 Lewis | AJNR 30 | Jan 2009 | www.ajnr.org

Fig 1. Bar chart shows the number of deaths due to all procedures (aneurysm creation and embolization) over 50-rabbit increments. The line indicates the linear regression for these data. In addition, the linear regression equation [$Y = 11.120879 - (0.0098462 \times X)$] is noted along with the P value ($P = .0180$) and R^2 ($R^2 = 0.384432$) value for the equation.

because of unfavorable aneurysm morphology, such as being too small to embolize, and 7 animals because of morbidities such as hairballs, a lacerated liver, a corneal ulcer, and a gastric ulcer.

The 2 categories related to procedural deaths, aneurysm creation, and embolization were divided into acute, at the time of surgery, and chronic postsurgical complications. Of the 121 deaths, 102 were procedure-related and 19 were not related to the procedure. Fifty-nine (58% of 102 procedure-related deaths) of the procedure-related deaths were associated with both acute and chronic aneurysm creation, yielding a mortality rate of 8.4% for aneurysm-creation procedures. Forty-three (42% of 102 procedure related deaths) were related to both acute and chronic aneurysm embolization, yielding a mortality rate of 8.1% for aneurysm embolization procedures.

A breakdown of all deaths by category (creation, embolization, and nonprocedural deaths) showed that the single greatest cause of death was aneurysm creation. If deaths due to aneurysm creation are plotted for 59 rabbit increments, the data show that the number of deaths due to aneurysm creation are diminished by more than half after 100 creations and remain fairly constant thereafter, regardless of the time between surgeries.

A Pearson product moment correlation (JMP, SAS) showed a significant ($P = .0246$, $R^2 = 0.354738$) negative relationship between the number of aneurysm-creation deaths and the number of creations in 50-rabbit increments. If all procedural deaths (aneurysm creations and embolizations) are correlated with the number of creations, the significant negative relationship remains ($P = .0180$, $R^2 = 0.384432$; Fig 1). When the number of days between the start of each 50-surgery increment and the finish of that increment was examined in relation to the same factors, there were no significant correlations. Finally, the factor “unknown cause of death” was the single largest contributor to the multiple regression equations for both the acute ($P = .301$, $R^2 = 0.3352$) and chronic ($P = .0279$, $R^2 = 0.3424$) aneurysm-creation equations as well as for the acute ($P = .0471$, $R^2 = 0.2896$) embolization equation. The single greatest contributor to the chronic embolization equation was the FTT variable ($P = .0175$, $R^2 = 0.3873$).

The FTT group, as defined previously, was the group that

was unable to return to normal eating habits, etc after embolization. When the average age of these animals was compared with the average age of the other animals that had embolization procedures, it was discovered that the FTT animals were significantly older (591 ± 362 days) than the other animals (159 ± 298 days, $P = .0004$).

Discussion

The rabbit model of elastase-induced aneurysm creation is being used by multiple research laboratories.^{18–22} With more laboratories trying to use the model come anecdotal reports of high mortality/morbidity rates and 1 published report²³ of a 54% death rate. This current retrospective study shows an approximately 8% death rate for both aneurysm-creation procedures and embolization procedures. Furthermore, we noted that as might be expected, mortality/morbidity rates are inversely related to the number of surgeries performed per operator. These current data suggest that the rabbit model can be applied in experienced hands with low mortality but that a learning curve is present. These data are important in guiding researchers new to the technique, regarding expected numbers of total subjects that will be needed to complete a desired sample size. The data may also inform researchers that their own mortality rates may be excessive, as noted previously, and that low rates are indeed attainable.

Multivariate analysis of variance showed that the only factor that correlated with both acute and chronic aneurysm-creation deaths was what was termed “unknown causes of death” (19 of 36 acute creation deaths and 8 of 33 chronic creation deaths). We suspect that most of these deaths were due to stroke. However, conclusive evidence for this diagnosis was not available, thus, the unknown category was created. The reason why the aneurysm-creation surgery may be more conducive to stroke can only be based on conjecture. The surgery requires a complete block of flow to the brain in the right carotid artery. Although the rabbit does have a circle of Willis, it may be that there is some biologic variability in the adequacy of the rerouted flow in some of the animals.

Acute and chronic embolization deaths accounted for 42% of the 121 total deaths. Multivariate analysis showed that the factors making an impact on acute embolization deaths were unknown causes of death (9 of 30) and also device failure (10 of 30). Factors making a significant impact on chronic embolization deaths were failure to thrive (6 of 13) and paralysis (3 of 13). Most interesting, the average time from embolism creation until sacrifice in the FTT group was significantly older ($P = .0004$) than the other animals that had embolization procedures. This indicates that the younger the animals are, the better chance they have to survive chronic embolization.

One study (Thiex et al²³) cites hemorrhagic tracheal necrosis as a leading cause of death (9 of 24 rabbits). In the Thiex study, the authors found an arterial branch coming off the CCA and providing blood to the trachea. When this branch was accommodated, the authors had no further complications due to tracheal necrosis. Although this issue was not a problem in our study, it does point out the need to be mindful of anatomic variations, not only in the creation phase but throughout any treatment phase as well.

The fact that the exact cause of death was not known in

every case was 1 limitation in this study. The other limitation was the fact that though 1 operator enabled us to provide a unique examination of changes in death rates with time, it also means that we cannot say that every operator will have diminishing mortality rates during the same time period.

Conclusions

Mortality rates of approximately 8% are associated with both experimental aneurysm creation and with embolization in the rabbit elastase-induced aneurysm model. Increasing operator experience is inversely correlated with mortality, and the age of the rabbit is positively associated with morbidity.

References

1. Reul J, Weis J, Spetzger U, et al. Long-term angiographic and histopathologic findings in experimental aneurysms of the carotid bifurcation embolized with platinum and tungsten coils. *AJNR Am J Neuroradiol* 1997;18:35–42
2. Bocher-Schwarz H, Ringel K, Bohl J, et al. Histological findings in coil-packed experimental aneurysms 3 months after embolization. *Neurosurgery* 2002;50:379–84, discussion 384–85
3. Guglielmi G, Ji C, Massoud T, et al. Experimental saccular aneurysms. II. A new model in swine. *Neuroradiology* 1994;36:547–50
4. Graves V, Partington C, Rufenacht D, et al. Treatment of carotid artery aneurysms with platinum coils: an experimental study in dogs. *AJNR Am J Neuroradiol* 1990;11:249–52
5. Graves V, Strother C, Rappe A. Treatment of experimental canine carotid aneurysms with platinum coils. *AJNR Am J Neuroradiol* 1993;14:787–93
6. Mawad M, Mawad J, Cartwright JJ, et al. Long-term histopathologic changes in canine aneurysms embolized with Guglielmi detachable coils. *AJNR Am J Neuroradiol* 1995;16:7–13
7. Tenjin H, Fushiki S, Nakahara Y, et al. Effect of Guglielmi detachable coils on experimental carotid artery aneurysms in primates. *Stroke* 1995;26:2075–80
8. Kallmes D, Helm G, Hudson S, et al. Histologic evaluation of platinum coil embolization in an aneurysm model in rabbits. *Radiology* 1999;213:217–22
9. Altes TA, Cloft HJ, Short JG, et al. 1999 ARRS Executive Council Award: creation of saccular aneurysms in the rabbit—a model suitable for testing endovascular devices. American Roentgen Ray Society *AJR Am J Roentgenol* 2000;174:349–54
10. Fujiwara N, Cloft H, Marx W, et al. Serial angiography in an elastase-induced aneurysm model in rabbits: evidence for progressive aneurysm enlargement after creation. *AJNR Am J Neuroradiol* 2001;22:698–703
11. de Gast A, Altes T, Marx W, et al. Transforming growth factor beta-coated platinum coils for endovascular treatment of aneurysms: an animal study. *Neurosurgery*. 2001;49:690–94
12. Marx W, Cloft H, Helm G, et al. Endovascular treatment of experimental aneurysms by use of biologically modified embolic devices: coil-mediated intraaneurysmal delivery of fibroblast tissue allografts. *AJNR Am J Neuroradiol* 2001;22:323–33
13. Kallmes D, Fujiwara N. New expandable hydrogel-platinum coil hybrid device for aneurysm embolization. *AJNR Am J Neuroradiol* 2002;23:1580–88
14. Kallmes D, Fujiwara N, Yuen D, et al. A collagen-based coil for embolization of saccular aneurysms in a New Zealand white rabbit model. *AJNR Am J Neuroradiol* 2003;24:591–96
15. Short J, Fujiwara N, Marx W, et al. Elastase-induced saccular aneurysms in rabbits: comparison of geometric features with those of human aneurysms. *AJNR Am J Neuroradiol* 2001;22:1833–37
16. Dai D, Ding Y, Danielson M, et al. Histopathologic and immunohistochemical comparison in human, rabbit, and swine aneurysms embolized with platinum coils. *AJNR Am J Neuroradiol* 2005;26:2560–68
17. Ding Y, Danielson M, Kadirvel R, et al. Modified technique to create morphologically reproducible elastase-induced aneurysms in rabbits. *Neuroradiology* 2006;48:528–32
18. Onizuka M, Miskolczi L, Gounis M, et al. Elastase-induced aneurysms in rabbits: effect of postconstruction geometry on final size. *AJNR Am J Neuroradiol* 2006;25:1129–31
19. Krings T, Hans F, Moller-Hartmann W, et al. Treatment of experimentally induced aneurysms with stents. *Neurosurgery* 2005;56:1347–59
20. Yasuda H, Kuroda S, Nanba R, et al. A novel coating biomaterial for intracranial aneurysms: effects and safety in extra- and intracranial carotid artery. *Neuropathology* 2005;25:66–76
21. Hans F, Moller-Hartmann W, Brunn A, et al. Treatment of wide-necked aneurysms with balloon-expandable polyurethane-covered stentgrafts: experience in an animal model. *Acta Neurochir (Wein)* 2005;147:871–76. Epub 2005 Mar 17
22. Doerfler A, Becker W, Wanke I, et al. Multimodal imaging in the elastase-induced aneurysm model in rabbits: a comparative study using serial DSA, MRA and CTA. *Rofo* 2004;176:590–96
23. Thix R, Hans F, Krings T, et al. Haemorrhagic tracheal necrosis as a lethal complication of an aneurysm model in rabbits via endoluminal incubation with elastase. *Acta Neurochir (Wien)* 2004;146:285–89, discussion 289. Epub 2004 Jan 22