

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



FRESENIUS  
KABI

caring for life

**AJNR**

*Reply:*

D.M. Mirsky, F.N. Ahmed and N.V. Stence

*AJNR Am J Neuroradiol* 2016, 37 (4) E36

doi: <https://doi.org/10.3174/ajnr.A4745>

<http://www.ajnr.org/content/37/4/E36>

This information is current as  
of April 20, 2024.

**REPLY:**

As indicated in our article,<sup>1</sup> approximately 70% of our patients with interhypothalamic adhesions had no symptoms referable to hypothalamic-pituitary dysfunction (40 of 57). We have since identified 72 additional patients, of whom 51 had no clinical symptoms. Thus, in the appropriate patient population in which associated symptoms are absent, interhypothalamic adhesions may be incidental.

Furthermore, in our experience, none of the asymptomatic patients and only 4 of the symptomatic population had additional midline abnormalities. We re-reviewed the imaging in all of our patients and found the numbers to hold true. None of the patients used for the figures had hypogenesis of the splenium or posterior periventricular white matter volume loss to cause splenium volume loss. The patient in Fig 1D did have a cavum septum pellucidum, a normal variant, but did not have partial fenestration on review of the images in 3 orthogonal planes.

Last, we agree with the Whitehead and Vezina<sup>2</sup> thesis that “the midline should be closely scrutinized for additional anomalies/abnormalities in patients harboring an interhypothalamic adhesion. Only after the brain has been carefully examined and signs/symptoms have been carefully considered can an interhypo-

thalamic adhesion be considered an incidental and isolated finding.”<sup>3</sup> It mirrors ours, “While associations between IHAs [interhypothalamic adhesions] and other syndromes likely exist, in the appropriate patient population lacking referable symptoms and with few or no other structural abnormalities, interhypothalamic adhesions may be incidental and of no clinical significance.”

**REFERENCES**

1. Ahmed FN, Stence NV, Mirsky DM. **Asymptomatic interhypothalamic adhesions in children.** *AJNR Am J Neuroradiol* 2015 Dec 3. [Epub ahead of print] CrossRef Medline
2. Whitehead MT, Vezina G. **Asymptomatic interhypothalamic adhesions in children.** *AJNR Am J Neuroradiol* 2016 Feb 25. [Epub ahead of print] CrossRef
3. Whitehead MT, Vezina G. **Interhypothalamic adhesion: a series of 13 cases.** *AJNR Am J Neuroradiol* 2014;35:2002–06 CrossRef Medline

**D.M. Mirsky**

Department of Radiology  
Children's Hospital Colorado  
Aurora, Colorado

**F.N. Ahmed**

Department of Radiology  
University of Colorado Anschutz Medical Campus  
Aurora, Colorado

**N.V. Stence**

Department of Radiology  
Children's Hospital Colorado  
Aurora, Colorado

<http://dx.doi.org/10.3174/ajnr.A4745>