

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



FRESENIUS
KABI

caring for life

AJNR

Reply:

W. Brinjikji, V. Yamaki and G. Lanzino

AJNR Am J Neuroradiol 2017, 38 (2) E17

doi: <https://doi.org/10.3174/ajnr.A5006>

<http://www.ajnr.org/content/38/2/E17>

This information is current as
of April 17, 2024.

REPLY:

Mahotra and colleagues have brought up a number of interesting questions regarding our recently published article on the neurovascular manifestations of hereditary hemorrhagic telangiectasia (HHT).¹ A number of the questions raised by the authors would require completely new studies; however, we will do our best to answer their questions in general terms.

Regarding the use of screening MR imaging for patients with HHT, a vast majority of the patients included in our study received brain MRIs as part of a standard screening protocol. As an HHT Center of Excellence, our preference is to adhere to the international guidelines for diagnosis and management of HHT.² We fully understand that there is no level 1 evidence to support the use of screening; however, screening is part of the standard of care for management of these patients.

A total of 23 patients with HHT who presented with ruptured or unruptured AVMs underwent treatment of their AVMs during this time period. A vast majority of these patients had surgery or radiosurgery. In general, outcomes related to treatment of these lesions were good, with low complication rates and no rehemorrhage. We recently published a study on the natural history of capillary vascular malformations and reported that the risk of hemorrhage from these lesions is zero.³ It is likely that the inclusion of capillary vascular malformations in natural history studies of brain AVMs in HHT is responsible for the deceptively benign natural history of these lesions compared with sporadic AVMs.⁴⁻⁶

Regarding the utility of vascular imaging in this patient subset, we certainly do not advocate the use of screening with conventional cerebral angiography for AVM diagnosis. Although cerebral angiography is a very safe procedure at our institution, it would not be cost-effective. We advocate the use of cerebral angiography if there is an equivocal finding on MR imaging or in preparation for treatment of a known brain AVM. MRA and CTA are also useful as problem-solving tools; however, based on our experience, contrast-enhanced MR imaging is generally sufficient as a screening modality for clinically significant brain AVMs. The recent addition of SWI to our HHT brain MR imaging protocol has proved useful.

A number of patients included in our study received repeat

imaging. However, this was not part of a screening protocol. In general, patients who received repeat imaging had a neurologic symptom that warranted further investigation. In general, studies have found no utility in the use of repeat screening for the presence of brain AVMs, even in children.⁷

Ultimately, Mahotra and colleagues are correct that there is no literature proving that screening or preemptive treatment of AVMs are effective strategies in HHT. It is unlikely that a definitive answer to this question will be borne from a single large institution's experience. This is a rare disease that remains poorly understood. Large, multi-institutional collaborations such as the Brain Vascular Malformation Consortium HHT Investigator Group will be needed to better answer these questions.

REFERENCES

1. Brinjikji W, Iyer VN, Yamaki V, et al. **Neurovascular manifestations of hereditary hemorrhagic telangiectasia: a consecutive series of 376 patients during 15 years.** *AJNR Am J Neuroradiol* 2016;37:1479–86 CrossRef Medline
2. Faughnan ME, Palda VA, Garcia-Tsao G, et al. **International guidelines for the diagnosis and management of hereditary haemorrhagic telangiectasia.** *J Med Genet* 2011;48:73–87 CrossRef Medline
3. Brinjikji W, Iyer VN, Lanzino G, et al. **Natural history of brain capillary vascular malformations in hereditary hemorrhagic telangiectasia patients.** *J Neurointerv Surg* 2016 Feb 26. [Epub ahead of print] CrossRef Medline
4. Kim H, Nelson J, Krings T, et al. **Hemorrhage rates from brain arteriovenous malformation in patients with hereditary hemorrhagic telangiectasia.** *Stroke* 2015;46:1362–64 CrossRef Medline
5. Brinjikji W, Iyer VN, Sorenson T, et al. **Cerebrovascular manifestations of hereditary hemorrhagic telangiectasia.** *Stroke* 2015;46:3329–37 CrossRef Medline
6. Brinjikji W, Lanzino G. **Lower risk of intracranial arteriovenous malformation hemorrhage in patients with hereditary hemorrhagic telangiectasia.** *Neurosurgery* 2016 Jul 27. [Epub ahead of print] CrossRef Medline
7. Latino GA, Al-Saleh S, Carpenter S, et al. **The diagnostic yield of rescreening for arteriovenous malformations in children with hereditary hemorrhagic telangiectasia.** *J Pediatr* 2014;165:197–99 CrossRef Medline

 W. Brinjikji

 V. Yamaki

 G. Lanzino

Department of Neurosurgery and Radiology
Mayo Clinic
Rochester, Minnesota

<http://dx.doi.org/10.3174/ajnr.A5006>