

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



FRESENIUS
KABI

caring for life

AJNR

**Comment on "Blood Flow Mimicking
Aneurysmal Wall Enhancement: A
Diagnostic Pitfall of Vessel Wall MRI Using
the Postcontrast 3D Turbo Spin-Echo MR
Imaging Sequence"**

M. Edjlali, P. Turski, C. Oppenheim and O. Naggara

This information is current as
of April 16, 2024.

AJNR Am J Neuroradiol 2018, 39 (11) E118

doi: <https://doi.org/10.3174/ajnr.A5777>

<http://www.ajnr.org/content/39/11/E118>

Comment on “Blood Flow Mimicking Aneurysmal Wall Enhancement: A Diagnostic Pitfall of Vessel Wall MRI Using the Postcontrast 3D Turbo Spin-Echo MR Imaging Sequence”

We read with interest the article “Blood Flow Mimicking Aneurysmal Wall Enhancement: A Diagnostic Pitfall of Vessel Wall MRI Using the Postcontrast 3D Turbo Spin-Echo MR Imaging Sequence”¹ published in the *American Journal of Neuroradiology* in 2018.

We would like to congratulate the authors for a very well-written article on a subject that may strongly affect the interpretation of vessel wall enhancement obtained on 3T brain MR imaging.

Several studies^{2,3} performed on various MR imaging units have demonstrated that aneurysm wall enhancement may be a marker for instability, thus a noninvasive biomarker for patient counseling and therapeutic decision-making. In this article, the authors demonstrated that part of the enhancement may be due to intra-aneurysmal slow blood flow because the wall enhancement decreased and sometimes vanished when a motion-sensitized driven equilibrium sequence was used. We agree that T1-shortened enhanced blood signal near the wall may cause nonspecific wall enhancement-like signal. However, in our opinion, the observation that slow-flow artifacts may contribute to the observed wall enhancement does not invalidate the fact that contrast enhancement is strongly linked to aneurysm status.²

Computational fluid dynamics studies⁴ demonstrated that slow flow contributes to low wall shear stress, hence promoting aneurysm wall inflammation. Slow flow, low wall shear stress, and wall inflammation probably all contribute to the observed enhancement. We agree that histopathologic confirmation is


strongly warranted to confirm the inflammatory nature of aneurysm wall enhancement.

REFERENCES

1. Kalsoum E, Chabernaud Negrier A, Tuilier T, et al. **Blood flow mimicking aneurysmal wall enhancement: a diagnostic pitfall of vessel wall MRI using the postcontrast 3D turbo spin-echo MR imaging sequence.** *AJNR Am J Neuroradiol* 2018;39:1065–67 CrossRef Medline
2. Edjlali M, Ben Hassen W, Boulouis G, et al. **Circumferential thick enhancement on high resolution vessel wall MRI is a biomarker with high specificity of intracranial aneurysm instability.** *Radiology* 2018 Jul 3:172879. [Epub ahead of print] CrossRef Medline
3. Edjlali M, Gentric JC, Régent-Rodriguez C, et al. **Does aneurysmal wall enhancement on vessel wall MRI help to distinguish stable from unstable intracranial aneurysms?** *Stroke* 2014;45:3704–06 CrossRef Medline
4. Xiang J, Tutino VM, Snyder KV, et al. **CFD: computational fluid dynamics or confounding factor dissemination? The role of hemodynamics in intracranial aneurysm rupture risk assessment.** *AJNR Am J Neuroradiol* 2014;35:1849–57 CrossRef Medline


 **M. Edjlali**

Department of Neuroradiology
Université Paris Descartes Sorbonne Paris Cité
Centre Hospitalier Sainte-Anne
Paris, France

 **P. Turski**

Departments of Radiology and Medical Physics
University of Wisconsin
Madison, Wisconsin

 **C. Oppenheim**

 **O. Naggara**

Department of Neuroradiology
Université Paris Descartes Sorbonne Paris Cité, Centre Hospitalier Sainte-Anne
Paris, France

<http://dx.doi.org/10.3174/ajnr.A5777>