

Are your **MRI contrast agents** cost-effective?

Learn more about generic **Gadolinium-Based Contrast Agents**.



FRESENIUS
KABI

caring for life

AJNR

Occlusion of cerebral arteriovenous malformations with N-butyl cyano-acrylate is permanent.

G Wikholm

AJNR Am J Neuroradiol 1995, 16 (3) 479-482

<http://www.ajnr.org/content/16/3/479>

This information is current as of April 17, 2024.

Occlusion of Cerebral Arteriovenous Malformations with *N*-Butyl Cyano-acrylate Is Permanent

Gunnar Wikholm

PURPOSE: To verify the permanence of total occlusion of cerebral arteriovenous malformations after embolization with *N*-butyl cyano-acrylate and to evaluate the occlusion rate. **METHODS:** One hundred thirty-four patients were treated for cerebral arteriovenous malformations with *N*-butyl cyano-acrylate embolization after superselective catheterization. Those initially totally occluded have been followed with angiography. **RESULTS:** In 15 instances (11.2%) the arteriovenous malformations were totally occluded by embolization alone. Follow-up angiograms have been performed in 12 cases (80%) after 4 to 78 (mean, 27) months after embolization. In all instances the arteriovenous malformations have remained occluded, and there have been no clinical signs of activity. **CONCLUSION:** After total occlusion of a cerebral arteriovenous malformation with *N*-butyl cyano-acrylate, the lesion stays occluded, and the clinical course seems stable.

Index terms: Arteriovenous malformations, embolization; Interventional materials, embolic agents

AJNR Am J Neuroradiol 16:479-482, March 1995

Several treatment options exist regarding cerebral arteriovenous malformations (AVMs) of the brain. One of these is transarterial embolization, which has now been used for more than 30 years (1), and materials and methods have developed considerably over time. One fundamental question when embolizing AVMs is whether the treatment is permanent or if the AVM will recanalize over time. This paper presents our experience in this respect regarding the embolic agent *N*-butyl cyano-acrylate.

Materials and Methods

From January 1, 1987, to December 31, 1992, 134 patients were treated at our institution with embolization for AVMs. No patient was treated outside this interval of time. The patients had mostly been deemed bad risks for surgery or not suitable for stereotactic radiosurgery because of AVM size or location or patient compliance at different neurosurgical clinics in the catchment area for interventional neuroradiology. Of these, 15 (11.2%) were totally occluded with embolization solely, without remain-

ing angiographic signs of nidus or shunt. These patients make up the study group.

The method of embolization was initially as described by Kerber (2) in 1976: the flow-guided, calibrated-leak balloon technique. It was mainly used until 1989 and occasionally later. As the variable stiffness guide wire-guided microcatheter (3) (Tracker, Target Therapeutics, Fremont, Calif) became available, it was used more from 1987 to 1990, when it was gradually replaced by the very floppy, flow-guided Pursil catheter (4) (Magic, Balt, France). Of the patients with totally occluded AVMs, *N*-Butyl cyano-acrylate (Histacryl Bleu, Melsungen AG, Germany) was used in all; polyvinyl alcohol was used as an adjunct in one. No other embolic materials were used.

The sizes of the AVMs have been measured on the films. Veins have not been included. Magnification and minification factors have always been taken in account. The AVMs have been approximated to ellipsoids and their volumes calculated using the formula:

$$V = d_1 \times d_2 \times d_3 \times \frac{1}{2},$$

where d_{1-3} are the three diameters of the AVM in centimeters, and V is the volume in milliliters (5).

Radiologic follow-up has been through cerebral angiograms performed usually at the referral hospital in the area where the patient lives. The angiograms have included all possible feeders to the area of the AVM. The criteria for total occlusion has been no remaining abnormal nidus vasculature or shunting and early-filling veins. All angiograms have been examined by us. Clinical follow-up was

Received May 4, 1994; accepted after revision August 29.

Address reprint requests to Gunnar Wikholm, MD, Section for Interventional Neuroradiology, Sahlgrenska University Hospital S-413 45 Göteborg, Sweden.

AJNR 16:479-482, Mar 1995 0195-6108/95/1603-0479
© American Society of Neuroradiology

TABLE 1: Lesion diameters and volumes in the whole group, in the group not totally occluded, and in the group totally occluded with embolization alone

| Lesion | Whole Group | Group Not Totally Occluded | Group Totally Embolized |
|--------------|------------------------------|--------------------------------|-------------------------------|
| Number | 134 | 119 | 15 |
| Diameter, mm | 41.6 ± 15 (range, 10–120) | 44.8 ± 15.3 (range, 15–120) | 23.1 ± 10.4 (range, 10–40) |
| Volume, mL | 26 ± 32 (range, 0.5–280) | 28.4 ± 33.2 (range, 1–280) | 7 ± 7 (range, 0.5–26) |

Note.—Values are mean ± SD.

performed via notes in patient files, done when follow-up angiography was performed.

Results

The AVMs totally occluded differed in size from the whole group of AVMs referred for embolization. They were smaller (Table 1). The rate of total occlusion among AVMs of different sizes is presented in Table 2.

The 15 patients in whom embolization was the only treatment have been followed up with angiography in all instances except 3. One patient with a thalamic AVM that had bled and had given her a hemiparesis had an increased paresis after embolization; this patient has been negative about all contact. The second patient has a psychiatric disease preventing angiographic follow-up, and the third patient died from pancreatic cancer 2 years after treatment. That leaves 12 (80%) cases being controlled with angiography after 4 to 78 months after embolization (mean, 27 months). In no instance was there any sign of revascularization of AVM; that is, there was absence of abnormal nidus vasculature and absence of shunting. The clinical follow-up has been for a total of 44 patient years. During this period no patient has had any bleeding or other new symptoms referable to the AVM.

Table 3 shows numbers of patients in relation to catheters used for embolization and numbers

TABLE 2: Rate of total occlusion among AVMs using 25 and 30 mm, respectively, as size limit for small AVMs

| Size Limit, mm | Total Number (%) | Number Occluded | Percentage Occluded |
|----------------|------------------|-----------------|---------------------|
| ≤25 | 14 (10.4) | 9 | 64.2 |
| >25 | 120 (89.6) | 6 | 5.0 |
| ≤30 | 26 (19.4) | 11 | 42.3 |
| >30 | 108 (80.6) | 4 | 3.7 |

TABLE 3: All patients divided into groups according to catheter used for embolization, the occlusion rate achieved in each group, and the volumes of the AVMs

| Technique | Total Number | Number Totally Occluded | Percentage Totally Occluded | Lesion Volume, mL, mean ± SD |
|-----------------------|--------------|-------------------------|-----------------------------|------------------------------|
| Tracker and/or Kerber | 64 | 11 | 17.2 | 27 ± 39 |
| Magic | 70 | 4 | 5.7 | 24 ± 22 |

Note.—The difference in proportions totally occluded is significant; $P < .02$.

and percentages of total occlusion in each group. There is an overlap between the groups, but no Magic catheter has been used in conjunction with any other type catheter in a patient with a totally occluded AVM.

Discussion

Since embolization became an alternative treatment to surgery for cerebral AVMs, there has been discussion of which therapeutic method is the safest. The “no treatment at all” option, especially for nonruptured AVMs, has its spokespeople (6). The decision to treat or not to treat depends heavily on knowledge of the natural history of this disease. Much has been written and bias in the studies discussed, but there is a rather uniform opinion that the cumulative yearly bleeding rate is around 3% and the yearly mortality from the disease is around 1% (7–9). Ondra et al (7) claim no difference in these risks regardless of presentation with hemorrhage or seizures. Brown et al (8) found approximately the same risk as cited above in his material of nonbleeders.

These studies show rather convincingly that the risk for complication because of an AVM does not depend on size or mode of presentation. Our policy has been to treat AVMs if angiography has shown appropriate paths to reach the nidus. We also considered the risk of hemorrhage to be increased if an intranidal aneurysm is present (10–12). Deruty et al found in their series that despite aneurysms in the area of the AVM, the malformation was the source of bleeding in all six studied instances (13). According to our experience, nidus aneurysms might hide in the nidus and not appear until superselective angiograms are performed or even after partial embolization. With improved catheters, it has been possible to reach the nidus in most cases referred to us. In about 10% it

has not been possible to reach a superselective position at which no functional parenchyma was involved. The 11.2% of the AVMs that have been successfully occluded by embolization alone is among the lower frequencies reported in the literature (14–17), but lesion sizes are not well reported. In our group of small AVMs (≤ 25 mm), the frequency of total occlusion is 64.2% (Table 2).

The permanence of embolization has been discussed and questioned (18). The cases reported by Rao et al (18) show that contrast in the cast has disappeared but that collaterals have developed to supply nonembolized nidus. The same phenomenon is stated by Fournier et al (19).

Our experience is that *N*-butyl cyanoacrylate is superior to polyvinyl alcohol regarding permanence. A recent paper from Nakstad et al shows interesting results with decreased recanalization rate after polyvinyl alcohol combined with platinum coils (20). This type of embolization demands Tracker or similar catheters, and in our experience it is not possible to achieve as distal catheterization with these catheters as with the Magic. Often one might therefore be dependent on the sump effect of the shunt to get the emboli to the right location. When the sump effect decreases, the risk for inadvertent embolization might increase. Development in this field is steadily ongoing, and the new hydrophilic coated Fastracker (Target) may well permit more distal catheterization. However, the possibly increased risk with use of guide wires intracranially, especially in vessels with high-flow angiopathy (21), remains with this technique.

If the rebleeding rate cited above (3%) were used on a group of 15 patients, statistically there should be a hemorrhage every second year. Because our average clinical follow-up time is 35.5 months, follow-up time is too short for definite conclusions regarding clinical cure in this material. However, the fact that none of our totally treated patients have had any sign of recanalization and no clinical sign of remaining AVM activity is a strong indicator that embolization with *N*-butyl cyanoacrylate is a permanent cure.

When splitting the material into different groups according to catheters used (Table 3), there seems to be an advantage to total occlusions when the Tracker- or Kerber-type balloon is used. If this difference is true, one reason

might be the increased ratio of *N*-butyl cyanoacrylate to blood with the Tracker (caused by larger inner lumen) or flow control with the Kerber balloons. Embolization with these catheters might allow better penetration of the nidus.

Conclusion

Arteriovenous malformations of the brain occluded with *N*-butyl cyanoacrylate remain occluded, and the clinical course after treatment seems stable.

References

1. Luessenhop AJ, Spence WT. Artificial embolization of cerebral arteries: report of use in a case of arteriovenous malformation. *JAMA* 1960;172:1153–1155
2. Kerber N. Balloon catheter with a calibrated leak. *Radiology* 1976;120:547–550
3. Kikuchi Y, Strother CM, Boyer M. New catheter for endovascular interventional procedures. *Radiology* 1987;165:870–871
4. Dion JE, Duckwiler GR, Lylyk P, Vinuela F, Bentson J. Progressive suppleness Pursil catheter: a new tool for superselective angiography and embolization. *AJNR Am J Neuroradiol* 1989;10:1068–1070
5. Pasqualin A, Barone G, Cioffi F, Rosta L, Scienza R, Da Pian R. The relevance of anatomic and hemodynamic factors to a classification of cerebral arteriovenous malformations. *Neurosurgery* 1991;28:370–379
6. Aminoff MJ. Treatment of unruptured cerebral arteriovenous malformations. *Neurology* 1987;37:815–819
7. Ondra SL, Troupp H, George ED, Schwab K. The natural history of symptomatic arteriovenous malformations of the brain: a 24-year follow-up assessment (see comments). *J Neurosurg* 1990;73:387–391
8. Brown RD Jr, Wiebers DO, Forbes G, et al. The natural history of unruptured intracranial arteriovenous malformations. *J Neurosurg* 1988;68:352–357
9. Crawford PM, West CR, Chadwick DW, Shaw MD. Arteriovenous malformations of the brain: natural history in unoperated patients. *J Neurol Neurosurg Psychiatry* 1986;49:1–10
10. Marks MP, Lane B, Steinberg GK, Snipes GJ. Intranidal aneurysms in cerebral arteriovenous malformations: evaluation and endovascular treatment. *Radiology* 1992;183:355–360
11. Muller Forell W, Valavanis A. Neuroradiologic exploration of cerebral arteriovenous malformations. *Radiologie* 1991;31:269–273
12. Marks MP, Lane B, Steinberg GK, Chang PJ. Hemorrhage in intracerebral arteriovenous malformations: angiographic determinants. *Radiology* 1990;176:807–813
13. Deruty R, Mottolese C, Soustiel JF, Pelissou Guyotat I. Association of cerebral arteriovenous malformation and cerebral aneurysm: diagnosis and management. *Acta Neurochir (Wien)* 1990;107:133–139
14. Lasjaunias P, Manelfe C, Terbrugge K, Lopez Ibor L. Endovascular treatment of cerebral arteriovenous malformations. *Neurosurg Rev* 1986;9:265–275
15. Berthelsen B, Lofgren J, Svendsen P. Embolization of cerebral arteriovenous malformations with bucrylate: experience in a first series of 29 patients. *Acta Radiol* 1990;31:13–21

16. Richling B, Bavinski G. Arterio-venous malformations of the basal ganglia: surgical versus endovascular treatment. *Acta Neurochir [Suppl] (Wien)* 1991;53:50-59
17. Ter Brugge KG, Lasjaunias P, Chiu MC. Surgical neuroangiography of intracranial vascular malformations. *Can J Neurol Sci* 1987;14:70-74
18. Rao VR, Mandalam KR, Gupta AK, Kumar S, Joseph S. Dissolution of isobutyl 2-cyanoacrylate on long-term follow-up. *AJNR Am J Neuroradiol* 1989;10:135-141
19. Fournier D, Terbrugge K, Rodesch G, Lasjaunias P. Revascularization of brain arteriovenous malformations after embolization with bucrylate. *Neuroradiology* 1990;32:497-501
20. Nakstad PH, Bakke SJ, Hald JK. Embolization of intracranial arteriovenous malformations and fistulas with polyvinyl alcohol particles and platinum fibre coils. *Neuroradiology* 1992;34:348-351
21. Pile-Spellman JMD, Baker KF, Liszczak TM, et al. High flow angiopathy: cerebral blood vessel changes in experimental chronic arteriovenous fistula. *AJNR Am J Neuroradiol* 1986;7:811-815