

Are your **MRI contrast agents** cost-effective?

Learn more about generic **Gadolinium-Based Contrast Agents**.



**FRESENIUS
KABI**

caring for life

AJNR

Comments on Brain AVM Embolization with Onyx

Huy M. Do

AJNR Am J Neuroradiol 2007, 28 (1) 178

<http://www.ajnr.org/content/28/1/178>

This information is current as of April 16, 2024.

Comments on Brain AVM Embolization with Onyx

This is an informative and encouraging single-center study on the embolization of brain arteriovenous malformations (AVMs) with Onyx, an alternative to the standard liquid embolic agent *N*-butyl cyanoacrylate. To date, there have only been 2 indexed manuscripts on brain AVM embolization with Onyx.^{1,2} The current study had a complication rate of 13.6%, with hemorrhagic complications occurring in 3 (7%) patients, resulting in the only death. AVM volume reduction from embolization in these 3 patients ranged from 50% to 95%. Although the authors discussed subtotal nidal obliteration with subsequent venous outlet obstruction as a possible cause of the hemorrhagic complications, another possible cause that has been well documented in the literature is normal perfusion pressure breakthrough bleeding.³⁻⁵ These studies highlight the physiologic derangement (changes in regional cerebral blood flow) of perinidal tissues before and after surgery and/or embolization with positron-emission tomography or single-photon emission CT, and they suggest that pretreatment nonin-

vasive evaluation of brain AVM patients may be helpful in identifying a subgroup with decreased autoregulatory capabilities who are at risk for normal perfusion pressure breakthrough after treatment. In this higher risk group, perhaps less aggressive nidal volume reduction at any single session by way of staged embolization procedures and more aggressive blood pressure management during the posttreatment period are indicated in an effort to reduce and prevent life-threatening brain hemorrhage.

References

1. van Rooij WJ, Sluzewski M, Beute GN. **Brain AVM embolization with Onyx.** *AJNR Am J Neuroradiol* 2007;28:172-177
2. Jahan R, Murayama Y, Gobin YP, et al. **Embolization of arteriovenous malformations with Onyx: clinicopathological experience in 23 patients.** *Neurosurgery* 2001;48:984-95
3. Ogasawara K, Yoshida K, Otawara Y, et al. **Cerebral blood flow imaging in arteriovenous malformation complicated by normal perfusion pressure breakthrough.** *Surg Neurol* 2001;56:380-84
4. Kaminaga T, Hayashida K, Iwama T, et al. **Hemodynamic changes around cerebral arteriovenous malformation before and after embolization measured with PET.** *J Neuroradiol* 1999;26:236-41
5. Batjer HH, Devoud MR, Meyer YJ, et al. **Cerebrovascular hemodynamics in arteriovenous malformation complicated by normal perfusion pressure breakthrough.** *Neurosurgery* 1988;22:503-09

Huy M. Do
Radiology

Stanford University Medical Center
Stanford, Calif

Address correspondence to Huy M. Do, MD, Radiology, Stanford University Medical Center, 300 Pasteur Dr, Stanford, CA 94305-5105; e-mail: huymdo@stanford.edu