

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



AJNR

Death by Nondiagnosis: Why Emergent CT Angiography Should Not Be Done for Patients with Subarachnoid Hemorrhage

David F. Kallmes, Kenneth Layton, William F. Marx and Frank Tong

This information is current as of April 19, 2024.

AJNR Am J Neuroradiol 2007, 28 (10) 1837-1838

doi: <https://doi.org/10.3174/ajnr.A0809>

<http://www.ajnr.org/content/28/10/1837>

musicians, and this increase correlated with the total duration of musical training. Although these changes may possibly be attributed to increased neuronal attenuation, increased NAA may also result from the increased size of neurons, alterations in synaptic attenuation, and changes in concentration of the other smaller metabolites under the NAA peak. In patients with impaired mathematic ability due to Gerstman syndrome or acalculia, spectroscopy demonstrated a focal decrease in NAA in the left posterior temporoparietal region,⁸ indicating an alteration of neuronal attenuation or size. A decrease in creatine suggested impaired regional cellular energetics of the neurons or a decrease in glial cells. Diffusion-tensor imaging has also been used to evaluate brain changes resulting from learning tasks. In studies evaluating the effect of practicing piano on the brain,⁹ a correlation was found between practicing and increased fiber tract organization. MR blood volume maps were also used to examine the effect of exercise on the brain. In a recent *Proceedings of the National Academy of Science U S A* publication,¹⁰ exercise-induced neurogenesis was described and found to target the dentate gyrus, a subregion that is important for memory and implicated in cognitive aging.

The exciting recent discovery that the brain can alter its shape and size during a period of weeks in response to the acquisition of new skills has led to valuable new insights on the structural response of the brain to learning and practice. New “functional” brain mapping techniques can now allow us to further evaluate morphologic brain changes in response to learning, development, disease, and recovery following injury. Future advances in imaging should also lead to a better understanding and use of the mechanisms underlying structural plasticity.

References

- Maguire EA, Spiers HJ, Good CD, et al. **Navigation expertise and the human hippocampus: a structural brain imaging analysis.** *Hippocampus* 2003;13:250–59
- Quartarone A, Siebner HR, Rothwell JC. **Task-specific hand dystonia: can too much plasticity be bad for you?** *Trends Neurosci* 2006;29:192–99
- Draganski B, Gaser C, Kempermann G, et al. **Temporal and spatial dynamics of brain structure changes during extensive learning.** *J Neurosci* 2006;26:6314–17
- Keller A, Arissian K, Asanuma H. **Synaptic proliferation in the motor cortex of adult cats after long-term thalamic stimulation.** *J Neurophysiol* 1992;68:295–308
- Darian-Smith C, Gilbert CD. **Axonal sprouting accompanies functional reorganization in adult cat striate cortex.** *Nature* 1994;368:737–40
- Levy LM, Ziemann U, Chen R, et al. **Rapid modulation of GABA in sensorimotor cortex induced by acute deafferentation.** *Ann Neurol* 2002;52:755–61
- Aydin K, Ciftci K, Terzibasoglu E, et al. **Quantitative proton MR spectroscopic findings of cortical reorganization in the auditory cortex of musicians.** *AJNR Am J Neuroradiol* 2005;26:128–36
- Levy LM, Reis IL, Grafman J. **Metabolic abnormalities detected by 1H-MRS in dyscalculia and dysgraphia.** *Neurology* 1999;53:639–41
- Bengtsson SL, Nagy Z, Skare S, et al. **Extensive piano practicing has regionally specific effects on white matter development.** *Nat Neurosci* 2005;8:1148–50. Epub 2005 Aug 7
- Pereira AC, Huddlestone DE, Brickman AM, et al. **An in vivo correlate of exercise-induced neurogenesis in the adult dentate gyrus.** *Proc Natl Acad Sci U S A* 2007;104:5638–43. Epub 2007 Mar 20

L.M. Levy

Department of Radiology

The George Washington University Medical Center

Washington, DC

DOI 10.3174/ajnr.A0724

EDITORIAL

Death by Nondiagnosis: Why Emergent CT Angiography Should Not Be Done for Patients with Subarachnoid Hemorrhage

As angiographers, we often feel stabs of inadequacy when first reading the many articles promoting the use of CT angiography (CTA) for aneurysm detection. On the basis of the apparently outstanding accuracy of the technique, many centers routinely apply CTA for evaluating patients presenting with subarachnoid hemorrhage.^{1–3} Management decisions are based on the CTA findings, with direct progression to craniotomy in some or many cases. This imaging and treatment algorithm seems efficient and state-of-the-art for everyone involved. The one individual apparently left out of consideration in this streamlined new-millennium approach is the patient, as far as we can tell.

The most urgent consideration for patients presenting with the suspicion of a ruptured aneurysm is to avoid rehemorrhage, which carries a mortality rate of 80%.⁴ Unsecured ruptured aneurysms will bleed again frequently. Rates of rehemorrhage are 4% at 24 hours, 20% at 2 weeks, and 50% at 6 months.⁵ These clinical facts mandate minimizing false-negative rates for any diagnostic examination proposed for imaging patients with suspected ruptured aneurysm. With this in mind, we read with great interest 2 abstracts presented at the 2006 annual meeting of the American Society of Neuroradiology, detailing suboptimal sensitivity of CTA for intracranial aneurysms.^{6,7} A quick back-of-the-envelope decision analysis, by using the 10% false-negative rate reported in those abstracts applied to a population with 75% prevalence of aneurysms, results in 2.5 additional deaths per 100 patients from rehemorrhage of undiagnosed ruptured aneurysms. Conventional angiography, with near-perfect accuracy and a mortality rate of 0.06%, would result in 0.06 deaths per 100 patients, or less than one fiftieth of the CTA rate.⁸ For our money, we'll take conventional angiography any day over CTA.

One could argue that sending all patients with negative findings on CTA on to conventional angiography could “fix” the deficiency of CTA caused by its imperfect sensitivity. We would welcome such a practice pattern. However, we can easily argue that all patients with positive findings on CTA should also go to conventional angiography; this recommendation would mean that every patient with a CTA, positive or negative, should go to conventional angiography.

Patients with CTA examination findings positive for aneurysms should undergo conventional angiography for several reasons. First, most centers now offer endovascular therapy as first-line treatment for ruptured aneurysms. In our opinion, it is difficult or impossible to determine, on the basis of CTA, whether an endovascular approach would be feasible in many cases, and we routinely apply 3D rotational angiography to triage patients to surgery or endovascular treatment. From a purely technical perspective, CTA has a maximum spatial resolution of between 0.35 and 0.5 mm, whereas 3D rotational angiography can resolve vessels between 0.2 and 0.3 mm and

conventional digital subtraction angiography (DSA) can resolve vessels as small as 0.1 mm (personal communication, H. Schmitt, PhD, Philips Medical Systems, May 2006). In our opinion, better spatial resolution will naturally lead to improved neck characterization. Recent literature seems to support our bias. Nearly 20% of cases initially thought to be amenable to coiling on the basis of CTA were instead referred for surgery following DSA.² Furthermore, rates of coiling from series proposing the use of CTA in place of DSA are fairly low, ranging from 20% to 35% of patients being treated with coils rather than surgery.¹⁻³ These low rates of coiling suggest that some patients sent directly to surgery without DSA may have been amenable to coil embolization.

In addition, small important arteries, such as the anterior choroidal artery or the recurrent artery of Heubner, are not identified on CTA. Finally, it remains possible that patients with 1 aneurysm detected on CTA will harbor other undetected aneurysms that were the true source of hemorrhage. Sending such patients directly to surgery, without conventional angiography, leaves these patients at risk for future hemorrhage.

Although we believe CTA is of limited value in a tertiary referral center, there may be reasonable uses for CTA in the other practice environments. For example, if one's rotational angiography equipment is suboptimal, CTA may assist in determining a working projection for coiling. Also, unstable patients with large intraparenchymal hematomas would benefit from CTA before going directly to surgery. Patients with relatively low pretest probability for aneurysm, such as those with minor trauma, or patients with relative contraindications to angiography, such as patients on warfarin (Coumadin), may reasonably be studied with CTA. In some cases, the choice of anesthesia, either neuroleptic or general, may be based on the probability of detecting an aneurysm, and antecedent CTA may be of benefit. Physicians in hospitals without endovascular therapists might be tempted to perform CTA before transferring patients to appropriate clinical centers that have endovascular capabilities, but even these patients almost certainly will undergo conventional angiography. However, we would argue that all of these instances in which CTA may be reasonable are not usually present in a tertiary care setting.

To review, in the setting of subarachnoid hemorrhage, a negative finding on CTA mandates subsequent conventional

angiography on the basis of unacceptable false-negative rates for CTA. Further, a positive finding on CTA also mandates subsequent conventional angiography to further assess aneurysm morphology for triage between endovascular therapy and surgery as well as to detect other aneurysms. If everyone with subarachnoid hemorrhage needs conventional angiography after CTA, then maybe perhaps we should dispense with the CTA all together and just get on with the appropriate test.

References

1. Karamessini MT, Kagadis GC, Petsas T, et al. **CT angiography with three-dimensional techniques for the early diagnosis of intracranial aneurysms: comparison with intra-arterial DSA and the surgical findings.** *Eur J Radiol* 2004;49:212-23
2. Hoh BL, Cheung AC, Rabinov JD, et al. **Results of a prospective protocol of computed tomographic angiography in place of catheter angiography as the only diagnostic and pretreatment planning study for cerebral aneurysms by a combined neurovascular team.** *Neurosurgery* 2004;54:1329-40
3. Dehdashti AR, Rufenacht DA, Delavelle J, et al. **Therapeutic decision and management of aneurysmal subarachnoid haemorrhage based on computed tomographic angiography.** *Br J Neurosurg* 2003;17:46-53
4. Rosenom J, Eskesen V, Schmidt K, et al. **The risk of rebleeding from ruptured intracranial aneurysms.** *J Neurosurg* 1987;67:329-32
5. Eskesen V, Rosenorn J, Schmidt K. **The impact of rebleeding on the life time probabilities of different outcomes in patients with ruptured intracranial aneurysms: a theoretical evaluation.** *Acta Neurochir (Wien)* 1988;95:99-101
6. Wang LC, Colen T, Cohen W, et al. **Evaluation of nontraumatic subarachnoid hemorrhage: CT angiography versus digital subtraction angiography.** Presented at: 44th annual meeting of the American Society of Neuroradiology, San Diego, Calif, May 1-5, 2006
7. Lane BF, Zoarski GH, Husain MA, et al. **Sensitivity of multidetector CT angiography for the detection of intracranial aneurysms.** Presented at: 44th annual meeting of the American Society of Neuroradiology, San Diego, Calif, May 1-5, 2006
8. Hankey GJ, Warlow CP, Sellar RJ. **Cerebral angiographic risk in mild cerebrovascular disease.** *Stroke* 1990;21:209-22

David F. Kallmes
Mayo Clinic
Rochester, Minn
Kenneth Layton
Baylor University Medical Center
Dallas, Tex
William F. Marx
Mission-St Joseph's Health System
Asheville, NC
Frank Tong
Emory Health System
Atlanta, Ga

DOI 10.3174/ajnr.A0809