

The **next generation** GBCA
from Guerbet is here

Explore new possibilities >

Guerbet | 

© Guerbet 2024 GUOB220151-A

AJNR

Reply:

H. Prosch, N. Grois and D. Prayer

AJNR Am J Neuroradiol 2008, 29 (4) e15

doi: <https://doi.org/10.3174/ajnr.A1050>

<http://www.ajnr.org/content/29/4/e15>

This information is current as
of March 20, 2024.

Reply:

Thank you for your interest in our article¹ and for sharing your interesting case. The proton MR spectra fit well with that of another patient with neurodegenerative Langerhans cell histiocytosis (ND-LCH) previously reported.² Decreased *N*-acetylaspartate peak accompanied by increased choline in the area of cerebellar signal intensity alterations supports the concept of a neurodegenerative process. Further studies are necessary to evaluate the sensitivity and specificity of 1H-MR spectroscopy in ND-LCH.

References

1. Prosch H, Grois N, Wnorowski M, et al. Long-term MR imaging course of neurodegenerative Langerhans cell histiocytosis. *AJNR Am J Neuroradiol* 2007;28:1022–28

2. Steiner M, Prayer D, Asenbaum S, et al. Modern imaging methods for the assessment of Langerhans' cell histiocytosis-associated neurodegenerative syndrome: case report. *J Child Neurol* 2005;20:253–57

H. Prosch

Otto Wagner Hospital

Department of Radiology

Vienna, Austria

N. Grois

Children's Cancer Research Institute

Vienna, Austria

D. Prayer

Medical University of Vienna

Division of Neuroradiology

Vienna, Austria

DOI 10.3174/ajnr.A1050