REPLY:

We greatly appreciate Dr Mascalchi's interest in our article "Inherited Cerebellar Ataxia in Childhood: A Pattern-Recognition Approach Using Brain MRI."¹ We agree that many cases of Friedreich ataxia (FRDA) have normal MR imaging findings at the time of diagnosis and that a visually normal cerebellum in a child with ataxia usually suggests FRDA. We also concur that volume loss and decreased fractional anisotropy by diffusion tensor imaging of the superior cerebellar peduncles and iron deposition on T2* are potential biomarkers for this disease.^{2,3}

REFERENCES

1. Vedolin L, Gonzalez G, Souza CF, et al. **Inherited cerebellar ataxia in childhood: a pattern-recognition approach using brain MRI.** *AJNR Am J Neuroradiol* May 17 2012 [Epub ahead of print]
2. Della Nave R, Ginestroni A, Diciotti S, et al. **Axial diffusivity is increased in the degenerating superior cerebellar peduncles of Friedreich's ataxia.** *Neuroradiology* 2011;53:367–72
3. Koeppen AH, Michael SC, Knutson MD, et al. **The dentate nucleus in Friedreich's ataxia: the role of iron-responsive proteins.** *Acta Neuropathol* 2007;114:163–73

L. Vedolin

Neuroradiology Section
Hospital Moinhos de Vento
Porto Alegre, Brazil

<http://dx.doi.org/10.3174/ajnr.A3481>