5. EITM. Empirical implications of theoretical models. <http://www.nsf.gov/sbe/ses/polisci/reports/pdf/eitmreport.pdf>. Accessed October 2, 2012
6. Clarke KA, Primo DM. Overcoming the 'physics envy.' <http://www.nytimes.com/2012/04/01/opinion/sunday/the-social-sciences-physics-envy.html>. Accessed October 2, 2012
7. Windschitl M, Thompson J, Braaten M. **Beyond the scientific method: model-based inquiry as a new paradigm of preference for school science investigations.** *Science Education* 2008;92:941–67
8. Musante S. A dynamic alternative to the scientific method: AIBS eye on education. http://www.aibs.org/eye-on-education/eye_on_education_2009_01.html. Accessed October 2, 2012
9. Understanding science. How science really works. <http://undsci.berkeley.edu>. Accessed October 2, 2012

EDITORIAL

Acute Stroke Imaging Research Roadmap II and International Survey of Acute Stroke Imaging Capabilities: We Need Your Help!

M. Wintermark and S.J. Warach,
on behalf of the STIR and Virtual International Stroke Trials
Archive (VISTA)-Imaging Investigators

Performing neuroimaging in the setting of a clinical trial, across multiple sites, is challenging because it involves standardizing acquisition and processing imaging protocols on multiple types of scanners by using multiple different platforms. The challenge is even more pronounced for cutting-edge imaging techniques such as arterial spin-labeling or diffusion tensor imaging. Mechanisms are therefore needed to translate and test advanced imaging methods across centers, to encourage the use of advanced imaging in acute settings, to stimulate closer academic-industry collaborations, and to promote the retrospective and prospective collection and pooling of imaging data while keeping in mind practical considerations such as clinical feasibility.

This daunting task has been tackled by the Stroke Imaging Research (STIR) group, a consortium of neuroradiologists, neurologists, imaging scientists, and emergency physicians with an interest in stroke imaging. STIR had a series of meetings in 2012 and 2013, where heated debates led to consensus recommendations as part of a stroke imaging research roadmap. This roadmap was published in *Stroke*¹ and should be read by all radiologists interested in stroke research because it contains some very important recommendations in terms of standardization of image acquisition and processing for stroke and how imaging should be incorporated in stroke clinical trials. To view the paper use the link in this issue's table of contents, or go directly to: <http://stroke.ahajournals.org/lookup/doi/10.1161/STROKEAHA.113.002015>.

STIR proposes a specific, standardized terminology for acute stroke imaging, aligned with the National Institute of Neurological Disorders and Stroke Common Data Elements,² including a modified TICI scale to assess reperfusion on cerebral conven-

tional angiography. STIR also introduces the concept of "Treatment-Relevant Acute Imaging Targets" (TRAIT), which is meant to capture imaging elements needed for inclusion (or exclusion) into specific treatment protocols. TRAIT acts as a shorthand term to describe the collection of specific imaging metrics used in protocols and simultaneously reminds trial designers to ensure that imaging is directed to the key anatomic or physiologic targets of their specific intervention.

STIR proposes the establishment of a calibration process for measuring ischemic core and penumbral software, as well as the population of the STIR clinical and imaging data repository to facilitate this calibration process. STIR recognizes that imaging techniques continuously evolve and that there will always be a newer, better ischemic core or penumbral imaging technique or processing software. Therefore, it is desirable to find a balance between continued attempts to improve on existing methods versus determining whether existing methods are good enough to be used in current clinical trials. At this time, STIR does not assess or recommend how to use ischemic core and penumbral information for prognosis, prediction of response to treatment, and/or selection of patients for reperfusion therapy. These are better answered in well-designed clinical trials or prospective validation studies.

Finally, STIR recommends the creation of a stroke neuroimaging network involving a collaboration between sites to promote scientific collaboration and education in a distributed fashion and further advance imaging protocols and software reuse, and data and model sharing. As a first step towards the creation of this network, STIR is conducting an international survey for which we need your help. Please take 15 minutes to fill out the survey, which can be found at <https://www.surveymonkey.com/s/DQRDYB2>. Thank you in advance for your collaboration!

REFERENCES

1. Wintermark M, Albers GW, Broderick JP, et al. **Acute Stroke Imaging Research Roadmap II.** *Stroke* 2013 Jul 16 [Epub ahead of print]
2. http://www.ninds.nih.gov/research/clinical_research/toolkit/common_data_elements.htm Accessed July 1 2013

EDITORIAL

Mechanical Thrombectomy after IMS III, Synthesis, and MR-RESCUE

L. Pierot, J. Gralla, C. Cognard, and P. White

Three recent publications report the neutral results of 3 randomized studies (Synthesis Expansion, Interventional Management of Stroke [IMS] III, and Mechanical Retrieval and Recanalization of Stroke Clots Using Embolectomy [MR-RESCUE]) comparing IV thrombolysis therapy with the endovascular treatment (EVT) of acute ischemic stroke (AIS).^{1–3} The simultaneous publication of these 3 reports might lead to the erroneous conclusion that endovascular treatment has no place in the management of AIS. However, the role of endovascular therapy for the treatment must be more carefully considered, given the tremendous