REPLY:

We would like to thank Dr Labauge and colleagues for their thoughtful reading of our recent article “Brain Magnetic Susceptibility Changes in Patients with Natalizumab-Associated Progressive Multifocal Leukoencephalopathy” (NTZ-PML).¹

For the survival and functional outcome of patients treated by NTZ, it is crucial to detect PML on the basis of brain MR imaging before symptom onset, when brain involvement is more localized. This is the first clinical report showing the magnetic susceptibility changes in brain NTZ-PML lesions from the asymptomatic to the chronic stages with different MR imaging scanners (1.5T and 3T) and different MR images (T2* and SWI). In this preliminary study, we did not assess the diagnostic accuracy of the hypointense rim sign on susceptibility-weighted MR images for the diagnosis of asymptomatic NTZ-PML, but our data suggest the major role of this technique in clinical routine. Thus, as suggested by Dr Labauge and colleagues, we believe that susceptibility-weighted MR images should be systematically included in the MR imaging protocol performed in patients with suspected PML.

SWI is now available on most MR imaging scanners and should be preferred to T2* due to the higher spatial resolution and the improved lesion conspicuity inherent to this technique. In our experience, SWI was much more sensitive than T2* in detecting a subtle hypointense rim involving the U-fibers, particularly at the asymptomatic stage. Assessing iron accumulation by quantitative susceptibility mapping must be useful as well to better understand the pathophysiology of this new imaging feature, as previously suggested.² However, this approach still needs postprocessing time and cannot be performed in daily clinical practice.

SWI should be interpreted according to other signal anomalies previously reported in the literature for the early diagnosis of PML, including perivascular punctate lesions (enhanced or not on postcontrast T1WI) and hyperintensities on diffusion-weighted images.^{3,4} Awareness of these specific lesion patterns and the use of an adequate MR imaging protocol facilitate an earlier diagnosis of asymptomatic NTZ-PML, associated with a more favorable prognosis.

REFERENCES

1. Hodel J, Outteryck O, Verclytte S, et al. **Brain magnetic susceptibility changes in patients with natalizumab-associated progressive mul-**

- tifocal leukoencephalopathy.** *AJNR Am J Neuroradiol* 2015 Aug 27. [Epub ahead of print] [CrossRef Medline](#)
2. Carra-Dalliere C, Menjot de Champfleury N, Aygnac X, et al. **Quantitative susceptibility mapping suggests a paramagnetic effect in PML.** *Neurology* 2015;84:1501–02 [CrossRef Medline](#)
3. Yousry TA, Pelletier D, Cadavid D, et al. **Magnetic resonance imaging pattern in natalizumab-associated progressive multifocal leukoencephalopathy.** *Ann Neurol* 2012;72:779–87 [CrossRef Medline](#)
4. Hodel J, Outteryck O, Dubron C, et al. **Asymptomatic progressive multifocal leukoencephalopathy associated with natalizumab: diagnostic precision with MR imaging.** *Radiology* 2015 Oct 5. [Epub ahead of print] [CrossRef Medline](#)

J. Hodel

University of Lille

Lille, France

Department of Neuroradiology

Roger Salengro Hospital

Lille, France

O. Outteryck

University of Lille

Lille, France

Department of Neurology

Roger Salengro Hospital

Lille, France

S. Verclytte

Department of Radiology

Saint Philibert Hospital

Lille, France

V. Deramecourt

University of Lille

Lille, France

Department of Pathology

Lille University Hospital

Lille, France

A. Lacour

University of Lille

Lille, France

Department of Neurology

Roger Salengro Hospital

Lille, France

J.-P. Pruvo

University of Lille

Lille, France

Department of Neuroradiology

Roger Salengro Hospital

Lille, France

P. Vermersch

University of Lille

Lille, France

Department of Neurology

Roger Salengro Hospital

Lille, France

X. Leclerc

University of Lille

Lille, France

Department of Neuroradiology

Roger Salengro Hospital

Lille, France

<http://dx.doi.org/10.3174/ajnr.A4634>