Cortical Thickness of Native Tibetans in the Qinghai-Tibetan Plateau

W. Wei, X. Wang, Q. Gong, M. Fan, and J. Zhang

ABSTRACT

BACKGROUND AND PURPOSE: High-altitude environmental factors and genetic variants together could have exerted their effects on the human brain. The present study was designed to investigate the cerebral morphology in high-altitude native Tibetans.

MATERIALS AND METHODS: T1-weighted brain images were obtained from 77 Tibetan adolescents on the Qinghai-Tibetan Plateau (altitude, 2300–5300 m) and 80 matched Han controls living at sea level. Cortical thickness, curvature, and sulcus were analyzed by using FreeSurfer.

RESULTS: Cortical thickness was significantly decreased in the left posterior cingulate cortex, lingual gyrus, superior parietal cortex, precuneus, and rostral middle frontal cortex and the right medial orbitofrontal cortex, lateral occipital cortex, precuneus, and paracentral lobule. Curvature was significantly decreased in the left superior parietal cortex and right superior marginal gyrus; the depth of the sulcus was significantly increased in the left inferior temporal gyrus and significantly decreased in the right superior marginal gyrus, superior temporal gyrus, and insular cortex. Moreover, cortical thickness was negatively correlated with altitude in the left superior and middle temporal gyri, rostral middle frontal cortex, insular cortex, posterior cingulate cortex, precuneus, lingual gyrus, and the right superior temporal gyrus. Curvature was positively correlated with altitude in the left rostral middle frontal cortex, insular cortex, and middle temporal gyrus. The depth of the sulcus was negatively correlated with altitude in the left lingual gyrus and right medial orbitofrontal cortex.

CONCLUSIONS: Differences in cortical morphometry in native Tibetans may reflect adaptations related to high altitude.

ABBREVIATIONS: DMN = default mode network; HA = high altitude; PCC = posterior cingulate cortex; PPARA = peroxisome proliferator-activated receptor α

The Qinghai-Tibetan Plateau is one of the most extreme environments inhabited by humans. High-altitude (HA) natives have developed distinctive biologic characteristics in respiratory and circulatory functions,^{1,2} hemoglobin concentration,³ arterial oxygen saturation,⁴ energy metabolism,⁵ and genes⁶ to offset the stresses of HA environmental factors. Previous studies have re-

vealed the adaptation of the cerebral glucose metabolic rate⁷ in HA Quechua natives and the adaptation of cerebral autoregulation in HA Himalayan natives⁸ and Ethiopian natives.⁹ However, existing studies provided no assessments of the cerebral structures in HA natives.

The brain is one of the organs in the body with the highest oxygen consumption and thus is particularly vulnerable to hypoxia.¹⁰ In contrast, cold weather at a HA may result in hypothermia, which has been shown to have a neuroprotective effect against hypoxic damage.^{11,12} Hypobaric at a HA could also be a factor contributing to brain lesions because it has been proved to induce white matter and cortical thickness impairment.^{13–15} Environmental ultraviolet rays can cause neuronal damage in the visual system and brain.^{16–18} Therefore, these HA environmental factors together could have exerted their effect on brain development.

Many present day native Tibetans on the Qinghai-Tibetan Plateau are the descendants of colonizers who arrived, at most, 25,000 years ago.¹⁹ Genetic adaptations at HA have been found in

Received August 2, 2016; accepted after revision October 24.

From the MRI Center (W.W.), First Affiliated Hospital of Xiamen University, Xiamen, China; Institute of Brain Disease and Cognition (W.W., J.Z.), Medical College of Xiamen University, Xiamen, China; Department of Neurology (X.W.), Affiliated Hospital of Hangzhou Normal University, Hangzhou, China; Huaxi Magnetic Resonance Research Center (Q.G.), West China Hospital, Sichuan University, Chengdu, China; and Department of Cognitive Sciences (M.F.), Institute of Basic Medical Sciences, Beijing, China.

This work was supported by National Science Foundation of China (grant numbers: 81171324 and 81471630) and the National Key Project (2012CB518200).

Please address correspondence to Jiaxing Zhang, MD, Department of Physiology, Medical College of Xiamen University, Xiamen, 361005, China; e-mail: zhangjiaxing@xmu.edu.cn

Indicates open access to non-subscribers at www.ajnr.org

<http://dx.doi.org/10.3174/ajnr.A5050>

the native Tibetans. Unique to Tibetans are the variants of genes within the hypoxia-inducible factor pathway, including Egl 9 homolog 1 (*EGLN1*), endothelial PAS domain protein 1 (*EPAS1*), and peroxisome proliferator-activated receptor α (*PPARA*), key genes in the oxygen homeostasis system at all altitudes.²⁰ Among these genes, *PPARA* is involved in neuronal proliferation, differentiation, and apoptosis²¹ and neuroprotection against oxidative stress.²² Genetic variants may also play a role in brain development at a HA.

In previous studies, we have investigated the brain acclimatization to HA in the Han immigrant descendants on the Qinghai-Tibetan Plateau²³ and in the lowlanders immigrating to the Qinghai-Tibetan Plateau for several days to several years.^{24,25} We found that HA acclimatization was associated with brain structural and functional modifications. Therefore, we hypothesized that native Tibetans could have developed brain adaptations to HA under the combined actions of both environmental factors and genetic variants. To test this hypothesis, we investigated the cerebral morphology of 77 healthy Tibetan adolescents living at different altitudes, with 80 sea-level Han adolescents as controls.

MATERIALS AND METHODS

Subjects

Seventy-seven healthy native Tibetans (34 males and 43 females, 14–18 years of age) living in the southeast Tibet Autonomous Region (southeast Nagqu, Xigaze, Lasa, Nyingchi, and Chamdo) (altitude, 2300–5300 m) were recruited. Their ancestors were all native Tibetans living on the Qinghai-Tibetan Plateau. They were without any prior descent to the lowlands or ascent to higher altitudes. All had a normal body mass index. They were all 10th grade students and had been enrolled in a high school at Chengdu (altitude, <400 m) for half a month. The control subjects were 80 (34 males and 46 females, 14–18 years of age) adolescents living at sea level (altitude, <50 m), matched with the native Tibetans by sex, age, and education. Subjects were excluded if they had a history of mountain sickness, neurologic disorder, or head injury. The experimental protocol was approved by the Research Ethics Review Board of Xiamen University. Procedures were fully explained, and all subjects provided written informed consent before participating in the study.

MR Imaging Data Acquisition

Brain images were obtained on the same model 3T Signa Excite MR imaging system (GE Healthcare, Milwaukee, Wisconsin) at the Huaxi Magnetic Resonance Research Center, West China Hospital, Chengdu, China, and at the MR Imaging Center, First Affiliated Hospital of Xiamen University, Xiamen, China. Parameters for the T1-weighted MPRAGE sequences were the following: TR/TE = 1900/2.48 ms, FOV = 25 × 25 cm², NEX = 1, matrix = 512 × 256, section thickness = 1.0 mm. Conventional 2D T1 and T2 images were also obtained and examined for any incidental findings. Data analyses were conducted by 2 researchers who were blinded to the status of subjects.

FreeSurfer Analysis

FreeSurfer (Version 5.1.0; <http://surfer.nmr.mgh.harvard.edu>) was used for cortical thickness, curvature, and sulcus analyses.

The process consisted of the removal of nonbrain tissue, mapping to Talairach-like space, and segmentation of the gray/white matter and pial boundaries. These maps of measurements were obtained by reconstructing representations of the GM/white matter boundary and the white matter boundary to the GM/CSF boundary and then calculating the closest distance from those surfaces at each vertex on the tessellated surfaces. All subject data were resampled to the FreeSurfer default common surface template by using a high-resolution surface-based averaging technique that aligned cortical folding patterns. Finally, the surface data were spatially smoothed by using a Gaussian kernel of 10-mm full width at half maximum. Regional variations of the cortical thickness, curvature, and sulcus were compared by using an independent samples *t* test, with age, sex, and education as covariates. The statistical parametric map was generated at $P < .05$ (false discovery rate–corrected for multiple comparisons).

Correlation Analysis

The correlations of cerebral structural measurements with altitude were analyzed, with age and sex as covariates. The false discovery rate of $P < .05$ was applied to correct for multiple comparisons.

RESULTS

Cortical Thickness, Curvature, and Depth of the Sulcus

Compared with controls, the significant changes of cerebral structural measurements in native Tibetans are shown in Fig 1. Cortical thickness was significantly decreased in the left posterior cingulate cortex (PCC), lingual gyrus, superior parietal cortex, precuneus, and rostral middle frontal cortex and the right medial orbitofrontal cortex, lateral occipital cortex, precuneus, and paracentral lobule (Table 1). Furthermore, curvature was significantly decreased in the left superior parietal cortex and right superior marginal gyrus. Finally, the depth of the sulcus was significantly increased in the left inferior temporal gyrus and significantly decreased in the right superior marginal gyrus, superior temporal gyrus, and insular cortex.

Correlations

The significant correlations of cerebral structural measurements with altitude in native Tibetans are shown in Fig 2. Cortical thickness was negatively correlated with altitude in the left superior and middle temporal gyri, rostral middle frontal cortex, PCC, precuneus, lingual gyrus, insular cortex, and right superior temporal gyrus (Table 2). Curvature was positively correlated with altitude in the left rostral middle frontal cortex, insular cortex, and middle temporal gyrus. The depth of the sulcus was negatively correlated with altitude in the left lingual gyrus and right medial orbitofrontal cortex.

Figure 3 indicates the overlaps of brain between the regions showing decreases of cortical thickness and those showing correlation between cortical thickness and altitude. These regions included the left PCC, precuneus, lingual gyrus, and rostral middle frontal cortex.

DISCUSSION

Our study is the first to reveal the cerebral characteristics of native Tibetans living on the Qinghai-Tibetan Plateau. Adaptations of Tibetans to HA were associated with structural modifications in

FIG 1. Regions showing changes of cortical measurements in native Tibetans compared with the sea-level Han controls ($P < .05$, false discovery rate–corrected for multiple comparisons). Blue indicates a decrease; red, an increase.

Table 1: Regional information of changed cortical thickness in native Tibetans

Areas	Volume (mm ³)	MNI Coordinate			P (Peak)
		x	y	z	
Left					
Posterior cingulate cortex/lingual gyrus	1467	−3.7	−8.5	33.8	3.651
Superior parietal cortex/precuneus	1815	−8.6	−68.1	56.0	2.571
Rostral middle frontal cortex	1575	−31.2	36.9	22.6	1.878
Right					
Lateral occipital cortex	2149	26.4	−83.9	14.4	3.081
Precuneus/paracentral lobule	2200	7.7	−61.8	58.2	2.583
Medial orbitofrontal cortex	1366	8.4	47.6	−13.8	3.296

Note:—MNI indicates Montreal Neurological Institute.

cortical thicknesses, curvature, and the sulcus. Mainly in the left hemisphere, the cortical structures were significantly correlated with altitude; moreover, with increasing altitude, native Tibetans had a decrease of regional cortical thickness, while they had an increase of cortical folding in terms of curvature. The regions showing decreases of cortical thickness and those showing correlations between cortical thickness and altitude overlapped in the left PCC, precuneus, lingual gyrus, and rostral middle frontal cortex.

The changed brain regions in the native Tibetans were in agreement with those found in our previous studies on Han immigrants living on the Qinghai-Tibetan Plateau. For example, Han immigrant descendants at HA showed decreases of GM volume in the cingulate cortex, rostral middle frontal cortex, lingual gyrus, and anterior insular cortex.²³ Adult immigrants who lived at a HA for 2 years showed decreases of cortical thickness in the

paracentral lobule, PCC, and precuneus and increases of cortical thickness in the orbitofrontal cortex, superior parietal cortex, inferior temporal gyrus, and lingual gyrus.²⁵ Moreover, compared with Tibetan adolescents who had lived at sea level for 4 years, the Tibetan adolescents who lived at a HA all the time showed decreases of GM volume in the superior parietal cortex and insular cortex.²⁶

Effects of Environmental Factors on Brain Structural Development

In native Tibetans, there were significant correlations between regional cortical measurements and altitude, suggesting that HA environmental factors influence brain development. The effects of HA environmental factors on brain development have been detected in Tibetan adolescents who were born and raised on the Qinghai-Tibetan Plateau compared with Tibetan adolescents who had lived at sea level during the past 4 years.²⁶

Cortical changes may be induced by hypoxia. For example, cerebral edema was found after acute exposure to normobaric hypoxia.²⁷ In agreement with the present findings, damage of GM in the cingulate cortex, temporal gyrus, and insular cortex has been detected in patients with hypoxia with chronic obstructive pulmonary disease.²⁸ The changed cerebral circulation may be attributed to temporal and posterior brain structural alterations shown in the present study. The vertebral artery supplies blood to

FIG 2. Regions in which the cortical measurements in native Tibetans showed significant correlations with altitude ($P < .05$, false discovery rate–corrected for multiple comparisons). Blue indicates a negative correlation; red, a positive correlation.

Table 2: Information of regions in which cortical thickness showed correlation with altitude in Tibetans

Areas	Volume (mm ²)	MNI Coordinate			P (Peak)
		x	y	z	
Left					
Superior/middle temporal gyrus	3150	−49.0	−32.6	−9.2	4.949
Precuneus/posterior cingulate cortex/lingual gyrus	2510	−9.4	−55.9	22.5	3.519
Rostral middle frontal cortex	1431	−23.8	51.8	−4.9	3.426
Insular cortex	2543	−27.8	24.3	5.7	2.445
Right					
Superior temporal gyrus	1174	56.6	−13.1	−4.7	2.134

Note:—MNI indicates Montreal Neurological Institute.

these 2 parts of the brain. Willie et al²⁹ found that with hypoxia, the relative increase in vertebral artery blood flow was 50% greater than in the other vessels. Ogoh et al³⁰ found that the internal carotid artery blood flow was unchanged, while the vertebral artery blood flow was significantly increased during acute hypoxia. Feddersen et al³¹ found that in mountaineers during ascent to 5050 m, the cerebral blood flow velocity was decreased in the posterior cerebral arteries. Moreover, hypoxia-induced alteration of cerebral blood flow patterns was found in the posterior brain in patients with hypoxia and obstructive sleep apnea, which included the lingual gyrus, parietal lobule, precuneus, temporal cortex, and insular cortex.³² Recently, Lawley et al³³ found that 2 hours of hypoxia caused an expected increase in frontal cortical gray matter perfusion but unexpected perfusion decreases predominantly in the PCC. After 10 hours in hypoxia, the decreased blood flow became more pronounced and widespread. They fur-

ther confirmed that these reductions in cerebral blood flow from hypoxia were related to vasoconstrictions.

Hypobaria at HA could be another important factor contributing to brain alterations. Acute mountain sickness scores were higher,³⁴ and visual sensitivities were lower³⁵ in hypobaric than in normobaric hypoxia. Studies on U-2 pilots showed that hypobaria was associated with a change of white matter hyperintensity^{14,36} and with the reduction of cortical thickness.¹⁵ Moreover, expo-

sure to hypobaria after traumatic brain injury worsened brain function.³⁷

Cold ambient temperature at a HA may result in hypothermia. Mild hypothermia is associated with decreased ventricular function, oxygen extraction, and microvascular flow.³⁸ During hypoxia, hypothermia has been shown to have a neuroprotective effect against hypoxic encephalopathy.¹² Moreover, hypothermia treatment in an animal stroke model reduced infarct size and improved neurologic behavior.¹¹

The decrease of visual cortical thickness in the present study may be an outcome of ultraviolet radiation at HA. Ultraviolet light has been shown to alter neuronal activity in the cortical structures involved in visual processing.³⁹ Environmental doses of ultraviolet rays can induce apoptosis in the retina and lamina ganglionaris within the visual system and in the brains of invertebrate crustaceans.^{16–18}

FIG 3. Tibetan brain overlaps (blue areas) between the regions showing significant decreases of cortical thickness (yellow line) and those showing correlations between cortical thickness and altitude (red line). For each correlation diagram, 80 overlapped red dots indicate the cortical thickness values in the sea-level controls. Blue dots indicate that the cortical thickness values in 77 Tibetans were significantly correlated with altitude.

Effects of Genes on Brain Structural Development

The results of the present study obtained from native Tibetans showed some differences from those found in the Han immigrant descendants,²³ which suggest that genes may play a role in Tibetan brain development. A recent study discovered genetic selection on the *PPARA* in HA native Tibetans on the Qinghai-Tibetan Plateau through comparing them with Chinese and Japanese lowland populations.²⁰ *PPARA* is expressed in the brain.^{21,40} It is involved in neuronal proliferation, differentiation, and apoptosis.^{21,41} *PPARA* encodes the fatty acid-activated transcription factors and fatty acid oxidation regulators; thus, it may affect brain structure through control of energy homeostasis.^{42,43} *PPARA* is also involved in angiogenesis,⁴⁴ which may contribute to the GM

change. Moreover, during hypoxia, *PPARA* supports a role in neuroprotection against oxidative stress.²²

Gene or Environment: Which One Determines Brain Structure?

Reduced psychomotor speed has been tested in the subjects (6–16 years of age) of mixed-ethnic background (including Native American, European, or African ancestries who immigrated to HA during different periods) who were born and raised at HAs.⁴⁵ For the proportion of Native Americans, Europeans, and Africans, genetic admixture with reduced psychomotor speed was comparable across altitude groups; therefore, chronic hypoxic exposure rather than genetic inheritance appears to affect brain

function development at HA. However, pulmonary function (forced expiratory volume and forced vital capacity) measures were greater in the HA natives compared with the lowland natives born and raised at HAs; this finding suggests a genetic effect.⁴⁶ In contrast, these 2 pulmonary measures were similar in HA natives and sea-level natives at lowland; this finding suggests that the genetic potential for larger lung volumes at HA depended on the developmental exposure to HA. In summary, results from these studies emphasized the importance of developmental adaptations to HA.

Implications for the Changes in Cortical Structural Measurements

In our study, decreases of cortical thickness in the medial orbito-frontal cortex, PCC, lingual gyrus, and precuneus; a decrease of curvature in the superior parietal cortex; and a decrease of the depth of the sulcus in the temporal cortices were detected. These regions are core areas of the resting-state brain default mode network (DMN) in humans.^{47,48} The impairment of GM structures may contribute to reduced neuronal activity. Previous multimodal neuroimaging studies have shed light on this function-structure association, underlining cognition, aging, disease, and behavior.⁴⁹ The changed activity within the DMN in response to chronic stress exposure has been found in humans⁵⁰ and animals.⁵¹ The impairment of cortical structures in HA Tibetans is likely to reflect the stress-induced atrophic effects in cortical regions. Cortical atrophy within the DMN may reduce the brain response to HA environmental stress. Because regions within the DMN exert forebrain modulation over visceral responses mediated via the hypothalamus and brain stem, structural and functional changes within these circuits may contribute to modulations in autonomic and neuroendocrine functions associated with HA adaptations.

Tibetan college students were found to have higher levels of negative emotional expressivities such as fear, sadness, and anger in comparison with Han, Uighur, Hui, and Mongolian students.⁵² The Cattell 16 Personality Factor Questionnaire revealed that Tibetan adolescents had lower stability, persistence, independence, and self-discipline and higher extroversion and excitability than Han adolescents.^{53,54} The findings of the Eysenck Personality Questionnaire and Trait Coping Style Questionnaire suggested that Tibetan adolescents had higher neuroticism, introversion, and psychoticism and lower concealment than Han adolescents.⁵⁵ In humans, the DMN is thought to support a variety of self-referential functions and mind-wandering.⁵⁶ Anxiety disorders and high trait anxiety are associated with decreased functioning of the DMN.⁵⁷ The precuneus is implicated in social and emotional functions, especially among individuals with neuroticism.⁵⁸ The left posterior and middle cingulate cortices and precuneus were found to be negatively correlated with neuroticism.⁵⁹ The ventral medial prefrontal (orbitofrontal) cortex of the DMN functions as a sensory-visceromotor link concerned with social behavior, mood control, and motivational drive, all of which are important components of an individual's personality.⁴⁸ Therefore, a changed DMN may explain why Tibetan adolescents had personality traits distinct from those of the Han adolescents.

The decreases of cortical thickness in the PCC and lingual

gyrus may be related to visual deficits at HA. The poor visual performances were consistent and prominent problems in the HA immigrants, such as the sea-level college students after 7 months of study at moderate altitudes,⁶⁰ and in the HA Han immigrant descendants.⁶¹ The PCC and lingual gyrus have direct anatomic connectivities with the visual striate cortices.⁶² The PCC lies at the parietomedial temporal visual stream pathway.⁶³ It is activated by visual perception, attention, and motion.^{64–66} The lesions of the PCC produce an impairment in visual discrimination learning⁶⁷ and a decrease of resting-state functional connectivity in the PCC, which is indicated by severe myopia.⁶⁸ The lingual gyrus is a part of the occipitotemporal pathway that engages in object drawing⁶⁹ and is activated by visuospatial navigation⁷⁰ and angle discrimination tasks.⁷¹ GM volume in the right lingual cortex has been proved to have a positive correlation with visual reproduction,⁷² while GM atrophy in this area was associated with visual hallucinations.⁷³

The decreased depth of the anterior insular sulcus and damage to the ventral medial prefrontal cortex in the present study support the hypothesis that these 2 regions are needed to process the perturbation of the homeostatic balance in extreme environments.^{48,74} Because the lesions in the insular cortex have been proved to disrupt representation of internal states that underpin motivation,⁷⁵ the changes of the anterior insular cortex may also be related to the blunted hypoxic ventilatory response in HA residents.⁷⁶

Limitations

There are several limitations to our study. First, brain images were obtained by using 2 MR imaging scanners. However, several studies have confirmed that the MR imaging scanner field strength, manufacturer, machine upgrade, and pulse sequence⁷⁷ and the different postprocessing algorithms applied to the images acquired from different MR imaging systems⁷⁸ have little effect on the reliability of cortical thickness measurements.

Second, the Tibetan subjects had moved to the lowlands for 1–15 days, which would have an influence on cerebral blood flow. However, given the previous investigations, this effect may be slight after residing at sea level for such a short time. For example, when HA adult natives descended to sea level for several weeks, their cardiac functions remained unchanged.^{79,80}

Third, the interethnic difference of brain structure can be large. Brain morphologic differences between populations of different origins have been found in whole-brain and region-specific volume.⁸¹ Future study should compare the interethnic difference of brain structures between Tibetans and Han populations at lowland.

Fourth, Tibetans have been living on the Qinghai-Tibetan Plateau and have developed a unique culture and language. The lifestyle of the Tibetans was different from that in the lowlands, for example, in the absence of crowds, which could have affected the subjects' emotional well-being. Moreover, diet was likely to be an important factor for the observed brain changes.

CONCLUSIONS

Native Tibetans have developed brain structural adaptations that enable their successful existence in the HA environment. The al-

terations of cortical structure in native Tibetans were mainly located in the resting-state brain DMN and were in agreement with those found in the Han immigrants on the Qinghai-Tibetan Plateau. Brain structural modifications may be derived from the interaction between genetic and environmental factors. Future study is need to test the personality traits, body homeostasis, and cognition and clarify the neuronal mechanisms of the Tibetans.

Disclosures: Ming Fan—RELATED: Grant: National Key Project, Comments: grant number 2012CB518200. Jiaxing Zhang—RELATED: Grant: National Science Foundation of China and National Key Project, Comments: grant Nos. 81171324 and 81471630.

REFERENCES

- Houston CS, Riley RL. **Respiratory and circulatory changes during acclimatization to high altitude.** *Am J Physiol* 1947;149:565–88 [CrossRef Medline](#)
- Lahiri S, DeLaney RG, Brody JS, et al. **Relative role of environmental and genetic factors in respiratory adaptation to high altitude.** *Nature* 1976;261:133–35 [CrossRef Medline](#)
- Beall CM, Reichsman AB. **Hemoglobin levels in a Himalayan high altitude population.** *Am J Phys Anthropol* 1984;63:301–06 [CrossRef Medline](#)
- Niermeyer S, Yang P, Shanmina, et al. **Arterial oxygen saturation in Tibetan and Han infants born in Lhasa, Tibet.** *N Engl J Med* 1995;333:1248–52 [CrossRef Medline](#)
- Murray AJ. **Energy metabolism and the high-altitude environment.** *Exp Physiol* 2016;101:23–27 [CrossRef Medline](#)
- Yi X, Liang Y, Huerta-Sanchez E, et al. **Sequencing of 50 human exomes reveals adaptation to high altitude.** *Science* 2010;329:75–78 [CrossRef Medline](#)
- Hochachka PW, Clark CM, Brown WD, et al. **The brain at high altitude: hypometabolism as a defense against chronic hypoxia?** *J Cereb Blood Flow Metab* 1994;14:671–79 [CrossRef Medline](#)
- Jansen GF, Krins A, Basnyat B, et al. **Role of the altitude level on cerebral autoregulation in man resident at high altitude.** *J Appl Physiol* (1985) 2007;103:518–23 [Medline](#)
- Claydon VE, Gulli G, Slessarev M, et al. **Cerebrovascular responses to hypoxia and hypocapnia in Ethiopian high altitude dwellers.** *Stroke* 2008;39:336–42 [CrossRef Medline](#)
- Hadanny A, Efrati S. **Oxygen: a limiting factor for brain recovery.** *Crit Care* 2015;19:307 [CrossRef Medline](#)
- Gao XY, Huang JO, Hu YF, et al. **Combination of mild hypothermia with neuroprotectants has greater neuroprotective effects during oxygen-glucose deprivation and reoxygenation-mediated neuronal injury.** *Sci Rep* 2014;4:7091 [CrossRef Medline](#)
- Kvistad CE, Thomassen L, Waje-Andreassen U, et al. **Body temperature and major neurological improvement in tPA-treated stroke patients.** *Acta Neurol Scand* 2014;129:325–29 [Medline](#)
- Coppel J, Hennis P, Gilbert-Kawai E, et al. **The physiological effects of hypobaric hypoxia versus normobaric hypoxia: a systematic review of crossover trials.** *Extrem Physiol Med* 2015;4:2 [CrossRef Medline](#)
- McGuire SA, Sherman PM, Wijtenburg SA, et al. **White matter hyperintensities and hypobaric exposure.** *Ann Neurol* 2014;76:719–26 [CrossRef Medline](#)
- Simmons D, Bursaw A, Sherman P, et al. **Repeated exposure to hypobaria in U2 pilots is associated with reduced aggregate cortical thickness.** *Neurology* 2015;84(suppl):P4.161
- Hollmann G, Linden R, Giangrande A, et al. **Increased p53 and decreased p21 accompany apoptosis induced by ultraviolet radiation in the nervous system of a crustacean.** *Aquatic Toxicol* 2016;173:1–8 [CrossRef Medline](#)
- Miguel NC, Meyer-Rochow VB, Allodi S. **Ultrastructural study of first and second order neurons in the visual system of the crab *Ucides cordatus* following exposure to ultraviolet radiation.** *Micron* 2002;33:627–37 [Medline](#)
- Miguel NC, Wajsenzon IJ, Takiya CM, et al. **Catalase: Bax and p53 expression in the visual system of the crab *Ucides cordatus* following exposure to ultraviolet radiation.** *Cell Tissue Res* 2007;329:159–68 [CrossRef Medline](#)
- Aldenderfer MS. **Moving up in the world.** *American Scientist* 2003;91:542–49
- Simonson TS, Yang Y, Huff CD, et al. **Genetic evidence for high-altitude adaptation in Tibet.** *Science* 2010;329:72–75 [CrossRef Medline](#)
- Cimini AM, Cerù MP. **Emerging roles of peroxisome proliferator-activated receptors (PPARs) in the regulation of neural stem cells proliferation and differentiation.** *Stem Cell Rev* 2008;4:293–303 [CrossRef Medline](#)
- Fidaleo M, Fanelli F, Ceru MP, et al. **Neuroprotective properties of peroxisome proliferator-activated receptor alpha (PPARα) and its lipid ligands.** *Curr Med Chem* 2014;21:2803–21 [CrossRef Medline](#)
- Zhang J, Yan X, Shi J, et al. **Structural modifications of the brain in acclimatization to high-altitude.** *PLoS One* 2010;5:e11449 [CrossRef Medline](#)
- Zhang H, Lin J, Sun Y, et al. **Compromised white matter microstructural integrity after mountain climbing: evidence from diffusion tensor imaging.** *High Alt Med Biol* 2012;13:118–25 [CrossRef Medline](#)
- Zhang J, Zhang H, Li J, et al. **Adaptive modulation of adult brain gray and white matter to high altitude: structural MRI studies.** *PLoS One* 2013;8:e68621 [CrossRef Medline](#)
- Zhang J, Zhang H, Chen J, et al. **Structural modulation of brain development by oxygen: evidence on adolescents migrating from high altitude to sea level environment.** *PLoS One* 2013;8:e67803 [CrossRef Medline](#)
- Kallenberg K, Bailey DM, Christ S, et al. **Magnetic resonance imaging evidence of cytotoxic cerebral edema in acute mountain sickness.** *J Cereb Blood Flow Metab* 2007;27:1064–71 [Medline](#)
- Zhang H, Wang X, Lin J, et al. **Grey and white matter abnormalities in chronic obstructive pulmonary disease: a case-control study.** *BMJ Open* 2012;2:e000844 [CrossRef Medline](#)
- Willie CK, Macleod DB, Shaw AD, et al. **Regional brain blood flow in man during acute changes in arterial blood gases.** *J Physiol* 2012;590:3261–75 [CrossRef Medline](#)
- Ogoh S, Sato K, Nakahara H, et al. **Effect of acute hypoxia on blood flow in vertebral and internal carotid arteries.** *Exp Physiol* 2013;98:692–98 [CrossRef Medline](#)
- Feddersen B, Neupane P, Thanbichler F, et al. **Regional differences in the cerebral blood flow velocity response to hypobaric hypoxia at high altitudes.** *J Cereb Blood Flow Metab* 2015;35:1846–51 [CrossRef Medline](#)
- Baril AA, Gagnon K, Arbour C, et al. **Regional cerebral blood flow during wakeful rest in older subjects with mild to severe obstructive sleep apnea.** *Sleep* 2015;38:1439–49 [CrossRef Medline](#)
- Lawley JS, Macdonald JH, Oliver SJ, et al. **Unexpected reductions in regional cerebral perfusion during prolonged hypoxia.** *J Physiol* 2016 Aug 9. [Epub ahead of print] [CrossRef Medline](#)
- DiPasquale DM, Strangman GE, Harris NS, et al. **Hypoxia, hypobaria, and exercise duration affect acute mountain sickness.** *Aerospace Med Hum Perform* 2015;86:614–19 [CrossRef Medline](#)
- Degache F, Larghi G, Faiss R, et al. **Hypobaric versus normobaric hypoxia: same effects on postural stability?** *High Alt Med Biol* 2012;13:40–45 [CrossRef Medline](#)
- McGuire S, Sherman P, Profenna L, et al. **White matter hyperintensities on MRI in high-altitude U-2 pilots.** *Neurology* 2013;81:729–35 [CrossRef Medline](#)
- Skovira JW, Kabadi SV, Wu J, et al. **Simulated aeromedical evacuation exacerbates experimental brain injury.** *J Neurotrauma* 2016;33:1292–302 [CrossRef Medline](#)
- He X, Su F, Taccone FS, et al. **Cardiovascular and microvascular responses to mild hypothermia in an ovine model.** *Resuscitation* 2012;83:760–66 [CrossRef Medline](#)
- Amir S, Robinson B. **Fos expression in rat visual cortex induced by**

- ocular input of ultraviolet light. *Brain Res* 1996;716:213–18 CrossRef Medline
40. Gofflot F, Charatoire N, Vasseur L, et al. **Systematic gene expression mapping clusters nuclear receptors according to their function in the brain.** *Cell* 2007;131:405–18 CrossRef Medline
41. Pérez-Martín M, Rivera P, Blanco E, et al. **Environmental enrichment, age, and PPAR α interact to regulate proliferation in neurogenic niches.** *Front Neurosci* 2016;10:89 CrossRef Medline
42. Knauf C, Rieusset J, Foretz M, et al. **Peroxisome proliferator-activated receptor- α -null mice have increased white adipose tissue glucose utilization, GLUT4, and fat mass: role in liver and brain.** *Endocrinology* 2006;147:4067–78 CrossRef Medline
43. Chakravarthy MV, Zhu Y, López M, et al. **Brain fatty acid synthase activates PPAR α to maintain energy homeostasis.** *J Clin Invest* 2007;117:2539–52 CrossRef Medline
44. Rizvi YQ, Mehta CS, Oyekan A. **Interactions of PPAR- α and adenosine receptors in hypoxia-induced angiogenesis.** *Vascu Pharmacol* 2013;59:144–51 CrossRef Medline
45. Hogan AM, Virues-Ortega J, Botti AB, et al. **Development of aptitude at altitude.** *Dev Sci* 2010;13:533–44 CrossRef Medline
46. Brutsaert TD, Soria R, Caceres E, et al. **Effect of developmental and ancestral high altitude exposure on chest morphology and pulmonary function in Andean and European/North American natives.** *Am J Hum Biol* 1999;1:383–95 Medline
47. Li W, Mai X, Liu C. **The default mode network and social understanding of others: what do brain connectivity studies tell us.** *Front Hum Neurosci* 2014;8:74 CrossRef Medline
48. Raichle ME. **The brain's default mode network.** *Annu Rev Neurosci* 2015;38:433–47 CrossRef Medline
49. Sui J, Huster R, Yu Q, et al. **Function-structure associations of the brain: evidence from multimodal connectivity and covariance studies.** *Neuroimage* 2014;102:11–23 CrossRef Medline
50. Soares JM, Sampaio A, Ferreira LM, et al. **Stress impact on resting state brain networks.** *PLoS One* 2013;8:e66500 CrossRef Medline
51. Henckens MJ, van der Marel K, van der Toorn A, et al. **Stress-induced alterations in large-scale functional networks of the rodent brain.** *Neuroimage* 2015;105:312–22 CrossRef Medline
52. Lu W, Wang Z. **Emotional expressivity, emotion regulation, and mood in college students: a cross-ethnic study.** *Social Behavior and Personality: An International Journal* 2012;40:319–30 CrossRef
53. Jin D. **Mental health problems and personal characters in Tibetan school children.** *Chinese Mental Health Journal* 1989;3:257–58
54. Kang YW, Kang XW, Xie SJ. **Analysis on Tibetan undergraduates' personality in inland.** *Chinese Journal of School Health* 2012;33:1176–79
55. Chang XD, Shi JH, Dan-Ying MA. **Correlation between personality traits and coping styles for junior high school students from Tibetan in Shanghai.** *Medical Journal of Chinese Peoples Health* 2013;25:94–97
56. Mason MF, Norton MI, Van Horn JD, et al. **Wandering minds: the default network and stimulus-independent thought.** *Science* 2007;315:393–95 CrossRef Medline
57. Sylvester CM, Corbetta M, Raichle ME, et al. **Functional network dysfunction in anxiety and anxiety disorders.** *Trends Neurosci* 2012;35:527–35 CrossRef Medline
58. Adelstein JS, Shehzad Z, Mennes M, et al. **Personality is reflected in the brain's intrinsic functional architecture.** *PLoS One* 2011;6:e27633 CrossRef Medline
59. Servaas MN, van der Velde J, Costafreda SG, et al. **Neuroticism and the brain: a quantitative meta-analysis of neuroimaging studies investigating emotion processing.** *Neurosci Biobehav Rev* 2013;37:1518–29 CrossRef Medline
60. Zhang J, Liu H, Yan X, et al. **Minimal effects on human memory following long-term living at moderate altitude.** *High Alt Med Biol* 2011;12:37–43 CrossRef Medline
61. Yan X, Zhang J, Gong Q, et al. **Cerebrovascular reactivity among native-raised high altitude residents: an fMRI study.** *BMC Neurosci* 2011;12:94 CrossRef Medline
62. Whittingstall K, Bernier M, Houde JC, et al. **Structural network underlying visuospatial imagery in humans.** *Cortex* 2014;56:85–98 CrossRef Medline
63. Kravitz DJ, Saleem KS, Baker CI, et al. **A new neural framework for visuospatial processing.** *Nat Rev Neurosci* 2011;12:217–30 CrossRef Medline
64. Antal A, Baudewig J, Paulus W, et al. **The posterior cingulate cortex and planum temporale/parietal operculum are activated by coherent visual motion.** *Vis Neurosci* 2008;25:17–26 CrossRef Medline
65. Fischer E, Bühlhoff HH, Logothetis NK, et al. **Visual motion responses in the posterior cingulate sulcus: a comparison to V5/MT and MST.** *Cereb Cortex* 2012;22:865–76 CrossRef Medline
66. Leech R, Sharp DJ. **The role of the posterior cingulate cortex in cognition and disease.** *Brain* 2014;137:12–32 CrossRef Medline
67. Bussey TJ, Muir JL, Everitt BJ, et al. **Dissociable effects of anterior and posterior cingulate cortex lesions on the acquisition of a conditional visual discrimination: facilitation of early learning vs. impairment of late learning.** *Behav Brain Res* 1996;82:45–56 CrossRef Medline
68. Zhai L, Li Q, Wang T, et al. **Altered functional connectivity density in high myopia.** *Behav Brain Res* 2016;303:85–92 CrossRef Medline
69. Ogawa K, Inui T. **The role of the posterior parietal cortex in drawing by copying.** *Neuropsychologia* 2009;47:1013–22 CrossRef Medline
70. Grön G, Wunderlich AP, Spitzer M, et al. **Brain activation during human navigation: gender-different neural networks as substrate of performance.** *Nat Neurosci* 2000;3:404–08 CrossRef Medline
71. Prvulovic D, Hubl D, Sack AT, et al. **Functional imaging of visuospatial processing in Alzheimer's disease.** *Neuroimage* 2002;17:1403–14 CrossRef Medline
72. Chee MW, Chen KH, Zheng H, et al. **Cognitive function and brain structure correlations in healthy elderly East Asians.** *Neuroimage* 2009;46:257–69 CrossRef Medline
73. Goldman JG, Stebbins GT, Dinh V, et al. **Visuo-perceptive region atrophy independent of cognitive status in patients with Parkinson's disease with hallucinations.** *Brain* 2014;137:849–59 CrossRef Medline
74. Paulus MP, Poteriat EG, Taylor MK, et al. **A neuroscience approach to optimizing brain resources for human performance in extreme environments.** *Neurosci Biobehav Rev* 2009;33:1080–88 CrossRef Medline
75. Dani JA, Montague PR. **Disrupting addiction through the loss of drug associated internal states.** *Nat Neurosci* 2007;10:403–04 CrossRef Medline
76. Frischo AR. **Functional adaptation to high altitude hypoxia.** *Science* 1975;187:313–19 CrossRef Medline
77. Han X, Jovicich J, Salat D, et al. **Reliability of MRI-derived measurements of human cerebral cortical thickness: the effects of field strength, scanner upgrade and manufacturer.** *Neuroimage* 2006;32:180–94 CrossRef Medline
78. Durand-Dubief F, Belaroussi B, Armspach JP, et al. **Reliability of longitudinal brain volume loss measurements between 2 sites in patients with multiple sclerosis: comparison of 7 quantification techniques.** *AJNR Am J Neuroradiol* 2012;33:1918–24 CrossRef Medline
79. McKenzie DC, Goodman LS, Nath C, et al. **Cardiovascular adaptations in Andean natives after 6 wk of exposure to sea level.** *J Appl Physiol* (1985) 1991;70:2650–55 Medline
80. Sime F, Peñaloza D, Ruiz L. **Bradycardia, increased cardiac output, and reversal of pulmonary hypertension in altitude natives living at sea level.** *Br Heart J* 1971;33:647–57 CrossRef Medline
81. Bai J, Abdul-Rahman MF, Rifkin-Graboi A, et al. **Population differences in brain morphology and microstructure among Chinese, Malay, and Indian neonates.** *PLoS One* 2012;7:e47816 CrossRef Medline