REPLY:

We thank our colleagues for their comments about our recent article, “Blood Flow Mimicking Aneurysmal Wall Enhancement: A Diagnostic Pitfall of Vessel Wall MRI Using the Postcontrast 3D Turbo Spin-Echo MR Imaging Sequence.”¹

Black-blood techniques such as motion-sensitized driven equilibrium or delay alternating with nutation for tailored excitation (DANTE) offer new perspectives for a better understanding of the nature of intracranial aneurysm enhancement. In our experience, flow-related artifacts are frequently encountered using the postcontrast 3D-TSE sequence for a wide range of incidental aneurysms (at the center of the aneurysmal cavity but also in contact with the wall of the aneurysm). It is now clear that the slow-flowing blood near the aneurysmal wall may represent all or part of the visible enhancement using conventional 3D-TSE sequences. Such findings raise the question of the relevance of this parameter for the management of asymptomatic patients.

The relationship among contrast enhancement on 3D-TSE, slow-flowing blood, low shear stress, and aneurysmal instability is poorly reported and is still speculative. In addition, studies with a long-term follow-up and/or histologic correlation are lacking in the literature. Thus, it appears difficult to assume that the enhancement related to slow-flowing blood and that of the aneurysmal wall itself are both related to inflammation/instability. Quite on the opposite, we think that it would be interesting to independently analyze each component of the enhancement visible on conventional 3D-TSE, including slow-flowing blood. Such an approach would allow us to identify patients in whom the slow-flowing blood is predominant versus those in whom the addition of a black-blood technique has little or no influence on the en-

hancement already visible on conventional sequences (and therefore potentially related to a “true” enhancement of the wall).

A better understanding of the factors likely to promote slow-flowing blood near the aneurysmal wall (such as the size of the aneurysm?) also appears important. In our opinion, the real ability of the 3D-TSE sequence to assess arterial wall inflammation remains uncertain because we cannot formally exclude residual slow-flowing blood (even including the additional black-blood technique). Further studies are required to evaluate alternative/complementary MR sequences less sensitive to flow artifacts and with a higher spatial resolution.

We hope that this discussion will encourage other teams to continue evaluating the circumferential wall enhancement of intracranial aneurysms, which remains a particularly complex entity.

REFERENCE

1. Kalsoum E, Chabernaude Negrier A, Tuilier T, et al. **Blood flow mimicking aneurysmal wall enhancement: a diagnostic pitfall of vessel wall MRI using the postcontrast 3D turbo spin-echo MR imaging sequence.** *AJNR Am J Neuroradiol* 2018;39:1065–67 [CrossRef](#) [Medline](#)

J. Hodel

E. Kalsoum

T. Tuilier

A. Benaïssa

Department of Neuroradiology
Centre Hospitalier Universitaire Henri Mondor
Créteil, France

R. Blanc

Department of Interventional Neuroradiology
Rothschild Foundation Hospital
Paris, France

P. Brugières

Department of Neuroradiology
Centre Hospitalier Universitaire Henri Mondor
Créteil, France

<http://dx.doi.org/10.3174/ajnr.A5789>