

# Application of 3D T1 Black-Blood Imaging in the Diagnosis of Leptomeningeal Carcinomatosis: Potential Pitfall of Slow-Flowing Blood

I read the publication “Application of 3D Fast Spin-Echo T1 Black-Blood Imaging in the Diagnosis and Prognostic Prediction of Patients with Leptomeningeal Carcinomatosis” by Oh et al<sup>1</sup> with a great interest. The authors concluded that black-blood imaging showed a significantly higher sensitivity than contrast-enhanced gradient recalled-echo and contrast-enhanced spin-echo imaging for detecting leptomeningeal carcinomatosis.

A variety of techniques can be used to achieve blood suppression on T1-weighted imaging. The most commonly used technique, which was also used by Oh et al,<sup>1</sup> is a variable flip angle refocusing pulse sequence in which the protons in the vessel lumen experience the slice-selective radiofrequency pulse but flow out of the imaging section before the refocusing pulse, resulting in blood-signal suppression. This technique is widely used in high-resolution intracranial vessel wall MR imaging; however, an important pitfall with this technique is that slow-flowing blood in leptomeningeal veins, dilated arteries, or leptomeningeal collaterals can cause incomplete or lack of suppression.<sup>2,3</sup> Kato et al<sup>4</sup> compared 3D fast spin-echo (sampling perfection with application-optimized contrasts by using different flip angle evolutions [SPACE; Siemens, Erlangen, Germany]) and 3D gradient-echo T1-weighted MPRAGE images in patients with small parenchymal brain metastasis. Lesion detectability was significantly higher on SPACE than on MPRAGE; however, vessels were falsely reported as metastasis using both techniques. I can only imagine that this pitfall will be aggravated when assessing leptomeningeal metastasis. One way to avoid these artifacts would be to use a double inversion recovery technique, which exploits both the flow

and T1 properties of blood to suppress its signal.<sup>2</sup> This technique requires a longer acquisition time, which is a limitation in high-resolution intracranial vessel wall MR imaging, given the need for very high spatial resolution; however, this should be less of a problem in the context of metastatic disease.

In conclusion, I agree with the authors that postcontrast T1 black-blood imaging is a promising technique for the detection of leptomeningeal carcinomatosis; however, it will require further investigation to determine the best technique for blood suppression to avoid the above-mentioned pitfall.

## REFERENCES

1. Oh J, Choi SH, Lee E, et al. **Application of 3D fast spin-echo T1 black-blood imaging in the diagnosis and prognostic prediction of patients with leptomeningeal carcinomatosis.** *AJNR Am J Neuroradiol* 2018; 39:1453–59 CrossRef Medline
2. Mandell DM, Mossa-Basha M, Qiao Y, et al; Vessel Wall Imaging Study Group of the American Society of Neuroradiology. **Intracranial vessel wall MRI: principles and expert consensus recommendations of the American Society of Neuroradiology.** *AJNR Am J Neuroradiol* 2017; 38:218–29 CrossRef Medline
3. Lindenholz A, van der Kolk AG, Zwanenburg JJ, et al. **The use and pitfalls of intracranial vessel wall imaging: how we do it.** *Radiology* 2018;286:12–28 CrossRef Medline
4. Kato Y, Higano S, Tamura H, et al. **Usefulness of contrast-enhanced T1-weighted sampling perfection with application-optimized contrasts by using different flip angle evolutions in detection of small brain metastasis at 3T MR imaging: comparison with magnetization-prepared rapid acquisition of gradient echo imaging.** *AJNR Am J Neuroradiol* 2009;30:923–29 CrossRef Medline

© S.A. Nabavizadeh

Department of Neuroradiology  
Hospital of the University of Pennsylvania  
Philadelphia, Pennsylvania

<http://dx.doi.org/10.3174/ajnr.A5832>