

Celebrating 35 Years of the AJNR


November 1985 edition

Contrast-Enhanced MR Imaging of Malignant Brain Tumors

Moshe Graif¹
Graeme M. Bydder²
Robert E. Gleister³
Peter Niendorf²
David G. T. Thomas³
Ian R. Young⁴

Magnetic resonance (MR) imaging was performed before and serially after intravenous injection of 1 mmol/kg gadolinium-DTPA (Schering) in 17 patients with clinical and histologic diagnoses of malignant cerebral tumors. There was a decrease of 1% in T1 and 10% in T2 in normal white matter and a decrease of 8% in T1 and 14% in T2 in normal gray matter. Contrast enhancement was observed in 16 of the 17 tumors. In the region of maximal enhancement a mean decrease of 18% in T1 was observed in low-grade gliomas, a mean decrease in T1 of 29% was seen in high-malignancy gliomas, and a mean decrease in T1 of 33% was observed in metastases. The decreases in T1 persisted for at least 50 min. In one case the central cystic region of the tumor displayed a decrease in T1 and T2. Measurements of signal intensity displayed maximal contrast enhancement with an IR 1500/500/44 sequence, much less with SE 1500/44, and least with SE 1500/80. Edema was observed on precontrast images in 14 cases, but satisfactory definition of the tumor-edema margin was only possible in four cases. After contrast enhancement this margin was defined in 10 cases. In four of the 17 cases areas of apparent "redness" seen before administration of Gd-DTPA displayed significant contrast enhancement and probably represented tumor infiltration. Comparison with CT showed a greater degree of contrast enhancement on MR images in eight cases, an equal degree in eight cases, and greater enhancement on CT in one case. Definition of the tumor-edema interface was better with MR in six cases and equal to CT in four cases. No significant toxic effects were found with the contrast agent. Gd-DTPA provided significant additional information over unenhanced MR imaging and enhanced CT.

The value of contrast enhancement was recognized soon after the introduction of cranial CT [1, 2]. The situation with magnetic resonance (MR) imaging is more complex. A greater level of soft-tissue contrast is available without contrast enhancement, and, unlike CT, no contrast agents suitable for parenteral use were available when the technique was first introduced. Nevertheless, particular situations have been defined, such as differentiation between tumor and edema, where contrast enhancement might be useful [3, 4], and recently the paramagnetic contrast agent gadolinium-DTPA has been tested in animals [5, 6] and used in clinical pilot studies [7-9]. In this report we extend our earlier clinical studies by



Received March 6, 1985; accepted after revision June 12, 1985.

This work was supported in part by Schering A.G., Berlin (Gd-DTPA), and Medical Research Council, Department of Health and Social Security.

¹Department of Diagnostic Radiology, Royal Postgraduate Medical School, HammerSmith Hospital, Du Cane Rd., London W12 0HS, United Kingdom. Address reprint requests to M. Graif, NMR Unit.

²Schering AG, Berlin, West Germany.

³Department of Neurosurgery, Maida Vale Hospital, Maida Vale, London W9, United Kingdom.


⁴Imperial International, Viral Research Centre, Wembley, Middlesex, United Kingdom.

AJNR 6:855-862, November/December 1985
0195-6108/85/0605-0855
© American Roentgen Ray Society

Cavernous Sinus Invasion by Pituitary Adenomas

Jamshid Ahmadi¹
Charles M. North¹
Henry D. Sogali¹
Chi-Shing Zee¹
Martin H. Weiss²

One hundred ninety-eight surgically explored pituitary adenomas were evaluated preoperatively by high-resolution computed tomography (CT). At surgery, evidence of direct cavernous sinus invasion was demonstrated in 19. CT findings in these cases included cavernous sinus expansion (17 patients) and visible encasement of the internal carotid artery (14 patients). The invasive tumor often enhanced to a lesser degree than the cavernous sinuses and ipsilateral internal carotid artery. Intracavernous cranial nerve compression, obliteration, or displacement (14 patients), invasion of the lateral wall of the cavernous sinus (seven patients), and diffuse bone destruction (seven cases) were other findings. Magnetic resonance imaging in three patients provided excellent demonstration of intracavernous internal carotid artery encasement, but displacement and obliteration of intracavernous cranial nerves was not shown as well as it was with CT. Histologically, only three patients showed anaplastic features and only one of them had distant metastases. There was no correlation between histologic features, hormone assays, and invasiveness. This experience indicates any type of pituitary adenoma, regardless of its endocrinologic activity, can invade the cavernous sinus. Cavernous sinus involvement makes complete surgical removal difficult. Preoperative recognition of invasive behavior of these tumors has prognostic value and aids in designing appropriate management. CT is the most useful technique generally available for evaluation and follow-up.



This article appears in the November 1985 issue of AJNR and the issue of AJNR.

Received December 12, 1984; revision May 14, 1985.

¹Department of Radiology, Scripps Clinic, University of Southern California, Los Angeles, CA 90033; requests to J. Ahmadi, LAQJ/SF, Box 2, 1200 N. State St., Los Angeles, CA 90033.

²Department of Neurological Surgery, University of Southern California School of Medicine, Los Angeles, CA 90033.
AJNR 6:863-868, November/December 1985
0195-6108/85/0605-0863
© American Roentgen Ray Society