REPLY:

We thank Morelli et al for their interest in our article and their feedback.

We think it is important to remember that coronavirus disease 2019 (COVID-19) was first recognized in December 2019, only 6 months ago.¹ Therefore, there will be inherent limitations to the size, scope, and implications of any research that can occur in such a truncated period.

In addition, as we acknowledge in our article, the patient described in our case report died on hospital day 3, before the completion of a complete laboratory work-up; furthermore, no postmortem examination was performed.² So, we agree with Morelli et al that it will require larger studies of more prolonged duration to draw definitive causal relationships between COVID-19 and various disease states, including acute cerebrovascular disease.

Until such studies occur, we would refer Morelli et al to the early-but-substantial medical literature that indicates a strong association between COVID-19 and cerebrovascular disease. This includes a retrospective study of 214 patients with COVID-19 in China, which revealed a 5.7% incidence of stroke in severely affected patients.³ In a case series of 64 patients with severe COVID-19, three of 13 patients who underwent brain MRIs demonstrated evidence of ischemic stroke.⁴ Researchers in New York City recently published a case series of 5 young patients (less than 50 years-old) with COVID-19 who presented with large-vessel stroke during a 2-week period; this represented an almost 7-fold increase in the frequency of large-vessel stroke seen in patients in this age group.⁵ Because severely affected patients with COVID-19 are often intubated and sedated, the presence of stroke can be overlooked, and the current estimates of the incidence of stroke may, therefore, be spuriously low.⁶ Additionally, Umaphathi et al⁷ reported an association between large-vessel stroke and the somewhat similar 2004 Severe Acute Respiratory Syndrome coronavirus (SARS-CoV-1) outbreak. Combined with the growing literature on coagulopathy in severely affected patients with COVID-19, we think that dismissing this information as anecdotal belies reasonable evidence for a causative association between COVID-19 and cerebrovascular disease.⁸⁻¹⁰

Like Morelli et al, we also subscribe to the principle of Occam's razor. The patient described in our case report had tested positive for SARS-CoV-2 sixteen days before admission. Initially, he had a relatively benign course with mild respiratory symptoms and myalgia, and he had no preceding neurologic deficits. Several hours before admission, however, he developed, nearly simultaneously, stroke symptoms and respiratory distress, which ultimately led to CT-confirmed acute respiratory distress syndrome (ARDS) and hemodynamic instability. We do not think the synchronous timing of these disparate symptoms is coincidental. Rather, the delayed

onset of both respiratory and neurologic symptoms in COVID-19 is suggestive of a maladaptive immune response, including cytokine storm syndrome, resulting in not only ARDS but also abnormalities in the clotting cascade and endothelial inflammation.^{11,12} Therefore, ascribing this patient's stroke to COVID-19 is entirely consistent with Occam's razor—to do otherwise risks invoking the Hickam dictum, which is the assignment of multiple different and independent diagnoses to explain a variety of signs and symptoms.¹³

REFERENCES

- Phelan AL, Katz R, Gostin LO. **The novel coronavirus originating in Wuhan, China: challenges for global health governance.** *JAMA* 2020 Jan 30. [Epub ahead of print] [CrossRef Medline](#)
- Goldberg MF, Goldberg MF, Cerejo R, et al. **Cerebrovascular disease in COVID-19.** *AJNR Am J Neuroradiol* 2020;41:1170–72 [CrossRef Medline](#)
- Mao L, Jin H, Wang M, et al. **Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China.** *JAMA Neurol* 2020;77:1–9 [CrossRef Medline](#)
- Helms J, Kremer S, Merdji H, et al. **Neurologic features in severe SARS-CoV-2 infection.** *N Engl J Med* 2020;382:2268–70 [CrossRef Medline](#)
- Oxley TJ, Mocco J, Majidi S, et al. **Large-vessel stroke as a presenting feature of Covid-19 in the young.** *N Engl J Med* 2020;382:e60 [CrossRef Medline](#)
- Yaghi S, Ishida K, Torres J, et al. **SARS2-CoV-2 and stroke in a New York health care system.** *Stroke* 2020 May 20. [Epub ahead of print] [CrossRef Medline](#)
- Umaphathi T, Kor AC, Venketasubramanian N, et al. **Large-artery ischaemic stroke in severe acute respiratory syndrome (SARS).** *J Neurol* 2004;251:1227–31 [CrossRef Medline](#)
- Wan S, Xiang Y, Fang W, et al. **Clinical features and treatment of COVID-19 patients in Northeast Chongqing.** *J Med Virol* 2020;92:797–806 [CrossRef Medline](#)
- Helms J, Tacquard C, Severac F, et al. **High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study.** *Intensive Care Med* 2020;46:1089–98 [CrossRef Medline](#)
- Arachchillage DR, Laffan M. **Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia.** *J Thromb Haemost* 2020;18:844–47 [CrossRef Medline](#)
- Mehta P, McAuley DF, Brown M, et al. **COVID-19: consider cytokine storm syndromes and immunosuppression.** *Lancet* 2020;395:1033–34 [CrossRef Medline](#)
- Berlin DA, Gulick RM, Martinez FJ. **Severe Covid-19.** *N Engl J Med* 2020 May 15. [Epub ahead of print] [CrossRef Medline](#)
- Miller WT. **Letter from the editor: Occam versus Hickam.** *Semin Roentgenol* 1998;33:213 [CrossRef](#)

Michael F. Goldberg

Imaging Institute, Division of Neuroradiology
Allegheny Health Network
Pittsburgh, Pennsylvania
Baltimore, Maryland

Morton F. Goldberg

Department of Ophthalmology
Johns Hopkins Medical Institutions

R. Cerejo

A.H. Tayal

Neuroscience Institute
Cerebrovascular Center
Allegheny Health Network
Pittsburgh, Pennsylvania

Indicates open access to non-subscribers at www.ajnr.org

<http://dx.doi.org/10.3174/ajnr.A6686>