

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



FRESENIUS
KABI

caring for life

AJNR

Reply

M. Petruzzellis, R. De Blasi, V. Lucivero, M. Sancilio, M. Prontera, A. Tinelli, D.M. Mezzapesa and F. Federico

AJNR Am J Neuroradiol published online 26 September 2007

<http://www.ajnr.org/content/early/2007/09/26/ajnr.A0731.citation>

This information is current as of April 19, 2024.

Reply:

We thank Dr. Fox for his comments concerning our previously published letter to the editor entitled, "Cerebral Aneurysms in a Patient with Osteogenesis Imperfecta and Exon 28 Polymorphism Of COL1A2."¹ He strongly informs us of a misunderstanding regarding the de novo aneurysm that developed after 4 months. Dr. Fox points out some potential errors in our diagnosis, claiming that the de novo aneurysm is actually a refilling of the previously treated one. We believe that indeed at 4 months, our images revealed that a new aneurysm had developed in front of the previous one (Fig 2B). Further evidence of this was found in the posttreatment un-subtracted images (not published due to space constraints), in which it was possible to appreciate the stent crossing the vertebrobasilar junction and the coils occluding 2 different and clearly separable aneurysms.

Finally, we are grateful to Dr. Fox for bringing to our attention some bibliographic references that may help us achieve a better understanding of the anatomic features of this particular case.

Reference

1. Petruzzellis M, De Blasi R, Lucivero V, et al. Cerebral aneurysms in a patient with osteogenesis imperfecta and exon 28 polymorphism of COL1A2. *AJNR Am J Neuroradiol* 2007;28:397-98

M. Petruzzellis
R. De Blasi
V. Lucivero
M. Sancilio
M. Prontera
A. Tinelli
D.M. Mezzapesa
F. Federico

*Dipartimento di Scienze Neurologiche e Psichiatriche
Università degli Studi di Bari-Italia*

Bari, Italy

DOI 10.3174/ajnr.A0731