Are your MRI contrast agents cost-effective? Learn more about generic Gadolinium-Based Contrast Agents.

FRESENIUS KABI
caring for life



"Bovine" Aortic Arch

Allen D. Elster

AJNR Am J Neuroradiol published online 26 November 2007

http://www.ajnr.org/content/early/2007/11/26/ajnr.A0861.cit ation

This information is current as of April 10, 2024.

Published November 26, 2007 as 10.3174/ajnr.A0861

"Bovine" Aortic Arch

I have reviewed with interest the recent paper by Layton et al¹ and correspondence by Vitek² concerning the human anatomic variation commonly but incorrectly referred to as the "bovine" aortic arch.

As a point of comparative anatomy, it should be noted that several common mammals have normal aortic branching patterns identical with the human "bovine" variant. Specifically, dogs, cats, and most rabbit species have 2 major branches from the aortic arch, the first being an innominate or brachiocephalic artery giving origin sequentially to the left common carotid and right common carotid arteries and the second being a left subclavian artery.³

I suggest, therefore, that if tempted to apply an animal name to the human arch variant described above, using a term like "canine," "feline," or "lapine" aortic arch would be more appropriate.

References

- 1. Layton KF, Kallmes DF, Cloft HJ, et al. **Bovine aortic arch araiant in humans:** clarification of a common misnomer. *AJNR Am J Neuroradiol* 2006;27: 1541, 42
- 2. Vitek JJ. Bovine aortic arch. AJNR Am J Neuroradiol 2007;2487:1205
- 3. Dome SH, Goody PC, Strickland NC, et al. Color Atlas of Veterinary Anatomy: Volume 3, The Dog and Cat. St Louis: Mosby-Year Book; 1996

Allen D. Elster

Department of Radiology

Wake Forest University School of Medicine

Winston-Salem, NC

DOI 10.3174/ajnr.A0861