

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



FRESENIUS
KABI

caring for life

AJNR

Reply:

D.G. Na and E. Kim

AJNR Am J Neuroradiol published online 19 June 2008

<http://www.ajnr.org/content/early/2008/06/19/ajnr.A1145.1.citation>

This information is current as
of April 17, 2024.

Reply:

We thank Drs. Zuccoli, Pipitone, and Cruz for their interest in our recent article regarding MR imaging features of metronidazole-induced encephalopathy (MIE).¹ Although the exact mechanisms of MIE and Wernicke encephalopathy (WE) are not elucidated, we agree with the opinion that MIE may be mediated by pathogenic pathways similar to those thought to be operating in WE.

In our article, we reported that the brain lesions of MIE were typically located at the cerebellar dentate nucleus, midbrain, dorsal pons, medulla, and splenium of the corpus callosum. The recently published case reports of WE show lesions in the cerebellar dentate^{2,3} and focal tegmental lesions of low pons,^{3,4} which we described as characteristic lesion locations of MIE and controversial atypical MR features of WE in the article.¹ These recent reports²⁻⁴ suggest that the MR imaging feature of bilateral symmetric involvement of dentate nuclei and cranial nerve nuclei may be found in WE, as well as in MIE. However, in these WE cases,²⁻⁴ there was involvement of at least one of medial thalamus, mammillary body, or pericentral cortices, which were well-documented MR features in WE but were not reported in MIE. Therefore, although 2 diseases of MIE and WE may share the similar pathogenic pathways for metabolic pathology and similar MR

features in the cerebellum and brain stem, the analysis of lesion distribution on MR images may still be useful in distinguishing MIE from WE.

References

1. Kim E, Na DG, Kim EY, et al. **MR imaging of metronidazole-induced encephalopathy: lesion distribution and diffusion-weighted imaging findings.** *AJNR Am J Neuroradiol* 2007;28:1652-58
2. Liu YT, Fuh JL, Lirng JF, et al. **Correlation of magnetic resonance images with neuropathology in acute Wernicke's encephalopathy.** *Clin Neurol Neurosurg* 2006;108:682-87
3. Zuccoli G, Motti L. **Atypical Wernicke's encephalopathy showing lesions in the cranial nerve nuclei and cerebellum.** *J Neuroimaging* 2008;18:194-97
4. Nolli M, Barbieri A, Pinna C, et al. **Wernicke's encephalopathy in a malnourished surgical patient: clinical features and magnetic resonance imaging.** *Acta Anaesthesiol Scand* 2005;49:1566-70

D.G. Na

*Human Medical Imaging and Intervention Center
Seoul National University Hospital*

E. Kim

*Seoul National University Hospital
Seoul, Korea*

DOI 10.3174/ajnr.A1145