

Are your **MRI contrast agents** cost-effective?

Learn more about generic **Gadolinium-Based Contrast Agents**.



**FRESENIUS  
KABI**

caring for life

**AJNR**

## **Superficial Siderosis in Cerebral Amyloid Angiopathy**

J. Linn and H. Brückmann

*AJNR Am J Neuroradiol* published online 22 October 2009  
<http://www.ajnr.org/content/early/2009/10/22/ajnr.A1913.citation>

This information is current as  
of April 19, 2024.

### Superficial Siderosis in Cerebral Amyloid Angiopathy

With great interest, we read the recent review article by Kumar entitled "Neuroimaging in Superficial Siderosis: An In-Depth Look," published on-line ahead of print in the *American Journal of Neuroradiology* in September 2009.<sup>1</sup> The author provides a thorough review of neuroimaging in superficial siderosis (SS) of the central nervous system and details important underlying causes of this phenomenon. Besides the common mechanisms (eg, history of trauma or intradural surgery), the author lists cerebral amyloid angiopathy (CAA) as a potential pathomechanism for SS.<sup>1</sup> We want to underline the role of this microangiopathic disease as an important cause of SS.

Most interesting, the published CAA cases with SS lack the typical clinical findings of "classic" SS, which are progressive gait ataxia with cerebellar dysarthria and sensorineural hearing loss, but patients often present with headache, seizures, and cognitive impairment.<sup>2-4</sup> This is most probably due to the characteristic localization of SS in patients with CAA: While the "classic" SS mainly affects brain stem and posterior fossa,<sup>1</sup> SS in CAA is typically found in a supratentorial distribution over the cerebral convexities.<sup>2-4</sup>

In our opinion, CAA should be thoroughly considered as a cause

of SS, especially in older patients with isolated supratentorial SS and an atypical clinical presentation. Further studies on the sensitivity and specificity of SS as a noninvasive diagnostic MR imaging sign of CAA are necessary.

### References

1. Kumar N. **Neuroimaging in superficial siderosis: an in-depth look.** *AJNR Am J Neuroradiol* 2009 Sep 3. [Epub ahead of print]
2. Linn J, Herms J, Dichgans M, et al. **Subarachnoid hemosiderosis and superficial cortical hemosiderosis in cerebral amyloid angiopathy.** *AJNR Am J Neuroradiol* 2008;29:184–86
3. Roch JA, Nighoghossian N, Hermier M, et al. **Transient neurologic symptoms related to cerebral amyloid angiopathy: usefulness of T2\*-weighted imaging.** *Cerebrovasc Dis* 2005;20:412–14
4. Feldman HH, Maia LF, Mackenzie IR, et al. **Superficial siderosis: a potential diagnostic marker of cerebral amyloid angiopathy in Alzheimer disease.** *Stroke* 2008;39:2894–97

J. Linn

H. Brückmann

Department of Neuroradiology

University Hospital Munich

Munich, Germany

DOI 10.3174/ajnr.A1913