

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



**FRESENIUS
KABI**

caring for life

AJNR

**Predicting Genotype and Survival in Glioma
Using Standard Clinical MR Imaging
Apparent Diffusion Coefficient Images: A
Pilot Study from The Cancer Genome Atlas**

C.-C. Wu, R. Jain, A. Radmanesh, L.M. Poisson, W.-Y.
Guo, D. Zagzag, M. Snuderl, D.G. Placantonakis, J. Golfinos
and A.S. Chi

This information is current as
of April 23, 2024.

AJNR Am J Neuroradiol published online 6 September 2018
<http://www.ajnr.org/content/early/2018/09/06/ajnr.A5794>

Predicting Genotype and Survival in Glioma Using Standard Clinical MR Imaging Apparent Diffusion Coefficient Images: A Pilot Study from The Cancer Genome Atlas

C.-C. Wu, R. Jain, A. Radmanesh, L.M. Poisson, W.-Y. Guo, D. Zagzag, M. Snuderl, D.G. Placantonakis, J. Golfinos, and A.S. Chi

ABSTRACT

BACKGROUND AND PURPOSE: Few studies have shown MR imaging features and ADC correlating with molecular markers and survival in patients with glioma. Our purpose was to correlate MR imaging features and ADC with molecular subtyping and survival in adult diffuse gliomas.

MATERIALS AND METHODS: Presurgical MRIs and ADC maps of 131 patients with diffuse gliomas and available molecular and survival data from The Cancer Genome Atlas were reviewed. MR imaging features, ADC (obtained by ROIs within the lowest ADC area), and mean relative ADC values were evaluated to predict *isocitrate dehydrogenase* (*IDH*) mutation, 1p/19q codeletion status, *MGMT* promoter methylation, and overall survival.

RESULTS: *IDH* wild-type gliomas tended to exhibit enhancement, necrosis, and edema; >50% enhancing area ($P < .001$); absence of a cystic area ($P = .013$); and lower mean relative ADC (median, 1.1 versus 1.6; $P < .001$) than *IDH*-mutant gliomas. By means of a cutoff value of 1.08 for mean relative ADC, *IDH*-mutant and *IDH* wild-type gliomas with lower mean relative ADC (<1.08) had poorer survival than those with higher mean relative ADC (median survival time, 24.2 months; 95% CI, 0.0–54.9 months versus 62.0 months; $P = .003$; and median survival time, 10.4 months; 95% CI, 4.4–16.4 months versus 17.7 months; 95% CI, 11.6–23.7 months; $P = .041$, respectively), regardless of World Health Organization grade. Median survival of those with *IDH*-mutant glioma with low mean relative ADC was not significantly different from that in those with *IDH* wild-type glioma. Other MR imaging features were not statistically significant predictors of survival.

CONCLUSIONS: *IDH* wild-type glioma showed lower ADC values, which also correlated with poor survival in both *IDH*-mutant and *IDH* wild-type gliomas, irrespective of histologic grade. A subgroup with *IDH*-mutant gliomas with lower ADC had dismal survival similar to that of those with *IDH* wild-type gliomas.

ABBREVIATIONS: *IDH* = *isocitrate dehydrogenase*; max = maximum; min = minimum; rADC = relative ADC; rADC_{mean} = mean relative ADC; TCGA = The Cancer Genome Atlas; WHO = World Health Organization

Gliomas are a heterogeneous group of tumors, and the clinical aggressiveness and prognoses are diverse among different histopathologic grades and molecular subtypes. Previous studies have shown that histopathologic classification of diffuse gliomas

has high interobserver variation and correlates imperfectly with clinical outcomes.^{1,2} Nevertheless, molecular markers, particularly *isocitrate dehydrogenase* (*IDH*) mutational status, have been demonstrated to be significant and more robust prognostic markers³ and have been incorporated into the classification of diffuse gliomas in the latest update of the World Health Organization (WHO) classification in 2016.⁴ *IDH* mutation, a powerful prognostic marker of improved survival in diffuse glioma, is found mainly in lower grade gliomas (WHO grades II and III), but also in glioblastoma (WHO grade IV), though at much lower frequency.^{5,6}

Preoperative and noninvasive determination of molecular subtyping is of great value in the clinical management of patients with glioma. However, studies correlating MR imaging features with *IDH*-mutation status and patient survival in diffuse gliomas are scarce. Recently, we showed that the “T2-FLAIR mismatch sign,” detectable using conventional MR imaging, is a highly spe-

Received April 25, 2018; accepted after revision July 2.

From the Department of Radiology (C.-C.W., W.-Y.G.), Taipei Veterans General Hospital, Taipei, Taiwan, Republic of China; School of Medicine (C.-C.W., W.-Y.G.), National Yang-Ming University, Taipei, Taiwan, Republic of China; Departments of Radiology (C.-C.W., R.J., A.R.), Neurosurgery (R.J., D.P., J.G.), and Pathology (D.Z., M.S.), NYU School of Medicine, New York, New York; Department of Public Health Sciences and Hermelin Brain Tumor Center (L.M.P.), Henry Ford Hospital, Detroit, Michigan; and Neuro-Oncology Program (A.S.C.), Laura and Isaac Perlmutter Cancer Center, NYU School of Medicine and Langone Health, New York, New York.

Chih-Chin Wu and Rajan Jain contributed equally to the article.

Please address correspondence to Rajan Jain, MD, Department of Radiology, NYU School of Medicine, 660 First Ave, 2nd Floor, New York, NY 10016; e-mail: rajan.jain@nyumc.org

<http://dx.doi.org/10.3174/ajnr.A5794>

cific imaging biomarker for the *IDH*-mutant, 1p/19q noncodeleted molecular subtype in lower grade gliomas.⁷ Wang et al⁸ demonstrated that the absence of contrast enhancement was associated with longer progression-free and overall survival in patients with *IDH1*-mutated anaplastic gliomas. MR spectroscopy could detect 2-hydroxyglutarate, a metabolite that accumulates in *IDH*-mutant gliomas but did not discover a survival difference.⁹ Blood volume estimates obtained by MR perfusion have also provided potential markers for noninvasive assessment of *IDH* status.¹⁰

ADC can be calculated from DWI, and tumors with more freely mobile water molecules and lesser cellularity have higher ADC values.¹¹ ADC has been shown to be a valuable imaging marker in the diagnosis of intracranial lesions as well as in grading brain tumors.¹¹⁻¹⁵ Therefore, we hypothesized that ADC values obtained from conventional MR imaging could correlate with molecular subtype and patient survival in adult diffuse gliomas.

MATERIALS AND METHODS

This was a retrospective study using data from the publicly available National Institutes of Health/National Cancer Institute–approved databases of The Cancer Genome Atlas (TCGA; <https://cancergenome.nih.gov>) and The Cancer Imaging Archive (<http://www.cancerimagingarchive.net/>),¹⁶⁻¹⁸ from which all 461 cases with imaging data were reviewed, and only cases of treatment-naïve diffuse gliomas (WHO grades II–IV) with available DWI and ADC maps were included. WHO grade, the status of 3 validated molecular prognostic markers (*IDH* mutation, 1p/19q codeletion, *MGMT* promoter methylation), and survival data were retrieved from The Cancer Genome Atlas. MR images were reviewed, in consensus, by 2 board-certified neuroradiologists (with 8 and 17 years of experience) who were blinded to pathologic and molecular diagnosis. The order of cases viewed was randomized to avoid bias.

Each tumor was scored for 9 MR imaging features according to the following criteria modified from the Visually Accessible Rembrandt Images MR imaging feature set¹⁹: T2 signal intensities (higher than gray matter or mixed [the presence equal to or darker than that of gray matter part]); T2 homogeneity (homogeneous or heterogeneous); margin (well-defined or not well-defined [either infiltrative or irregular]); edema (none to minimal or mild to marked); enhancing pattern (non-/minimally enhancing or enhancing); portion of enhancing area (<50% or ≥50%); the presence of cystic areas (presence or absence); and the presence of necrotic areas (presence or absence). We investigated the relationship among 9 different MR imaging features and 3 molecular markers (*IDH* mutation, 1p/19q codeletion, and *MGMT* promoter methylation) as well as WHO grade and overall survival.

Diffusion-weighted images were analyzed using OsiriX Imaging Software (<http://www.osirix-viewer.com>). ADC measurements were generated by manually drawing 3 nonoverlapping ROIs ranging from 40 to 60 mm² within the region of lowest ADC values within the solid component of each tumor on ADC maps. The ADC value was also calculated from contralateral normal-appearing white matter by drawing a single ROI with a size similar to that of a tumoral ROI. We obtained mean, minimum (min), and maximum (max) ADCs of each tumor, respectively, by averaging the 3 ROIs; and relative ADC ($rADC_{mean}$, $rADC_{min}$, and

$rADC_{max}$) was calculated by dividing the tumor ADC by the ADC of the contralateral normal-appearing white matter.

Statistical Analysis

The Kolmogorov-Smirnov test was used to determine whether the numeric data (age and relative ADC values) for each group were normally distributed. Independent variables (clinical parameter and MR imaging features) were compared using the χ^2 test among different molecular groups. Normally distributed continuous variables (eg, age) were compared using the independent *t* test or ANOVA test, and non-normally distributed continuous variables ($rADC_{mean}$, $rADC_{min}$, and $rADC_{max}$) were compared using the Mann-Whitney *U* test among different molecular groups. The intraobserver reliability of ADC value measuring was tested using intraclass correlation coefficients. The optimal cutoff value of each $rADC$ was obtained from receiver operating characteristic curve analysis when the Youden index reached a maximum. Survival curves were estimated and plotted by the Kaplan-Meier method with log-rank tests to compare Kaplan-Meier curves among groups. Variables were first analyzed by the univariate model. MR imaging features and clinical and molecular parameters (including age, sex, WHO grade, *MGMT* promoter methylation status, *IDH* mutation status, and 1p/19q codeletion status) with statistical significance in univariate analysis ($P < .05$) were entered into a Cox proportional hazards ratio model for multivariate analysis. Statistical significance was defined as $P < .05$ for all tests. The statistical analyses were performed using the statistical software package SPSS 23.0 (IBM, Armonk, New York) and R statistical and computing software, Version 3.3.2 (<http://www.r-project.org>).

RESULTS

A total of 131 (59 [45%] *IDH* wild-type and 72 [55%] *IDH*-mutant) gliomas were included in this study. Of the 72 *IDH*-mutant tumors, 26 (36%) were 1p/19q codeleted and 46 (64%) were non-1p/19q codeleted. Patients in the *IDH* wild-type group (mean, 60 ± 12.2 years) were significantly older than those in the *IDH*-mutant group (mean, 45.1 ± 13.9 years), and patients with *IDH*-mutant, 1p/19q codeleted gliomas (mean, 50.7 ± 13.9 years) were older than those with *IDH*-mutant, non-1p/19q codeleted gliomas (mean, 41.9 ± 12.9 years) ($P = .01$).

Correlation between Conventional MR Imaging Features and Molecular Subtypes

Among the conventional MR imaging characteristics, *IDH* wild-type gliomas were more likely to exhibit enhancement ($P < .001$), >50% enhancing area ($P < .001$), absence of cystic area ($P = .013$), the presence of necrosis ($P < .001$), and the presence of edema ($P < .001$). Within the *IDH*-mutant group, there were no MR imaging characteristics to differentiate 1p/19q codeletion status using the features tested (Table 1).

Correlation between $rADC$ Values and Molecular Subtypes

Median $rADC_{mean}$, $rADC_{min}$, and $rADC_{max}$ values of *IDH* wild-type gliomas were significantly lower than those of *IDH*-mutant gliomas ($P < .001$) (Fig 1 and Table 1). Within the *IDH*-mutant

Table 1: MR imaging features and IDH-mutation and 1p/19q codeletion status

	IDH Wild-Type	IDH-Mutant	P	IDH-Mutant, Non-1p/19q Codeleted	IDH-Mutant, 1p/19q Codeleted	P
MR imaging features (No.) (%)						
T2 signal intensities ^a						
>Gray matter	13 (22.4%)	30 (43.5%)	.012 ^c	21 (48.8%)	9 (34.6%)	.248
>Mixed	45 (77.6%)	39 (56.5%)		22 (51.2%)	17 (65.4%)	
T2 homogeneity ^a						
Homogeneous	6 (10.3%)	14 (20.3%)	.125	10 (23.3%)	4 (15.4%)	.544
Heterogeneous	52 (89.7%)	55 (79.7%)		33 (76.7%)	22 (84.6%)	
Margin						
Well-defined	32 (54.2%)	27 (38.0%)	.065	20 (44.4%)	7 (26.9%)	.143
Mixed	27 (45.8%)	44 (62.0%)		25 (55.6%)	19 (73.1%)	
Edema						
No-to-minimal	20 (33.9%)	57 (79.2%)	<.001 ^c	38 (82.6%)	19 (73.1%)	.339
Mild-to-marked	39 (66.1%)	15 (20.8%)		8 (17.4%)	7 (26.9%)	
Enhancing pattern ^b						
None/minimally enhancing	4 (7.0%)	30 (42.3%)	<.001 ^c	19 (41.3%)	11 (44.0%)	.826
Enhancing	55 (93.2%)	41 (56.9%)		27 (58.7%)	14 (56.0%)	
Proportion of enhancing area ^b						
<50%	14 (23.7%)	63 (88.7%)	<.001 ^c	40 (87.0%)	23 (92.0%)	.704
≥50%	45 (76.3%)	8 (11.3%)		6 (13.0%)	2 (8.0%)	
Cystic area						
Presence	5 (8.5%)	28 (39.4%)	.013 ^c	26 (56.5%)	17 (65.4%)	.461
Absence	54 (91.5%)	43 (59.7%)		20 (43.5%)	9 (34.6%)	
Necrotic area ^b						
Presence	49 (83.1%)	16 (22.5%)	<.001 ^c	38 (82.6%)	17 (68.0%)	.159
Absence	10 (16.9%)	55 (77.5%)		8 (17.4%)	8 (32.0%)	
rADC _{mean} (median)	1.1	1.6	<.001 ^c	1.7	1.5	.071
rADC _{min} (median)	0.9	1.4	<.001 ^c	1.4	1.3	.178
rADC _{max} (median)	1.3	1.8	<.001 ^c	1.9	1.7	.106

^a There were 4 cases lacking T2 MR images.

^b One case in the IDH-mutant group lacked postcontrast studies.

^c Statistically significant ($P < .05$).




FIG 1. Boxplot representation of rADC_{mean} values by glioma IDH genotype.

Table 2: Comparison of overall survival by relative ADC and glioma IDH-mutation status

Variable ^a	Case No.	Survival Time ^b	Coefficient (β)	P Value
IDH mutant, high rADC _{mean}	63	62.0		
vs IDH mutant, low rADC _{mean}			.141	.003
vs IDH wild-type, high rADC _{mean}			.06	<.001
vs IDH wild-type, low rADC _{mean}			.24	<.001
IDH mutant, low rADC _{mean}	9	24.2 (0.0–55.0)		
vs IDH wild-type, high rADC _{mean}			.45	.614
IDH wild-type, high rADC _{mean}	30	17.7 (11.6–23.7)		
vs IDH wild-type, low rADC _{mean}			.25	.041
IDH wild-type, low rADC _{mean}	29	10.4 (4.4–16.4)		
vs IDH mutant, low rADC _{mean}			.51	.179

^a High rADC_{mean}: rADC_{mean} values ≥ 1.08 ; low rADC_{mean}: rADC_{mean} values < 1.08 .

^b Survival time is expressed as median (95% CI) (months).

glioma cohort, rADC values trended lower in 1p/19q codeleted gliomas than in noncodeleted gliomas; however, this trend did not reach statistical significance. Receiver operating characteristic analysis identified an rADC_{mean} of 1.2 as the optimal cutoff value to differentiate IDH wild-type and IDH-mutant gliomas irrespective of WHO grade, with the best combination of sensitivity (81.9%) and specificity (74.6%) and area under the curve (0.790; 95% CI, 0.707–0.869; $P < .001$). In the analysis of intraobserver reliability, the intraclass correlation coefficients indicated a good correlation of the first evaluator (intraclass correlation coefficient, 0.951; 95% CI, 0.829–0.987; $P < .001$) and the second evaluator (intraclass correlation coefficient, 0.926; 95% CI, 0.785–0.975; $P < .001$).

Correlation between rADC and Overall Survival

Median overall survival was 25.4 months (95% CI, 19.0–31.7 months), and overall cumulative survival rates were 79% at 1 year, 59% at 2 years, 42% at 3 years, and 36% at 5 years in all 131 cases. Univariate survival analysis found survival to be significantly related to 6 MR imaging features, including T2 homogeneity, enhancing pattern, enhancing areas, presence of necrosis, rADC_{mean}, and rADC_{min} values, in addition to age, WHO grade, MGMT promoter methylation status, IDH-mutation status, and 1p/19q codeletion status.

Multivariate analysis identified IDH status and rADC_{mean} as the only prognostic factors that independently impacted overall survival after considering the WHO grade, MGMT promoter methylation, and 1p/19q codeletion status and adjusting for patient age. Specifically, IDH-mutant gliomas had significantly longer overall survival (median, 62.0 months) than IDH wild-type gliomas (median, 14.7 months) ($P < .001$ by log-rank test; age-adjusted hazard ratio, 13.5; 95% CI, 4.8–38.4; $P < .001$ by multivariate Cox analysis). Overall cumulative survival rates were 97% at 1 year, 71% at 3 years, and 61% at 5 years in IDH-mutant gliomas, and 76% at 1 year and 13% at 3 years in IDH wild-type gliomas. Gliomas with higher rADC_{mean} had longer overall survival compared with those with lower rADC_{mean} ($P = .001$ by univariate regression; age-adjusted hazard ratio, 0.17; 95% CI, 0.1–0.6; $P = .004$ by multivariate Cox analysis). No statistically significant differences were noted for the remaining MR imaging features, WHO grade, MGMT promoter methylation status, and 1p/19q codeletion status in multivariate analysis.

By means of the area under a time-dependent receiver operating characteristic curve for prediction of survival at 12 months,

the optimal cutoff value of 1.08 for rADC_{mean} could differentiate survival differences within both IDH-mutant and IDH wild-type gliomas. Patients having IDH-mutant gliomas with an rADC_{mean} below the cutoff value of 1.08 had poorer survival than those with an rADC_{mean} above 1.08 ($P < .001$). In addition, median survival associated with IDH-mutant gliomas with a low rADC_{mean} was very poor. Survival time for this group was similar to that of those with IDH wild-type gliomas with either high or low rADC_{mean} (Table 2 and Fig 2). Fi-

nally, the rADC_{mean} cutoff value of 1.08 could also distinguish a survival difference within IDH wild-type gliomas ($P < .041$).

DISCUSSION

Mutations in the IDH genes are among the most important diagnostic and prognostic markers of diffuse gliomas.²⁰ Patients with IDH-mutant gliomas have significantly longer survival compared with those with IDH wild-type gliomas, and management of these 2 molecular subgroups differs significantly. Previous studies have investigated the potential of various conventional and advanced MR imaging characteristics, including perfusion, diffusion tensor, and MR spectroscopy, in identifying genetic subtypes of diffuse gliomas.^{10,21,22} Here, our results indicate that rADC values correlate with IDH mutation status as well as survival in both IDH-mutant and IDH wild-type diffuse gliomas, independent of their WHO grade. Additionally, using rADC, we could identify a particularly poor prognosis subset of IDH-mutant gliomas, with outcomes similar to those in patients IDH wild-type disease. Most important, determining the rADC value is a simple approach that requires no specialized software; hence, our findings potentially have immediate clinical impact.

The mechanism by which IDH-mutant and IDH wild-type gliomas differ in terms of the rADC is not clear; however, it may be related to tumor cellularity. In many previous studies, DWI has shown utility for preoperative grading and outcome of gliomas and for evaluating the response to therapy in patients with glioblastoma.^{13,23–27} ADC values provide quantitative information that reflects Brownian motion of water molecules within a scanned area and are determined by many factors. Mainly, differences in ADC have been attributed to tumor cellularity but also to the presence of necrosis or cysts and water content in interstitial space.^{11,13,28} ADC has been shown to correlate inversely with tumor cellularity on histologic examination, one of the main features of the WHO classification of brain tumors. Our results demonstrate that most IDH-mutant gliomas exhibit higher rADC_{mean} values and MR imaging features accordant with their low-grade features, while most IDH wild-type gliomas show necrosis and a lower rADC_{mean} in solid portions, likely representing higher cellularity, which is associated with higher grade features.

Previous studies have observed an association between ADC and IDH status in gliomas. One study of 37 anaplastic astrocytomas showed that the minimum ADC (cutoff point, 0.95×10^{-3} mm²/s) had acceptable discrimination (area under the curve,




FIG 2. Kaplan-Meier survival curve subclassification for the present study by relative ADC and *IDH* genotype. Censored patients are annotated by a asterisks or plus signs. Results of the analysis are provided in Table 2. *IDHmut* indicates *IDH*-mutant; *IDHwt*, *IDH* wild-type.

0.711; 95% CI, 0.534–0.887) to predict the *IDH* status.²⁹ Another retrospective study of 112 cases by Tan et al³⁰ demonstrated that fractional anisotropy and ADC from diffusion tensor imaging can detect *IDH1* mutation in astrocytomas, with the ratio of ADC_{min} being the best metric for detecting *IDH* mutation, regardless of the WHO grade. We found that $rADC_{mean}$ can differentiate *IDH* wild-type from *IDH*-mutant gliomas with excellent discrimination, regardless of WHO grade. Our study also emphasizes MR imaging and ADC values correlating well with molecular subtype.

The main novel finding of our study is that preoperative $rADC$ values can distinguish favorable and unfavorable prognosis within both *IDH*-mutant and *IDH* wild-type glioma subgroups. While *IDH*-mutant gliomas generally behave less aggressively and have a better prognosis compared with their *IDH* wild-type counterparts, we identified a small subset (12.5%) of *IDH*-mutant gliomas with low $rADC_{mean}$ values and poor overall survival, which was only slightly better (24 months) than that of *IDH* wild-type gliomas but was not statistically significant. Concordantly, a study by Jiao et al³¹ revealed that a small subgroup (11.7%) of patients with *IDH*-mutant gliomas across all grades had a dismal prognosis (median survival of 22 months), more similar to *IDH* wild-type gliomas and glioblastomas in their cohort. These tumors had distinct genetic characteristics, lacking the typical concurrent genetic alterations observed in *IDH*-mutant gliomas. In addition, a recent study of The Cancer Genome Atlas identified a small subset (5.5%) of *IDH*-mutant gliomas with markedly worse survival than other *IDH*-mutant gliomas, and these tumors were associ-

ated with relatively decreased global DNA methylation.³² Together, these data clearly indicate that a subgroup of *IDH*-mutant gliomas behaves as aggressively as their *IDH* wild-type counterparts. Although whether the malignant subgroups across these datasets represent the same biology is unknown, our results suggest that $rADC$ values can potentially identify this aggressively behaving *IDH*-mutant subgroup.

Furthermore, our study highlights how detection of robust imaging-phenotype correlations can be significantly improved by analyzing glioma datasets by molecular subtype rather than by histopathologic classification. We evaluated MR imaging features and prognosis in diffuse gliomas across lower and higher grades in the current study and demonstrated the power of $rADC_{mean}$ to differentiate *IDH*-mutation status and discrete survival subgroups beyond WHO grade. While many previous studies have demonstrated an inverse relationship between ADC and astrocytoma grade,^{12,13,28} other studies have shown substantial overlap of ADC values between high-versus-low-grade gliomas^{3,24,33,34} and no significant differences between grade II versus III³⁵ or grade III versus IV.^{36,37} These observed variations of ADC in predicting tumor grade are likely due to limitations that make the exact histopathologic classification challenging, with high inter-observer variability² and molecular constituent and clinical behavior being likely different in tumors with the same histopathologic grade.^{3,4,38,39} Similar to a recent meta-analysis by Zulfiqar et al,²⁷ which showed that low ADC values correlate independently with poor survival in malignant astrocytomas (grades III and IV),

we found an inverse relationship between $rADC_{mean}$ values and prognosis for both *IDH*-mutant and *IDH* wild-type tumors independent of WHO grade.

One limitation of our study is its retrospective design, which was necessary to include a relatively large number of patients and to correlate with survival, which is relatively long in patients with *IDH* mutation. A second limitation is that the studied patients had been scanned on different MR imaging magnet types, and ADC maps were generated by diffusion-weighted imaging or diffusion tensor imaging of all collected data. However, a previous study has verified that ADC datasets from 3-directional DWI and 6-directional diffusion tensor imaging could be analyzed together.⁴⁰ We calculated the $rADC$ to minimize the differences among absolute ADC values across platforms. Third, the treatment regimen applied to each patient was not available to us in many cases. This issue might have potentially impacted the outcome and survival in each case. However, *IDH* status has been repeatedly shown to be an independent marker of prognosis in independent datasets.⁴ Finally, the ADC value has previously been reported to predict 1p/19q codeletion status, a marker of oligodendroglioma, in lower grade gliomas.^{35,41} In the study by Johnson et al,⁴¹ the ADC values were calculated from sampling both the highest and lowest ADC areas. However, we did not detect a significant correlation between $rADC$ values and 1p/19q codeletion status. Further investigation of the optimal methods of measuring ADC and physiology correlates of ADC values in genetically defined oligodendroglioma is needed.

Our results require independent confirmation, incorporating emerging molecular markers and accounting for different treatment strategies. However, our results expand on and refine the existing correlation between DWI and tumor genetic markers and highlight its potential role as an independent imaging biomarker that can aid in substratification of patients with gliomas, both *IDH*-mutant and *IDH* wild-type. Here, we were able to identify a subset of aggressive *IDH*-mutant gliomas using ADC values easily obtained from common clinical MR images. Ongoing accumulation of tumorigenesis knowledge, together with imaging studies stratified by molecular subgroup rather than histopathologic features, will likely identify additional robust genetic-imaging-clinical phenotype correlations that will improve early detection of clinically meaningful glioma molecular subtypes.

CONCLUSIONS

We demonstrate that ADC values obtained from DWI correlate with *IDH*-mutation status and overall survival in adult diffuse gliomas. *IDH* wild-type gliomas showed low ADC values and poor survival compared with *IDH*-mutant gliomas. Within *IDH*-mutant gliomas, a small subgroup with lower ADC values had dismal survival, similar to that in *IDH* wild-type gliomas. ADC values correlated with survival in patients with *IDH*-mutant and *IDH* wild-type gliomas regardless of WHO grade. Preoperative ADC estimates may corroborate with molecular subtypes as a prognostic marker and potentially enhance risk stratification, especially within *IDH*-mutant gliomas.

2016-00221//C32595GG*; Patents (Planned, Pending or Issued): method for treating high-grade gliomas*. *Money paid to the Institution.

REFERENCES

1. Coons SW, Johnson PC, Scheithauer BW, et al. **Improving diagnostic accuracy and interobserver concordance in the classification and grading of primary gliomas.** *Cancer* 1997;79:1381–93 CrossRef Medline
2. van den Bent MJ. **Interobserver variation of the histopathological diagnosis in clinical trials on glioma: a clinician's perspective.** *Acta Neuropathol* 2010;120:297–304 CrossRef Medline
3. Cancer Genome Atlas Research Network; Brat DJ, Verhaak RG, Aldape KD, et al. **Comprehensive, integrative genomic analysis of diffuse lower-grade gliomas.** *N Engl J Med* 2015;372:2481–98 CrossRef Medline
4. Louis DN, Perry A, Reifenberger G, et al. **The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary.** *Acta Neuropathol* 2016;131:803–20 CrossRef Medline
5. Vignesswaran K, Neill S, Hadjipanayis CG. **Beyond the World Health Organization grading of infiltrating gliomas: advances in the molecular genetics of glioma classification.** *Ann Transl Med* 2015;3:95 CrossRef Medline
6. Turkalp Z, Karamchandani J, Das S. **IDH mutation in glioma: new insights and promises for the future.** *JAMA Neurol* 2014;71:1319–25 CrossRef Medline
7. Patel SH, Poisson LM, Brat DJ, et al. **T2-FLAIR mismatch, an imaging biomarker for IDH and 1p/19q status in lower grade gliomas: a TCGA/TCIA project.** *Clin Cancer Res* 2017;23:6078–85 CrossRef Medline
8. Wang YY, Wang K, Li SW, et al. **Patterns of tumor contrast enhancement predict the prognosis of anaplastic gliomas with IDH1 mutation.** *AJNR Am J Neuroradiol* 2015;36:2023–29 CrossRef Medline
9. Natsumeda M, Igarashi H, Nomura T, et al. **Accumulation of 2-hydroxyglutarate in gliomas correlates with survival: a study by 3.0-Tesla magnetic resonance spectroscopy.** *Acta Neuropathol Commun* 2014;2:158 CrossRef Medline
10. Kickingereder P, Sahm F, Radbruch A, et al. **IDH mutation status is associated with a distinct hypoxia/angiogenesis transcriptome signature which is non-invasively predictable with rCBV imaging in human glioma.** *Sci Rep* 2015;5:16238 CrossRef Medline
11. Gupta RK, Cloughesy TF, Sinha U, et al. **Relationships between choline magnetic resonance spectroscopy, apparent diffusion coefficient and quantitative histopathology in human glioma.** *J Neurooncol* 2000;50:215–26 CrossRef Medline
12. Yamasaki F, Kurisu K, Satoh K, et al. **Apparent diffusion coefficient of human brain tumors at MR imaging.** *Radiology* 2005;235:985–91 CrossRef Medline
13. Lee EJ, Lee SK, Agid R, et al. **Preoperative grading of presumptive low-grade astrocytomas on MR imaging: diagnostic value of minimum apparent diffusion coefficient.** *AJNR Am J Neuroradiol* 2008;29:1872–77 CrossRef Medline
14. Hilario A, Ramos A, Perez-Nuñez A, et al. **The added value of apparent diffusion coefficient to cerebral blood volume in the preoperative grading of diffuse gliomas.** *AJNR Am J Neuroradiol* 2012;33:701–07 CrossRef Medline
15. Zhang L, Min Z, Tang M, et al. **The utility of diffusion MRI with quantitative ADC measurements for differentiating high-grade from low-grade cerebral gliomas: evidence from a meta-analysis.** *J Neurol Sci* 2017;373:9–15 CrossRef Medline
16. TCGA-GBM collection from The Cancer Imaging Archive (TCIA). <https://wiki.cancerimagingarchive.net/display/Public/TCGA-GBM>. Accessed April 01, 2016
17. TCGA-LGG collection from The Cancer Imaging Archive (TCIA). <https://wiki.cancerimagingarchive.net/display/Public/TCGA-LGG>. Accessed April 01, 2016
18. Clark K, Vendt B, Smith K, et al. **The Cancer Imaging Archive (TCIA): maintaining and operating a public information repository.** *J Digit Imaging* 2013;26:1045–57 CrossRef Medline

Disclosures: Dimitris Placantonakis—UNRELATED: Grants/Grants Pending: National Institutes of Health, R01 NS102665, NY State Stem Cell Program - DOH01-STEM5-

19. Wiki for the VASARI feature set The National Cancer Institute Web site. <https://wikicancerimagingarchivenet/display/Public/VASARI+Research+Project>. Updated July 25, 2016
20. Eckel-Passow JE, Lachance DH, Molinaro AM, et al. **Glioma groups based on 1p/19q, IDH, and TERT promoter mutations in tumors.** *N Engl J Med* 2015;372:2499–508 [CrossRef Medline](#)
21. Xiong J, Tan W, Wen J, et al. **Combination of diffusion tensor imaging and conventional MRI correlates with isocitrate dehydrogenase 1/2 mutations but not 1p/19q genotyping in oligodendroglial tumours.** *Eur Radiol* 2016;26:1705–15 [CrossRef Medline](#)
22. Leu K, Ott GA, Lai A, et al. **Perfusion and diffusion MRI signatures in histologic and genetic subtypes of WHO grade II–III diffuse gliomas.** *J Neurooncol* 2017;134:177–88 [CrossRef Medline](#)
23. Murakami R, Hirai T, Kitajima M, et al. **Magnetic resonance imaging of pilocytic astrocytomas: usefulness of the minimum apparent diffusion coefficient (ADC) value for differentiation from high-grade gliomas.** *Acta Radiol* 2008;49:462–67 [CrossRef Medline](#)
24. Higano S, Yun X, Kumabe T, et al. **Malignant astrocytic tumors: clinical importance of apparent diffusion coefficient in prediction of grade and prognosis.** *Radiology* 2006;241:839–46 [CrossRef Medline](#)
25. Murakami R, Sugahara T, Nakamura H, et al. **Malignant supratentorial astrocytoma treated with postoperative radiation therapy: prognostic value of pretreatment quantitative diffusion-weighted MR imaging.** *Radiology* 2007;243:493–99 [CrossRef Medline](#)
26. Yamasaki F, Sugiyama K, Ohtaki M, et al. **Glioblastoma treated with postoperative radio-chemotherapy: prognostic value of apparent diffusion coefficient at MR imaging.** *Eur J Radiol* 2010;73:532–37 [CrossRef Medline](#)
27. Zulficar M, Yousem DM, Lai H. **ADC values and prognosis of malignant astrocytomas: does lower ADC predict a worse prognosis independent of grade of tumor? A meta-analysis.** *AJR Am J Roentgenology* 2013;200:624–29 [CrossRef Medline](#)
28. Sugahara T, Korogi Y, Kochi M, et al. **Usefulness of diffusion-weighted MRI with echo-planar technique in the evaluation of cellularity in gliomas.** *J Magn Reson Imaging* 1999;9:53–60 [CrossRef Medline](#)
29. Wasserman JK, Nicholas G, Yaworski R, et al. **Radiological and pathological features associated with IDH1–R132H mutation status and early mortality in newly diagnosed anaplastic astrocytic tumours.** *PLoS One* 2015;10:e0123890 [CrossRef Medline](#)
30. Tan WL, Huang WY, Yin B, et al. **Can diffusion tensor imaging noninvasively detect IDH1 gene mutations in astroglomas? A retrospective study of 112 cases.** *AJNR Am J Neuroradiol* 2014;35:920–27 [CrossRef Medline](#)
31. Jiao Y, Killela PJ, Reitman ZJ, et al. **Frequent ATRX, CIC, FUBP1 and IDH1 mutations refine the classification of malignant gliomas.** *Oncotarget* 2012;3:709–22 [CrossRef Medline](#)
32. Ceccarelli M, Barthel FP, Malta TM, et al. **Molecular profiling reveals biologically discrete subsets and pathways of progression in diffuse glioma.** *Cell* 2016;164:550–63 [CrossRef Medline](#)
33. Lam WW, Poon WS, Metreweli C. **Diffusion MR imaging in glioma: does it have any role in the pre-operation determination of grading of glioma?** *Clin Radiol* 2002;57:219–25 [CrossRef Medline](#)
34. Murakami R, Hirai T, Sugahara T, et al. **Grading astrocytic tumors by using apparent diffusion coefficient parameters: superiority of a one- versus two-parameter pilot method.** *Radiology* 2009;251:838–45 [CrossRef Medline](#)
35. Khayal IS, Vandenberg SR, Smith KJ, et al. **MRI apparent diffusion coefficient reflects histopathologic subtype, axonal disruption, and tumor fraction in diffuse-type grade II gliomas.** *Neuro Oncol* 2011;13:1192–201 [CrossRef Medline](#)
36. Kang Y, Choi SH, Kim YJ, et al. **Gliomas: histogram analysis of apparent diffusion coefficient maps with standard- or high-b-value diffusion-weighted MR imaging—correlation with tumor grade.** *Radiology* 2011;261:882–90 [CrossRef Medline](#)
37. Kitis O, Altay H, Calli C, et al. **Minimum apparent diffusion coefficients in the evaluation of brain tumors.** *Eur J Radiol* 2005;55:393–400 [Medline](#)
38. Olar A, Wani KM, Alfaro-Munoz KD, et al. **IDH mutation status and role of WHO grade and mitotic index in overall survival in grade II–III diffuse gliomas.** *Acta Neuropathol* 2015;129:585–96 [CrossRef Medline](#)
39. Reuss DE, Mamatjan Y, Schrimpf D, et al. **IDH mutant diffuse and anaplastic astrocytomas have similar age at presentation and little difference in survival: a grading problem for WHO.** *Acta Neuropathol* 2015;129:867–73 [CrossRef Medline](#)
40. Khayal IS, McKnight TR, McGue C, et al. **Apparent diffusion coefficient and fractional anisotropy of newly diagnosed grade II gliomas.** *NMR Biomed* 2009;22:449–55 [CrossRef Medline](#)
41. Johnson DR, Diehn FE, Giannini C, et al. **Genetically defined oligodendroglioma is characterized by indistinct tumor borders at MRI.** *AJNR Am J Neuroradiol* 2017;38:678–84 [CrossRef Medline](#)