

**On-line Table 1: Patient characteristics**

Case/Age (yr)	Sex	Symptoms before ICAS/rePTA	Lesion Location and Stenosis (%)			Stent-System Diameter/Length (mm)	
				on iaDSA <sub>pre</sub>	and iaDSA <sub>post</sub>		
1 (52)	F	Stroke	M1	75	16	Wingspan	3/15
2 (51)	M	Stroke	M1	77	33	Wingspan	2.5/15
3 (51)	M	Stroke	ICA-C3	65	2	Driver	3.5/9
4 (64)	M	Stroke	BA	83	32	Driver	3/9
5 (57)	M	TIA	V4	90	0	Driver	3.5/15
6 (57)	M	TIA	BA	86	20	Pharos	3.5/8
7 (66)	M	Stroke	M1	70	12	Wingspan	3/20
8 (62)	M	None (DWI lesions)	ICA-C3	89	10	Driver	4/9
9 (48)	M	Stroke	M1	76	7	Pharos	2.75/8
10 (44)	F	Stroke	A2	68	25	Wingspan	3/15
11 (45) <sup>a</sup>	F	TIA	A2	77	30	Wingspan	3/15
12 (45) <sup>a</sup>	F	TIA	A2	80	19	Wingspan	3/15
13 (61)	F	Stroke	M1	79	23	Wingspan	3/15
14 (74)	M	Stroke	V4	85	13	Pharos	3.5/8
15 (69)	M	TIA	BA	78	18	Pharos	3/8
16 (75)	M	Stroke	M1	76	33	Wingspan	3/15
17 (56)	F	Stroke	M1	75	10	Wingspan	3/15

<sup>a</sup> Cases after rePTA.**On-line Table 2: Data at follow-up**

Case	Symptoms at Follow-Up	Stenosis (%) on iaDSA	$\Delta$ DSA%	Stenosis (%) on ivACT	Modified Mehran System
1	–	70	54	68	IC
2	–	33	0	70	–
3	–	27	25	32	IB
4	–	54	22	56	IB
5	–	0	0	0	–
6	–	22	2	24	–
7	–	33	21	34	IB
8	–	11	1	12	–
9	–	40	33	43	IA
10	TIA	77	52	79	IB
11	TIA	80	50	78	II
12	–	23	4	26	–
13	–	25	2	27	–
14	–	19	6	24	–
15	–	81	63	78	II
16	–	34	1	37	–
17	–	26	16	29	IB