

On-line Table: Comparison of tonsillar herniation, CSF velocities, and flow durations in patients with Chiari I without and with CAH

	CAH		<i>P</i> Value
	Absent (<i>n</i> = 21, mean [median])	Present (<i>n</i> = 9, mean [median])	
Age (yr)	30.5 ± 15 (31)	26.4 ± 12 (27)	.48
Tonsillar herniation (mm)	10.2 ± 5.9 (7)	14.4 ± 8.4 (10)	.10
Region 1			
Maximum systolic velocity (mm/s)	22.8 ± 16.4 (16.6)	23.6 ± 13.4 (16)	.9
Maximum diastolic velocity (mm/s)	16.4 ± 7.8 (16.5)	20.5 ± 14.2 (17.9)	.3
CSF systole-fraction of cardiac cycle	0.41 ± 0.15 (0.41)	0.28 ± 0.09 (0.25)	.02
CSF diastole-fraction of cardiac cycle	0.59 ± 0.15 (0.59)	0.72 ± 0.09 (0.75)	.02
Region 2			
Maximum systolic velocity (mm/s)	27.05 ± 9.8 (25.1)	29.5 ± 14.4 (30.3)	.87
Maximum diastolic velocity (mm/s)	15.3 ± 5.7 (15.2)	17 ± 10.8 (16.9)	.89
CSF systole-fraction of cardiac cycle	0.49 ± 0.15 (0.5)	0.40 ± 0.11 (0.44)	.16
CSF diastole-fraction of cardiac cycle	0.51 ± 0.15 (0.5)	0.60 ± 0.15 (0.56)	.16