Are your MRI contrast agents cost-effective? Learn more about generic Gadolinium-Based Contrast Agents.





Report of Spectroscopy Findings Not the First

J.R. McConnell

AJNR Am J Neuroradiol published online 1 April 2010 http://www.ajnr.org/content/early/2010/04/01/ajnr.A2081.cit ation

This information is current as of April 19, 2024.

Published April 1, 2010 as 10.3174/ajnr.A2081

Report of Spectroscopy Findings Not the First

Foerster et al¹ were right to report their proton MR spectroscopy findings in children with hepatic encephalopathy but were wrong to claim to be the first. I draw your attention to their article, in which the authors make the statement, "Our study is the first...." This kind of statement adds nothing to the scientific or educational value of the report and is especially concerning because it is not true. In 1995, we used proton MR spectroscopy to assess children with hepatic encephalopathy before and after liver transplantation.² Both articles show that brain spectroscopy is helpful in assessing hepatic encephalopathy in children.

References

- Foerster BR, Conklin LS, Petrou M, et al. Minimal hepatic encephalopathy in children: evaluation with proton MR spectroscopy. AJNR Am J Neuroradiol 2009;30:1610–23. Epub 2009 Jun 9
- 2. McConnell JR, Antonson DL, Ong CS, et al. **Proton spectroscopy of brain** glutamine in acute liver failure. *Hepatology* 1995;22:69–74

J.R. McConnell Greater Regional Medical Center Creston, Iowa

DOI 10.3174/ajnr.A2081