



This information is current as of April 17, 2024.

Follow-Up MR Imaging for Cerebral Sparganosis

S. Wiwanitkit and V. Wiwanitkit

AJNR Am J Neuroradiol published online 25 October 2012 http://www.ajnr.org/content/early/2012/10/25/ajnr.A3391.cit ation

Published October 25, 2012 as 10.3174/ajnr.A3391

Follow-Up MR Imaging for Cerebral Sparganosis

The recent article on follow-up MR imaging for cerebral sparganosis is interesting.¹ Li et al noted, "Migration is a notable feature of cerebral sparganosis." This finding is quite interesting. Due to its basic nature, sparganosis is usually a chronic localized disease. According to a large series from Thailand, no observation of cerebral migration was noted.² The "tunnel" sign is a new finding that was recently reported, but there has been no confirmation on pathology.³ The question is whether the cases are actual sparganosis with confirmed pathology or whether there is any additional co-occurrence of another migrating cerebral parasitic disease such as gnathostomiasis.

References

1. Li YX, Ramsahye H, Yin B, et al. **Migration: a notable feature of cerebral sparganosis on follow-up MR imaging.** *AJNR Am J Neuroradiol* 2012 Aug 2. [Epub ahead of print]

- 2. Wiwanitkit V. A review of human sparganosis in Thailand. Int J Infect Dis 2005;9:312–16
- 3. Song T, Wang WS, Zhou BR, et al. CT and MR characteristics of cerebral sparganosis. AJNR Am J Neuroradiol 2007;28:1700–05

S. Wiwanitkit Wiwanitkit House Bangkok, Thailand V. Wiwanitkit Joseph Ayobabalola University Nigeria, Africa

http://dx.doi.org/10.3174/ajnr.A3391