

**Are your MRI contrast agents cost-effective?**  
Learn more about generic Gadolinium-Based Contrast Agents.



# AJNR

This information is current as  
of April 19, 2024.

## Reply:

M.E. Adin, L. Kleinberg, D. Vaidya, E. Zan, S. Mirbagheri  
and D.M. Yusem

*AJNR Am J Neuroradiol* published online 22 October 2015  
<http://www.ajnr.org/content/early/2015/10/22/ajnr.A4608.citation>

**REPLY:**

Our study commenced in the last quarter of 2013, concluded in May 2014, and was presented at the Symposium Neuro-radiologicum held in Istanbul on September 7–12, 2014. As a result of multipdisciplinary authorship and distant collaborations, the manuscript was finally submitted to the *American Journal of Neuroradiology* in December 2014. At the initial period of our study, there were 2 articles in the literature concluding that hyperintense dentate nuclei (HDN) on T1-weighted MR imaging were secondary to rapidly progressive MS and radiation therapy (RT) effect.<sup>1,2</sup> We, therefore, investigated this entity in a large cohort of irradiated individuals and found no apparent associations between HDN and RT. During the data gathering and after the submission of our manuscript, several studies were published on this topic, mainly arguing the possible association of HDN and repeated performance of gadolinium-enhanced MR imaging.<sup>3,4</sup> McDonald et al<sup>5</sup> were the first to show actual gadolinium deposition in the human brain by using inductively coupled plasma mass spectrometry in postmortem subjects. Although linear gadolinium-based contrast agents were reported to be associated with HDN, the exact mechanism and clinical ramifications remain unclear.<sup>6–8</sup>

T1-weighted FLAIR is a recently described MR imaging sequence.<sup>9</sup> In brain imaging, it is mainly used to better differentiate gray and white matter, owing to its improved contrast difference. In our institution, it is not a part of the standard brain MR imaging protocol and is mainly performed for epilepsy. Because our cohort was obtained from those with a history of RT for underlying tumoral lesions, only a handful of the subjects were imaged with the T1 FLAIR sequence. As far as the authors' qualitative assessment, differentiating HDN from normal dentate nuclei (NDN) on axial T1 FLAIR images was not a matter of debate in any of subjects included in our study (Fig 1). On sagittal T1 FLAIR images, given the enhanced brightness of white matter, qualitative differentiation of faint HDN and NDN could be doubtful, particularly if the reader is not familiar with appearance of HDN. Nevertheless, without specific research on this topic, we do not speculate on the superiority of one technique (T1 FSE, FLAIR, MPRAGE, and so forth) over another in differentiating faint HDN from NDN. From our study cohort, we randomly looked into 7 subjects with NDN who underwent at least 6 contrast-enhanced MR imaging examinations by using linear gadolinium agents (Table). Not a single case was imaged with T1 FLAIR, practically excluding the possibility of HDN being misinterpreted as NDN.

The mechanism of gadolinium retention in the dentate nuclei is unknown, and individual factors contributing to the normal appearance of the dentate nuclei in some patients, despite the large amount of gadolinium administered, remain unclear.

<http://dx.doi.org/10.3174/ajnr.A4608>

**ACKNOWLEDGMENTS**

We thank Kanda et al for their interest in our study and their contribution to the knowledge of gadolinium safety.

**REFERENCES**

- Roccatagliata L, Vuolo L, Bonzano L, et al. Multiple sclerosis: hyperintense dentate nucleus on unenhanced T1-weighted MR images is associated with the secondary progressive subtype. *Radiology* 2009; 251:503–10 CrossRef Medline
- Kasahara S, Miki Y, Kanagaki M, et al. Hyperintense dentate nucleus on unenhanced T1-weighted MR images is associated with a history of brain irradiation. *Radiology* 2011;258:222–28 CrossRef Medline
- Kanda T, Ishii K, Kawaguchi H, et al. High signal intensity in the dentate nucleus and globus pallidus on unenhanced T1-weighted MR images: relationship with increasing cumulative dose of a gadolinium-based contrast material. *Radiology* 2014;270:834–41 CrossRef Medline
- Errante Y, Cirimele V, Mallio CA, et al. Progressive increase of T1 signal intensity of the dentate nucleus on unenhanced magnetic resonance images is associated with cumulative doses of intravenously administered gadodiamide in patients with normal renal function, suggesting dechelation. *Invest Radiol* 2014;49:685–90 CrossRef Medline
- McDonald RJ, McDonald JS, Kallmes DF, et al. Intracranial gadolinium deposition after contrast-enhanced MR imaging. *Radiology* 2015;275:772–82 CrossRef Medline
- Kanda T, Osawa M, Oba H, et al. High signal intensity in dentate nucleus on unenhanced T1-weighted MR images: association with linear versus macrocyclic gadolinium chelate administration. *Radiology* 2015;275:803–09 CrossRef Medline
- Quattrocchi CC, Mallio CA, Errante Y, et al. Gadodiamide and dentate nucleus T1 hyperintensity in patients with meningioma evaluated by multiple follow-up contrast-enhanced magnetic resonance examinations with no systemic interval therapy. *Invest Radiol* 2015; 50:470–72 CrossRef Medline
- Robert P, Lehericy S, Grand S, et al. T1-weighted hypersignal in the deep cerebellar nuclei after repeated administrations of gadolinium-based contrast agents in healthy rats: difference between linear and macrocyclic agents. *Invest Radiol* 2015;50:473–80 CrossRef Medline
- Hori M, Okubo T, Uozumi K, et al. T1-weighted fluid attenuated inversion recovery at low-field strength: a viable alternative for T1-weighted intracranial imaging. *AJNR Am J Neuroradiol* 2003;24: 648–51 Medline

M.E. Adin

Division of Neuroradiology  
The Russell H. Morgan Department of Radiology and Radiological Science  
The Johns Hopkins Medical Institutions  
Baltimore, Maryland

L. Kleinberg

Department of Radiation Oncology and Radiation Molecular Sciences  
The Johns Hopkins Medical Institutions  
Baltimore, Maryland

D. Vaidya

Johns Hopkins Bloomberg School of Public Health  
The Johns Hopkins Medical Institutions  
Baltimore, Maryland


E. Zan

S. Mirbagheri

D.M. Yousem

Division of Neuroradiology

The Russell H. Morgan Department of Radiology and Radiological Science  
The Johns Hopkins Medical Institutions  
Baltimore, Maryland



**FIG 1.** A 16-year-old girl who underwent periodic MR imaging after gadopentetate dimeglumine (Magnevist; Bayer HealthCare Pharmaceuticals, Wayne, New Jersey) administration for operated pilocytic astrocytoma of the optic nerve during 14 years. HDN was clearly visible on both sequences (A, FSE TIWI; TE, 526 ms; TE, 12 ms. B, T1 FLAIR; TR, 1200 ms; TE, 2.46 ms; inversion recovery, 600).

**Number of contrast-enhanced MRIs and amount of gadolinium administration per scan<sup>a</sup>**

No.	Agent Used	Subject 1	Subject 2	Subject 3	Subject 4	Subject 5	Subject 6	Subject 7
1	Magnevist	11 mL	13 mL	13 mL	17 mL	18 mL	3.6 mL	10 mL
2	Magnevist	11 mL	13 mL	13 mL	18 mL	19 mL	4 mL	10 mL
3	Magnevist	12 mL	15 mL	15 mL	20 mL	19 mL	4 mL	11 mL
4	Magnevist	13 mL	15 mL	15 mL	20 mL	19 mL	4 mL	11 mL
5	Magnevist	13 mL	15 mL	15 mL	20 mL	19 mL	4 mL	20 mL
6	Magnevist	13 mL	15 mL	17 mL	20 mL	20 mL	4 mL	
7	Magnevist	13 mL	15 mL	17 mL	20 mL	20 mL	4 mL	
8	Magnevist	15 mL	15 mL	17 mL	20 mL	20 mL		
9	Magnevist	15 mL	15 mL	20 mL	20 mL			
10	Magnevist	15 mL	15 mL	20 mL	20 mL			
11	Magnevist	20 mL	15 mL	20 mL				
12	Magnevist	20 mL	15 mL					
13	Omniscan	13 mL						17 mL
14	Omniscan	13 mL						
15	Omniscan	13 mL						

<sup>a</sup>T1 FLAIR was performed only in subjects 4 and 5. Although the subjects with a history of outside contrast-enhanced MRIs had been excluded, there may have been more contrast administration at outside centers because subjects had long follow-up times for underlying lesions. Therefore, the numbers illustrated in this Table should be considered as the least number of gadolinium administrations.