

On-line Table: Baseline characteristics and risk factors of patients who were and those who were not enrolled in the study^a

Characteristic	Patients with DLB		P	Patients with PDD		P
	Enrolled (n = 42)	Not Enrolled (n = 25)		Enrolled (n = 88)	Not Enrolled (n = 44)	
Sociodemographics						
Age, y	75.1 ± 6.1	74.4 ± 4.9	.53	73.7 ± 7	71.9 ± 5.3	.112
Sex, male	18 (42.9)	11 (44)	.927	32 (36.4)	20 (45.5)	.379
Risk factors						
Hypertension	25 (59.5)	13 (52)	.548	41 (46.6)	19 (43.2)	.711
Diabetes mellitus	16 (38.1)	6 (24)	.235	18 (20.5)	10 (22.7)	.763
Cigarette smoking	4 (9.5)	3 (12)	.749	12 (13.8)	9 (20.5)	.326
Total cholesterol, mg/dL	160.7 ± 30.3	171.5 ± 26.5	.147	175.6 ± 38.3	166.4 ± 41.2	.214
K-MMSE score	19.1 ± 5.1	19.7 ± 6	.659	20.3 ± 4.5	21 ± 3.5	.341
UPDRS-III score	29.2 ± 13.7	23.3 ± 14.3	.196	28.8 ± 15.1	26.4 ± 11.5	.446

Note:—K-MMSE indicates Korean Mini-Mental State Examination; UPDRS-III, Unified Parkinson's Disease Rating Scale Part III.

^a Values are expressed as number (%) of subjects or mean ± SD.