On-line Table: Recurrences

					nary of Recurrences			
Clinically Occult Recurrences (n = 21)								
	No. of Cases	Sex	Mean Age (yr)	Smoking History	Subsite of Primary Tumor	T-Stage	HPV Status	Treatment
NI-RADS 2	8	1 F 7 M	65.6	87.5% (7)	Oral cavity (2) Oropharynx (4) Hypopharynx (0) Larynx (2)	TI (1) T2 (1) T3 (2) T4a (3) T4b (1)	2 (+) 6 (-)	4 Surgery (curative) 3 Chemotherapy/radiation (palliative) 1 Unknown
NI-RADS 3	13	3 F 10 M	65.8	69.2% (9)	Oral cavity (4) Oropharynx (6) Hypopharynx (1) Larynx (2)	T2 (1) T3 (1) T4 (4) T4a (5) T4b (2)	4 (+) 9 (–)	2 Surgery (curative) 7 Chemotherapy/radiation (1 curative, 6 palliative) 2 None 2 Unknown
Total	21	4 F 17 M	65.7	76.2% (16)	Oral cavity (6) Oropharynx (10) Hypopharynx (1) Larynx (4)	T1 (1) T2 (2) T3 (3) T4 (4) T4a (8) T4b (3)	6 (+) 15 (-)	6 Surgery 10 Chemotherapy/ radiation 2 None 3 Unknown
				Symptoma	tic Recurrences (n	= 38)		
NI-RADS 2	15	4 F 11 M	62.9	66.7% (10)	Oral cavity (6) Oropharynx (4) Hypopharynx (2) Larynx (3)	T1 (2) T2 (2) T3 (5) T4a (3) Unknown (3)	1(+)	7 Surgery (curative) 5 Chemotherapy/radiatior (1 curative, 4 palliative) 1 Chemotherapy alone (palliative) 2 Unknown
NI-RADS 3	23	6 F 17 M	64.2	73.9% (17)	Oral cavity (10) Oropharynx (6) Hypopharynx (1) Larynx (5) Nasopharynx (1)	T2 (7) T3 (3) T4 (3) T4a (6) T4b (2) Unknown (2)	4 (+) 1 (-)	2 Surgery (curative) 5 Chemotherapy/radiation (palliative) 9 Chemotherapy alone (palliative) 2 Hospice care 3 Declined further treatment 1 Developed acute infection 1 Unknown
Total	38	10 F 28 M	63.8	76.2% (16)	Oral cavity (16) Oropharynx (10) Hypopharynx (3) Larynx (8) Nasopharynx (1)	TT (2) T2 (9) T3 (8) T4 (3) T4a (9) T4b (2) Unknown (5)	5 (+) 1 (-)	9 Surgery 10 Chemotherapy/ radiation 10 Chemotherapy alone 2 Hospice care 3 Declined further treatment 1 Developed acute infection 3 Unknown