

### Balance and Equilibrium, I: The Vestibule and Semicircular Canals

Joel D. Swartz, David L. Daniels, H. Ric Harnsberger, Katherine A. Shaffer, and Leighton Mark

In this, our second temporal bone installment, we will emphasize the vestibular portion of the labyrinth, that relating to balance and equilibrium. Before proceeding, we must again remind the reader of the basic structure of the labyrinth: an inner membranous labyrinth (endolymphatic) surrounded by an outer osseous labyrinth with an interposed supportive perilymphatic labyrinth. We recommend perusal of the first installment before continuing if there are any uncertainties in this regard.

The *vestibule*, the largest labyrinthine cavity, measures 4 to 6 mm maximal diameter (1–3) (Figs 1–3). The medial wall of the vestibule is unique in that it contains two distinct depressions (Fig 4). Posterosuperiorly lies the *elliptical recess*, where the *utricle* is anchored. The smaller *spherical recess* lies anteroinferiorly and contains the *sacculle*. Between these two concavities lies the *vestibular crest*, a ridge that divides posteriorly into two limbs bounding an additional small depression, the *cochlear recess*, which is the most proximal portion of the cochlear apparatus and leads to the *scala vestibuli*. This area will be discussed in greater detail in subsequent installments.

The *utricle* is firmly attached within the elliptical recess by connective tissue and filaments of the utricular branch of the superior vestibular nerve (4) (Fig 5). The *sacculle* is ovoid and smaller than the *utricle*. It is adherent to the spherical recess by fibrous tissue and the saccular branch of the inferior vestibular nerve. The *utricle* and *sacculle* communicate with each other via the *utriculosaccular duct*. The *utricle* also communicates with the *semicircular canals* and *endolymphatic sinus* (via the *utricular duct*). Additional *saccular communications* include the *endolymphatic sinus* (via the *saccular duct*) and *cochlea* (via the *ductus reuniens*).

The *endolymphatic duct* arises from the *endolymphatic sinus* and passes through the *vestibular aqueduct* of the *osseous labyrinth* to emerge from an aperture along the posterior surface of the *petrous pyramid* as the *endolymphatic sac*.

The *utricle* and *sacculle* are together referred to as the *static labyrinth*, because their function is to detect the position of the head relative to gravity (5–7). They each have a focal concentration of sensory receptors (*maculae*) located at right angles to each other and consisting of ciliated hair cells and tiny crystals of calcium carbonate (*otoliths*) embedded in a gelatinous mass. These *otoliths* respond to *gravitational pull*; therefore, changes in head position distort and stimulate the hair cells.

On the posterior wall of the *utricle* reside five openings for the three *semicircular ducts* (remember that the superior and posterior canals have a common crus). Each *semicircular duct* is less than one third the diameter of the bony housing from which it is separated by *perilymph*. The *osseous superior semicircular canal* is responsible for a ridge known as the *arcuate eminence* along the roof of the *petrous bone* (8). The *lateral semicircular canal* is 30° off the horizontal plane and provides much of the anatomic medial wall of the *attic* and *aditus*. Each of the *semicircular canals* is orthogonal with the others. The superior and lateral canals are innervated by the superior vestibular nerve, the posterior *semicircular canal* by the inferior vestibular nerve (Fig 6).

The *semicircular canals* are together referred to as the *kinetic labyrinth*, because they respond to rotational or angular acceleration (5–8). Therefore, as a unit, the *utricle*, *sacculle* and *semicircular canals* are involved with balance and maintenance of a stable retinal image.


---

From the Department of Radiology, Germantown Hospital and Medical Center, Philadelphia, Pa (J.D.S.), the Section of Neuroradiology, Department of Radiology, Medical College of Wisconsin, Milwaukee (D.L.D., K.A.S., L.M.), and the Section of Neuroradiology, Department of Radiology, University of Utah Medical Center, Salt Lake City (H.R.H.).

Address reprint requests to David L. Daniels, MD, Section of Neuroradiology, Department of Radiology/DH151, Doyne Clinic, Medical College of Wisconsin, 8700 W Wisconsin Ave, Milwaukee, WI 53226.

**Index terms:** Balance; Temporal bone, anatomy; Anatomic moments

AJNR 17:17–21, Jan 1996 0195-6108/96/1701-0017 © American Society of Neuroradiology



1



2



3

Fig 1. Coronal CT image at level of vestibule (V). L indicates lateral semicircular canal; S, superior semicircular canal; P, promontory (basilar turn of cochlea); I, internal auditory canal; and arrows, crista falciformis.

Fig 2. Axial T2-weighted high-resolution fast spin-echo MR image of left temporal bone using phased-array coils. I indicates internal auditory canal; C, cochlea; V, vestibule; single arrow, lateral semicircular canal; and double arrows, posterior semicircular canal.

Fig 3. T2-weighted fast spin-echo MR pulse sequence using phased-array coils and maximum intensity projection technique. Image demonstrates all three semicircular canals emanating from the vestibule as well as the apical, middle, and basilar turns of the cochlea. (Compare with Figure 4.)

Each semicircular canal subtends two thirds of a circle and contains an ampulla for transmission of vestibular nerve fibers. These ampullae are analogous to the maculae of the utricle and saccule in that they contain hair cells. Endolymph flow in response to angular head movement stimulates these hair cells. Each semicircular canal responds to a different rotational axis.

The arterial supply to the vestibule and semicircular canals arises from the labyrinthine branch of the anterior inferior cerebellar artery (4). The posterior, superior, and lateral semicir-

cular canals are supplied via the *anterior vestibular artery*; the capsule and posterior semicircular canal receive their nutrients via the *posterior vestibular artery*.

The thrust of imaging this area is twofold: to identify bony labyrinth detail and to document demineralization, congenital deformity, fractures, and erosive lesions. This is done primarily with computed tomography (CT) (Fig 1). Detailed imaging of the membranous labyrinth is mainly via high-resolution thin-section magnetic resonance (MR) imaging (Figs 2-3). Gadolinium enhancement most often occurs with




Fig 4. Cut-away lateral view of the right osseous labyrinth. In the medial wall of the vestibule are depressions representing the spherical, elliptical, and cochlear recesses, which contain the saccule, the utricle, and the posterior part of the cochlear duct, respectively. The vestibular crest separates the spherical and elliptical recesses (modified from Williams et al [9], Ferner [10], Proctor [11], and Netter and Colacino [12]).




Fig 5. Medial view of the right membranous labyrinth. Demonstrated are the saccule and utricle communicating with each other and with the endolymphatic duct extending to the endolymphatic sac. Also shown are the semicircular ducts communicating with the utricle (modified from Williams et al [9], Ferner [10], Proctor [11], and Netter and Colacino [12]).






Fig 6. Insert view of the membranous labyrinth from Figure 5. Labeled are nerve branches extending from the utricle and ampullae of the superior and lateral semicircular canals, which form the superior vestibular nerve, and branches extending from the saccule and ampulla of the posterior semicircular canal, which form the inferior vestibular nerve.

labyrinthitis. Hyperintensity on nonenhanced T1-weighted images is considered the result of hemorrhage. Obliteration of the membranous labyrinth may be *ossific* (labyrinthine ossifications), best seen with CT, or *fibrous* (earlier stage), perhaps most easily appreciated with newer MR sequences that emphasize thin sections and T2 weighting. The normally uniformly hyperintense signal of the endolymph and perilymph would be altered in this circumstance.

## References

- Bergeron RT. In: Bergeron RT and Som PM, eds. *Head and Neck Imaging, III*. St Louis, Mo: Mosby; 1992:927-944
- Nager GT. *Pathology of the Ear and Temporal Bone*. Baltimore, Md: Williams & Wilkins; 1993:3-49
- Schuknecht H. *Pathology of the Ear*. 2nd ed. Philadelphia, Pa: Lea and Febiger; 1993:45-66
- Hiatt JL, Gartner LP. *Textbook of Head and Neck Anatomy*. 2nd ed. Baltimore, Md: Williams & Wilkins Co; 1987:188
- Montgomery RL. *Head and Neck Anatomy with Clinical Correlations*. New York, NY: McGraw-Hill Book Co; 1981:302-311
- Berkovitz BKB, Moxham BJ. *A Textbook of Head and Neck Anatomy*. St Louis, Mo: Year Book Medical Publishers; 1988:362-388
- Berkovitz BKB, Moxham BJ. *A Textbook of Head and Neck Anatomy*. St Louis, Mo: Year Book Medical Publishers; 1988:471-473
- Daube JR, Reagan TJ, Sandok BA, Westmoreland BF. *Medical Neurosciences: An Approach to Anatomy, Pathology and Physiology by Systems and Levels*. Boston, Mass: Little Brown and Co; 1986:357-365
- Williams PL, Warwick R, Dyson M, Bannister LH, eds. *Gray's Anatomy*. 37th ed. London, England: Churchill Livingstone; 1989: 1229-1234
- Ferner H, ed. *Pernkopf Atlas of Topographical and Applied Human Anatomy*. Baltimore, Md: Urban and Schwarzenberg; 1980: 162, 167, 168
- Proctor B. *Surgical Anatomy of the Ear and Temporal Bone*. New York, NY: Thieme Medical Publishers Inc; 1989:90, 136, 137
- Netter FH, Colacino S, eds. *Atlas of Human Anatomy*. Summit, NJ: Ciba-Geigy Corp; 1989:90