

The **next generation** GBCA
from Guerbet is here

Explore new possibilities >

Guerbet | 

© Guerbet 2024 GUOB220151-A

AJNR

Hearing, II: the retrocochlear auditory pathway.

J D Swartz, D L Daniels, H R Harnsberger, J L Ulmer, K A
Shaffer and L P Mark

AJNR Am J Neuroradiol 1996, 17 (8) 1479-1481
<http://www.ajnr.org/content/17/8/1479.citation>

This information is current as
of September 16, 2024.

Hearing, II: The Retrocochlear Auditory Pathway

Joel D. Swartz, David L. Daniels, H. Ric Harnsberger, John L. Ulmer, Katherine A. Shaffer, and Leighton P. Mark

In contradistinction to the vestibular (Scarpa's) ganglion, which is located at the level of the internal auditory meatus, the cochlear (spiral) ganglion, the primary sensory ganglion for hearing, is located within the bony confines of the cochlea (1, 2). Multiple cochlear fibers converge to form the cochlear nerve, which courses within the anterior-inferior quadrant of the internal auditory canal inferior to the facial nerve and immediately anterior to the inferior vestibular nerve (2, 3) (Figs 1–4). The cochlear nerve then traverses the cerebellopontine angle (usually descending slightly) to synapse in the *dorsal* (higher frequencies) and *ventral* (lower frequencies) nuclei in the upper medulla immediately superficial to the inferior cerebellar peduncle (restiform body) (3–5). For imaging purposes, these nuclei have been referred to by Gebarski et al (6) as the *cochlear nuclear complex* (CNC), which is tubular, 8 mm in length and 3 mm thick. These authors have exquisitely defined landmarks for identification of the CNC, which appears as a modest convexity along the posterolateral surface of the upper medulla bordered by the foramen of Luschka and its accompanying choroid plexus. The choroid plexus extends from the roof of the fourth ventricle to the cerebellopontine angle via the lateral recess. The root entry zone and cerebellar flocculus are also consistent landmarks at this level.

From the CNC, fibers ascend primarily crossed but also uncrossed through the lateral lemniscus within the pons to the inferior colliculi in the midbrain. Fibers from the ventral cochlear nucleus form a structure referred to as the *trap-*

ezoid body. Both crossed and a few of the uncrossed fibers terminate in the superior olivary nuclei. The significance of this latter detour may relate to interaural time differentials (location of sound) or sound dampening (stapedius contraction) (1). From the inferior colliculi in the midbrain, fibers ascend to the medial geniculate bodies in the thalamus and subsequently, via auditory radiations, to the transverse temporal gyrus of Heschl, which resides along the posterior surface of the superior temporal gyrus (7).

The reader should be aware that unilateral retrocochlear sensorineural hearing loss can result only from a lesion of the *cochlear nerve* or *cochlear nuclei* (7). Insults occurring within the more proximal auditory pathway cause bilateral sensorineural hearing loss usually more apparent on the contralateral side. Cortical lesions typically result in an auditory agnosia, an impaired interpretation of sound, rather than a true hearing loss.

Knowledge of the retrocochlear auditory pathway is of importance to the imaging specialist when detailed evaluation of the patient with sensorineural hearing loss is necessary (8). The observer must focus on the anatomy of the entire intraaxial auditory pathway and not just the internal auditory canal and cerebellopontine angle.

Acknowledgment

We thank Hugh Curtin for his enthusiasm and help in preparing the Anatomic Moments on the temporal bone.

From the Department of Radiology, Germantown Hospital and Medical Center, Philadelphia, Pa (J.D.S.), the Section of Neuroradiology, Department of Radiology, Medical College of Wisconsin, Milwaukee (D.L.D., J.L.U., K.A.S., L.P.M.), and the Section of Neuroradiology, Department of Radiology, University of Utah Medical Center, Salt Lake City (H.R.H.).

Address reprint requests to David L. Daniels, MD, Section of Neuroradiology, Department of Radiology, Medical College of Wisconsin, Froedtert Memorial Lutheran Hospital, 9200 W Wisconsin Ave, Milwaukee, WI 53226.

Index terms: Anatomic moments; Hearing; Temporal bone, anatomy

Fig 1. A, Axial T2-weighted fast spin-echo magnetic resonance (MR) image obtained with phased-array coils reveals the cochlea (C) and vestibule (V). The cochlear nerve (upper arrow) and the inferior vestibular nerve (lower arrow) are seen coursing through the inferior portion of the internal auditory canal.

B, Corresponding axial 1-mm-thick computed tomogram.




Fig 2. Sagittal T2-weighted fast spin-echo MR image shows the cochlear nerve (arrow) coursing anteroinferiorly within the fundus of the internal auditory canal.




Fig 3. Axial contrast-enhanced T1-weighted MR image shows choroid plexus enhancement (arrows) within the foramen of Luschka outlining the upper medulla in the region of the cochlear nuclei.




Fig 4. Schematic drawing demonstrates the cochlear nerve extending through the internal auditory canal and cerebellopontine angle cistern to synapse in the dorsal and ventral cochlear nuclei. These nuclei are located lateral to the inferior cerebellar peduncle and form a slight bulge along the posterior lateral surface of the upper medulla. The choroid plexus is located at the posterior aspect of the cerebrospinal fluid-containing foramen of Luschka and also medial to the flocculus (modified from Gebarski et al [6], Harnsberger [9], Ferner [10], DeArmond et al [11], and Netter [12]).

References

1. Wilson-Pauwels L, Akesson EJ, Stewart PA, ed. *Cranial Nerves: Anatomy and Clinical Comments*. Toronto, Canada: BC Decker Inc; 1988:97-112
2. Schuknecht H. *Pathology of the Ear*. 2nd ed, Philadelphia, Pa: Lea & Febiger; 1993:45-66
3. Montgomery RL. *Head and Neck Anatomy with Clinical Correlations*. New York, NY: McGraw-Hill Book Co; 1981:302-311
4. Berkovitz BKB, Moxham BJ. *A Textbook of Head and Neck Anatomy*. St Louis, Mo: Yearbook Medical Publishers; 1988:362-388, 471-473
5. Daube JR, Reagan TJ, Sandok BA, Westmoreland BF. *Medical Neurosciences: An Approach to Anatomy, Pathology and Physiology by Systems and Levels*. New York, NY: Little Brown & Co; 1986:357-365
6. Gebarski SS, Tucci DL, Telian SA. The cochlear nuclear complex: MR location and abnormalities. *AJNR Am J Neuroradiol* 1993;14:1311-1318
7. Armington WG, Harnsberger HR, Smoker WRK, Osborn AG. Normal and diseased acoustic pathway: evaluation with MR imaging. *Radiology* 1988;167:509-515
8. Swartz JD. Sensorineural hearing deficit: a contemporary systematic approach. *Radiographics* 1996;16:561-574
9. Harnsberger HR. *Handbook of Head and Neck Imaging*. St Louis, Mo: Mosby; 1995:527
10. Ferner H, ed. *Pernkopf Atlas of Topographical and Applied Human Anatomy*. Baltimore, Md: Urban & Schwarzenberg; 1980:90, 163
11. DeArmond SJ, Fusco MM, Dewey MM. *Structure of the Human Brain: A Photographic Atlas*. 3rd ed. New York, NY: Oxford University Press; 1989:90, 92
12. Netter FH. *The CIBA Collection of Medical Illustrations*. Vol I: *Nervous System*. West Caldwell, NJ: CIBA; 1986:104, 177