

Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a choice.



FRESENIUS
KABI

[VIEW CATALOG](#)

AJNR

BRAIN IMAGING

Robert I. Grossman

AJNR Am J Neuroradiol 2000, 21 (1) 9-18

<http://www.ajnr.org/content/21/1/9>

This information is current as
of May 9, 2025.

BRAIN IMAGING

The *AJNR* has been at the forefront of the imaging revolution that has taken place over the past 20 years. It has served as one of the primary sources for data critical to the progress of neuroimaging. I have chosen to categorize brain imaging arbitrarily into several sections and summarize some of the advances specifically related to our journal. It has been an amazing 20 years!

Techniques

It was clear quite early in its course that MR imaging could have a profound effect on the depiction of blood flow (1, 2). Indeed, Alvarez et al (3) in 1986 first described the MR imaging appearance of the cessation of blood flow in the internal carotid artery. Among the many comparison studies reported was that of Litt et al (4), who compared conventional angiography with 2D time-of-flight (TOF) angiography. They observed good agreement between the results of MR angiography (MRA) and the standard of reference, conventional angiography. These data began the implementation of what is now the commonly accepted use of MRA as a primary diagnostic tool for extracranial occlusive vascular disease.

Ross et al (5) in 1990 compared intracranial MRA with intra-arterial digital subtraction angiography (DSA) for the evaluation of intracranial aneurysms. They concluded that MRA could reveal intracranial aneurysms as small as 3–4 mm and was a promising method for noninvasive screening of patients at risk for aneurysms.

Lewin and Laub (6) performed comparison studies of TOF techniques for intracranial MRA and suggested tailoring intracranial MRA for maximal diagnostic value. They also provided an analysis of the difficulties associated with these techniques, highlighting problems with slow flow on 3D TOF studies. They confirmed what Ruggieri et al (7) reported; ie, to maximize flow-related enhancement, the imaging volume should be perpendicularly oriented to inflowing blood, with an optimized flip angle and TR. Additionally, to reduce signal loss from flow-induced phase cancellation, constant-velocity flow-compensation gradients, combined with the shortest TE, could minimize intravoxel phase dispersion. Thin sections with small voxels reduced signal loss from phase shifts by limiting the effective range-of-motion-induced phase shifts across the small voxel.

One of the first articles that reported the use of fluid-attenuated inversion recovery (FLAIR) appeared in the *AJNR* in 1992 (8). Harbingers of its current use, the images clearly showed the utility of the pulse sequence. In 1986, LeBihan (9) showed that MR pulse sequences could be sensi-

tized to perfusion and diffusion. One of the earliest and most exciting articles employing diffusion appeared in the *AJNR* in 1990. In it, Moseley et al (10), using an animal model of stroke, reported that diffusion-weighted imaging was more sensitive to early stroke than was conventional T2-weighted imaging. Chien et al (11) extended these observations to patients with stroke and indicated that diffusion imaging may improve tissue specificity and enable differentiation between various types of tissue damage. Tsuruda et al (12) in 1990 implemented the technique to distinguish arachnoid cysts from epidermoid tumors. In commentaries, Henkelman (13) in 1990 noted that diffusion-weighted MR imaging was here to stay, and Fisher and Sotak in 1992 (14) concluded a new era for the very early evaluation of ischemic cerebrovascular disorders appeared to be dawning. These reports correctly predicted the widespread use of diffusion-weighted imaging as an important diagnostic tool.

Functional MR (fMR) imaging represents an important technique for enhancing our understanding of how the brain works. Yetkin et al (15) reported on the application of fMR imaging for noninvasively determining the proximity of eloquent brain to focal lesions. Roberts and Rowley (16) argued that a combination of magnetic source and fMR imaging provided the best results for mapping the sensorimotor cortex. Hedera et al (17) and Lee et al in 1998 (18) reported unexpectedly negative results, failing to identify activated visual cortex and motor cortex, respectively. An editorial by Bryan and Kraut recognized the importance of these negative results, admonishing us to “be skeptical of what [we] see and make nothing of what [we] don’t see” (19).

It was clear rather early that contrast agents would have a major role in the evaluation of brain diseases. Claussen et al (20) in 1985 demonstrated that tumor tissue could be differentiated from peritumoral edema with the use of noncontrast T2-weighted imaging and that gadolinium would increase the potential of MR imaging. Graif et al (21) in the same year reported that comparison of MR with CT showed a greater degree of contrast enhancement on MR images in a group of patients with malignant cerebral tumors.

The technique for determining CBF by xenon-enhanced CT, originally reported by Meyer et al in 1981 (22), has been subsequently discussed and evaluated in a number of articles published in the *AJNR* that sought to improve and validate the technique (23–25).

In 1992, Aoki et al (26) suggested that 3D CT angiography (CTA) could be performed routinely and was helpful for surgical planning by revealing the anatomy of cerebral aneurysms and sur-

rounding structures. Evaluation of CTA in the journal clearly aided in improving the methodology (27, 28).

Stroke and Hemorrhage

The diagnosis of acute stroke prior to MR imaging was based on the clinical presentation in concert with a CT scan that excluded other pathology. Using an acute stroke model, Brant-Zawadzki et al (29) in 1986 reported the potential for increased sensitivity with MR imaging. Bryan et al (30) reported that MR imaging had increased sensitivity for the early detection (within 24 hours) of cerebral infarction and the better definition of the extent of the infarct compared with CT. Yuh et al (31) in the same issue of the *AJNR* argued that signal changes may not be reliable within the first 8 hours and observed that vascular abnormalities, when present, were the most reliable and earliest findings. They also theorized that morphologic changes and early parenchymal enhancement preceded signal changes. They indicated that paramagnetic contrast medium might provide additional data for detection and evaluation of acute ischemia. This confirmed the reports of Elster and Moody (32, 33) identifying arterial enhancement. Crain et al (34) embellished the notion of enhanced MR imaging of cerebral ischemia, reporting a variety of enhancement patterns in the acute phase, which they concluded reflected underlying pathophysiology and could have prognostic significance.

Moody et al (35) found focal abnormalities in terminal arterioles and capillaries among patients and dogs who had undergone cardiopulmonary bypass. They proposed that these findings could be the anatomic correlate of the neurologic deficits or intellectual dysfunction seen in at least 24% of patients after cardiopulmonary bypass. With alkaline phosphatase staining, they found acellular fatty material in the microvasculature of patients who died shortly after cardiopulmonary bypass (36). Steinberg et al (37), using MR and cerebrospinal fluid enzymes, confirmed that brain damage can result from cardiopulmonary bypass. In a commentary, Moody (38) suggested that the heart surgery performed at Stanford, under hypothermia with low blood flow and continuous systemic blood pressure in the 50- to 70-mm Hg range, may decrease the microembolic injury to the brain.

In an elegant scientific investigation, Moody et al (33) in 1990 studied vascular anatomy, employing a refined histochemical technique for alkaline phosphatase that labeled the microvasculature and preserved the background neuropil. They found six different patterns of intraparenchymal afferent blood supply to the brain, and suggested that some of these patterns offered protection to certain brain regions, while leaving others vulnerable in cases of anoxia or hypoperfusion. The work provided an explanation of variations in the brain's response to such injury.

The term *leukoaraiosis* was first used by Hachinski et al (39) in relation to periventricular white matter hypodensities. Braffman et al correlated MR findings of brain specimens with gross and microscopic histopathology in an attempt to provide MR criteria to differentiate lacunar infarction from Virchow-Robin spaces (40) and elucidate the etiology of white matter hyperintensity (41). Hendrie et al (42) noted the correlation between foci of increased intensity on T2-weighted images and aging unrelated to cognitive function or cerebrovascular risk factors. Using xenon CT, Kobari et al (43) in 1990 measured local blood flow in a variety of patient groups and concluded a correlation existed between diffuse cerebral hypoperfusion, cognitive impairment, and leukoaraiosis. Bryan et al (44) in a large patient cohort found that the prevalence of subclinical cerebrovascular disease reflected by MR imaging was of a magnitude greater than clinically suspected, and that disease increased with age and was more prevalent among blacks.

The term *venous congestive encephalopathy* was proposed in the *AJNR* by Willinsky et al in 1994 (45) for patients who present with neurologic deficits caused by venous hypertension. Periventricular venous collagenosis with stenosis or occlusion of deep cerebral veins has been associated with leukoaraiosis (46). Hurst et al (47) found that venous hypertensive encephalopathy secondary to dural arteriovenous fistulas could cause progressive dementia. The journal has published its share of vascular syndrome findings, including familial arteriopathic leukoencephalopathy (48), primary antiphospholipid syndrome (49), and leukoencephalopathy in cerebral amyloid angiopathy (50).

The significance of a hyperdense middle cerebral artery (MCA) on CT has been thoroughly debated in the journal (51–54). Bozzao et al (55) suggested that the appearance of hypodensity revealed by CT soon after embolism onset was strongly predictive of hemorrhagic transformation. In an *AJNR* commentary, Pessin et al (56) argued that the occurrence of hemorrhagic infarction alone does not necessarily imply a serious complication. von Kummer et al (52) expanded this outcome correlate and stated that early hypodensity on CT scans occurring with acute infarction was a predictor of ischemic brain damage. If it encompassed 50% or more of the MCA territory, then there was an 85% positive predictive value for fatal outcome.

Schwartz et al (57) used diffusion imaging to indicate that the edema of hypertensive encephalopathy (as seen in a variety of diseases, including pre-eclampsia-eclampsia, lupus nephritis, and with immunosuppressive drug therapy) has primarily vasogenic origin from lack of autoregulation. They suggested that therapy should aim to lower blood pressure and prevent hemorrhage. Nonaneurysmal perimesencephalic subarachnoid hemorrhage was first reported in 1985 by van Gijn et al (58). In 1991, CT and MR patterns differentiating this entity from aneurysmal rupture was published in the

AJNR (59) and correlated with an excellent prognosis.

The complex MR appearance of hemorrhage generated a great deal of discussion and controversy since the initial publication of its imaging characteristics at 1.5 Tesla (60) and the *AJNR* has been an active forum for much of this discussion. Edelman et al (61) employed gradient-echo imaging to improve detection of susceptibility effects and sensitivity to hemorrhage. Zimmerman et al (62) in 1988 illustrated the imaging appearance of acute hemorrhage at 0.5 Tesla. The role of magnetic susceptibility in the appearance of hypointensity seen on T2-weighted images was corroborated by a histologic biochemical study in rats (63) and in vivo magnetization transfer (MT) measurements (64). Hayman et al (65) suggested the role of hemoglobin immobilization by clot structure or red cell contraction in acute hematoma and disputed the results of in vitro MT and relaxation rate measurements (60). Janick et al (66) refined the understanding of how various oxidation states of intracellular and extracellular hemoglobin and protein concentration contribute to the MR appearance of hemorrhage. Boyko et al (67) described T1 shortening from processes unrelated to hemorrhage (calcification and laminar necrosis).

Infection and Inflammation

Braun et al (68) recognized that finding a ring pattern on unenhanced CT scans could increase specificity of the enhanced images, particularly related to structural lesions such as brain abscess. MR features of pyogenic abscesses were reported in 1989 and included edema, central necrosis, extraparenchymal spread, and peripheral high intensity on T1-weighted images (69). MR imaging was reported to be superior to CT for detecting the full extent of inflammation and assessing the resolution of abscess. The journal reported the difficulties with CT in the diagnosis of subdural empyema (70), but by 1989 the diagnosis could be made early and accurately with MR imaging (71).

By the early 1980s, it was known that HIV was neurotrophic (72). The *AJNR* and other radiologic journals (73–77) reported a plethora of imaging findings in AIDS patients. Flowers et al (78) determined that MR imaging was valuable for evaluating encephalopathy in AIDS patients. Cohen et al (79) in 1992 noted that MR findings are normal to minimally abnormal during early stages of HIV infection, subtle neuropsychologic abnormalities do not correlate with MR, and there may be a prominence of adenoidal tissue in patients without opportunistic infections.

Wehn et al (80) in 1989 first observed dilated perivascular spaces in cryptococcal infections, Tien et al (81) and Mathews et al (82) extended the MR observations regarding this AIDS-related infection, including its lack of enhancement and underestimation of the extent of disease. Coccidioidomycosis was revealed to show widespread cisternal

and cervical meningeal enhancement and ventricular enlargement (83). HIV patients were particularly susceptible to brain abscess.

MT ratio (MTR) measurements proved useful for distinguishing progressive multifocal leukoencephalopathy (PML) from HIV encephalitis (84). PML had lower MTR values—much lower than regions involved with HIV encephalitis. Thallium 201 brain single-photon-emission CT (SPECT) could be employed in the imaging of AIDS patients to differentiate CNS lymphoma from infectious lesions, such as toxoplasma encephalitis (85). Increased intense uptake was associated with lymphoma, whereas toxoplasma encephalitis had no uptake. Kim et al (86) suggested that T1-weighted imaging revealed that tuberculoma had hyperintense and hypointense rims that corresponded to collagenous fibers and inflammatory infiltrate, respectively. The *AJNR* featured imaging findings of many other infectious and inflammatory diseases, including Wegener granulomatosis (87), Lyme disease (88, 89), chronic fatigue syndrome (90), neurosyphilis (91, 92) aspergillosis (93), Whipple's disease (94), St. Louis encephalitis (95), Japanese encephalitis (96), malaria (97) and Rocky Mountain spotted fever (98).

White Matter Disease

Horowitz et al (99) reported that the ovoid lesion on MR images might increase specificity for the diagnosis of MS and that this lesion was the MR correlate of "Dawson's finger." MS lesions could have a diverse appearance, including rings on MR images that are high-intensity with T1-weighted imaging and low-intensity with T2-weighted imaging (100). T2 shortening was reported to be present in MS patients in the thalamus, putamen (101), cortex, and adjacent subcortical white matter (102).

Two investigations reported contrast enhancement decreases over disease duration that reflect clinical classification (103, 104). Grossman et al (105) suggested that proton spectroscopy could lead to better categorization of MS lesions than contrast enhancement could and that demyelination had a longer course than contrast enhancement. Guttmann et al (106) characterized the temporal evolution of MS lesions.

In an animal model of Wallerian degeneration, Lexa et al (107) proved that application of the MTR measure was more sensitive for the early detection of degeneration than was conventional MR imaging and that temporal changes revealed by the MTR corresponded to histologic phases of Wallerian degeneration. The MTR was suggested to be a robust measure (108) and reliable method for determining MS lesion age (109). Hiehle et al (110) drew attention to T1 hypointense lesions and their low MTRs and concluded that, based on MTR data, T1 hypointense lesions represented the most demyelinated MS lesions. Loevner et al (111) confirmed this observation and suggested that T1-weighted imaging might be useful for characterizing MS le-

sions. van Buchem et al (112) combined a computer software program and MT to produce global MTR histograms for estimating whole-brain disease burden in MS. Phillips et al (113) reported that the MTR histogram was a better indicator of global disease burden than T2 was of lesion volume.

Metabolic and Toxic States

Xiong et al (114) reported the MR findings in toluene abuse, including loss of gray-white matter, periventricular white matter abnormality, decreased size of the corpus callosum, and hypointensity intensity in the thalami. Methanol can produce putaminal necrosis and hemorrhage as well as peripheral white matter lesions (115) and other regions involved in severe intoxication include the caudate nucleus, pontine tegmentum, and optic nerves (116). Ethylene glycol toxicity affects the thalamus and pons (117). MR imaging reveals that organic mercury poisoning (Minamata disease) affects the calcarine area, cerebellum, and postcentral gyri, and these regions are responsible for the characteristic manifestation of the disease, including constriction of the visual fields, ataxia, and sensory disturbance (118). Cyclosporin A produced reversible changes in the occipital region (119).

Degenerative Diseases

In 1986, George et al (120) showed that leukoencephalopathy was linked to the aging process and was observed in both "normal" and cognitively impaired individuals who had no other evidence of vascular disease. De Leon et al (121), in a 3-year longitudinal study of patients with Alzheimer's disease, found that changes in ventricular size reflected clinical changes. Holodny et al (122) reported that dilatation of the perihippocampal fissures could be a sensitive and specific marker for distinguishing Alzheimer's disease from normal-pressure hydrocephalus. Bradley et al (123) called attention to the relationship between CBF and CSF circulation and to the association of ischemic periventricular lesions in patients with normal-pressure hydrocephalus and in elderly patients with communicating hydrocephalus. As would be expected, the results of these fresh ideas were debated in the journal (124, 125).

The MR imaging appearance of acute lesions in the Wernicke-Korsakoff syndrome, including reversible involvement of the dorsal medial thalamic nuclei and periaqueductal region (126, 127), were first described in the journal. Kato et al (128) used MT measurements to detect pyramidal tract lesions in amyotrophic lateral sclerosis.

The imaging findings in chronic acquired hepatic failure (increased signal intensity in the basal ganglia, pituitary gland, and mesencephalon surrounding the red nuclei on T1-weighted images) were characterized by Brunberg et al (129). We have even seen the "eye-of-the-tiger" in Hallervorden-Spatz disease (central hypointensity within a hy-

perintense rim surrounded by hypointensity on T2-weighted images in the globus pallidus [130]).

Trauma

Several investigators (131–133) reported imaging characteristics of extracerebral collections secondary to traumatic brain injury. McCluney et al (134) called attention to the position of the cortical veins in differentiating subdural hygroma from atrophy. In a blinded comparison of CT and MR imaging, Orrison et al (135) argued that CT and MR were complementary in the evaluation of acute head trauma. Gentry et al (136) reported that corpus callosal injuries were more frequent than expected, with an associated high incidence of diffuse axonal injury. Mittl et al (64), using T2*-weighted imaging, found evidence of diffuse axonal injury in some patients with mild head injuries in whom CT findings were normal. They suggested that these lesions might be responsible for some aspects of the postconcussive syndrome. Blatter et al (137) performed brain volumetric quantitation in traumatic brain injury, and suggested that these measures might be predictive of cognitive outcome. Bigler et al (138) focused on hippocampal and temporal horn volume and reported that, in the subacute phase after brain trauma, the volume of the temporal horn might correlate with intellectual outcome and that of the hippocampus with verbal memory function.

Pituitary Region

A number of investigators studied the pituitary with CT and MR imaging and described its normal appearance (139, 140), the changes in adolescents and preadolescents (141), and the utility of contrast-enhanced MR imaging for localizing microadenomas (142–145). This small region has had more than its share of reports and controversy. Mark et al (146) in 1984 suggested that high intensity in the posterior sella as shown on T1-weighted images might represent fat. Fujisawa et al (147–149) theorized that the signal intensity was related to the functional status of the hypothalamoneurohypophyseal axis and the signal was secondary to neurosecretory granules. Using phospholipid vesicles, Kucharczyk et al (150) in an elegant experiment modeled the signal intensity and concluded that the MR imaging characteristics could be explained by the phospholipid vesicles. Using fat suppression, Mark et al (146) suggested that the high-intensity signal could have more than one source.

Tien et al (151) proposed that MR imaging could reveal central diabetes insipidus. Moses et al (152) concluded that T1-weighted MR imaging might be able to reveal the difference between central diabetes insipidus (absent posterior pituitary bright spot) and primary polydipsia (bright spot present). Lundin et al (153) reported serial changes in macroprolactinomas resulting from long-term bromo-

criptine therapy, including increasing T2 values over time.

Dynamic MR imaging showed abnormalities of the hypophysial vasculature in lymphocytic hypophysitis (154). Using classical radiologic-pathologic correlation, Sartoretti-Schefer et al (155) were able to separate adamantinous MR findings from squamous-papillary craniopharyngiomas, and Masayuki et al (156) provided MR imaging criteria for the diagnosis of Rathke cleft cysts.

Neoplasms

The detection of metastatic disease has received considerable attention in the journal. In 1990, Sze et al (157) recommended contrast enhancement for the detection of brain metastasis. In 1992, Yuh et al (158) suggested high-dose (0.3 mmol/kg) gadolinium for detection of early or small metastases. Higher contrast doses were judged to be better than delayed imaging with standard contrast doses (159). Sze et al (160) provided confirmation of the beneficial effects of triple dose in cases of equivocal findings or solitary metastasis. In a commentary in the *AJNR*, Ginsberg and Lang (161) argued for postcontrast MT imaging rather than triple-dose gadolinium. Elster and Chen in 1992 (162) concluded that nonenhancing white matter abnormalities have a low probability of representing metastatic disease.

It would be too difficult to enumerate all of the articles on particular brain neoplasms. Suffice to say that the *AJNR* literature runs the spectrum from investigations of astrocytomas (163) to xanthoastrocytomas (164). Proton MR spectroscopy was deemed a reliable technique for grading gliomas when combinations of metabolites were statistically compared (165).

Complications of radiation injury to the brain and the differentiation from recurrent or residual tumor are important clinical issues. The journal published an excellent review of this subject in 1991 (166, 167). Schwartz et al (168) suggested that dual-isotope SPECT with ^{201}Tl and $^{99\text{m}}\text{Tc}$ -HMPAO may be useful for differentiating sites of tumor from radiation changes in patients treated for malignant glioma. In a provocative paper, Ricci et al (169) suggested that flurodeoxyglucose (FDG) positron emission tomography (PET) may not be as useful as previously indicated (170) for differentiating recurrent tumor from radiation necrosis.

Neuropsychiatry

The ability of MR imaging to define structural abnormalities may elucidate behavioral abnormalities. Andreasen in an *AJNR* commentary stressed a new alliance between neuropsychiatry and neuroradiology (171). Degreef et al (172) detected an increased prevalence of a cavum septum pellucidum, cavum vergae, and partial callosal agenesis in schizophrenics, suggesting that these might be an

important substrate in this disorder. Seidenwurm et al (173) studied subjects with extremely violent behavior with FDG PET and found decreased temporal lobe metabolism was correlated with limbic abnormalities seen at electrophysiologic and neuropsychiatric evaluation. Using MR spectroscopy, Gonzalez et al (174) quantified brain lithium. They found variability in brain versus serum levels among patients with bipolar disease. Using fMR imaging, Sunshine et al (175) studied patients with attention deficit disorder. In this preliminary study, they detected additional areas of activity in this patient cohort.

Epilepsy

Imaging findings in postictal patients include transient cerebral swelling and enhancement (167, 176). In a cohort of patients with histologically verified mesial temporal sclerosis, Meiners et al (177) found increased hippocampal signal intensity and decreased gray-white demarcation in the temporal lobe to be the most specific MR imaging features of this lesion. Oppenheim et al (178) observed that complete loss of digitations in the hippocampal head was a sensitive and specific marker of mesial temporal sclerosis. Cheon et al (179) reported that visual assessment was slightly superior to MR volumetry. Using pathologic confirmation as a standard of reference in a large patient cohort with temporal lobe epilepsy, Lee et al (180) found that MR imaging yielded 93% sensitivity and 83% specificity in detecting hippocampal/amygdalar abnormalities. The presence of hippocampal atrophy correlated with the duration of seizures.

The Future

Neuroradiology has been key in advancing our understanding of the pathophysiology of disease and improving the sensitivity to its detection and extent as well as the specificity to particular pathogens. I am struck by how far we have come in the past 20 years in elucidating neurologic diseases. There is, however, still a long road ahead!

MR imaging has become the primary technique for probing the brain and will continue to have that role in the future. Molecular imaging with various MR, SPECT, or PET probes may further our understanding of brain metabolism and function. There is no question that we will have a more profound understanding of how the brain works as functional imaging becomes more refined.

We will clearly move to faster imaging techniques, stronger gradients, and higher field-strength magnets. Resolution will improve, and we may eventually evolve to image true in vivo pathology. High-resolution spectroscopy will be commonplace and that biochemical window will be invaluable as a predictor of outcome. We will use it to assess the effect of disease on the entire

brain. Quantitative methods will readily enable assessment of tumor or disease burden and treatment outcome. These methods will make MR imaging a primary outcome measure in neurologic disease.

It is critical that neuroradiology be substantively involved in the continued development of imaging science. Unless neuroradiology commits itself to performing significant scientific investigation, the specialty could be dismantled. Without scientific progress, there will be no specialty. Other specialties have stronger commitments to basic science and training programs are designed in part to preparing physician scientists. We need to see this as a challenge to our fundamental existence and support all aspects of the science of neuroimaging.

ROBERT I. GROSSMAN, M.D.
Senior Editor

References

1. Axel, L. **Blood flow effects in magnetic resonance imaging.** *AJR Am J Roentgenol* 1984;143:1157-1166
2. Bradley WG Jr, Waluch V. **Blood flow: magnetic resonance imaging.** *Radiology* 1985;154:443-450
3. Alvarez O, Edwards JH, Hyman RA. **MR recognition of internal carotid artery occlusion.** *AJNR Am J Neuroradiol* 1986;7:359-360
4. Litt AW, Eidelman EM, Pinto RS, et al. **Diagnosis of carotid artery stenosis: comparison of 2DFT time-of-flight MR angiography with contrast angiography in 50 patients.** *AJNR Am J Neuroradiol* 1991;12:149-154
5. Ross JS, Masaryk TJ, Modic MT, Ruggieri PM, Haacke EM, Selman WR. **Intracranial aneurysms: evaluation by MR angiography.** *AJNR Am J Neuroradiol* 1990;11:449-455
6. Lewin JS, Laub G. **Intracranial MR angiography: a direct comparison of three time-of-flight techniques [see comments].** *AJNR Am J Neuroradiol* 1991;12:1133-1139
7. Ruggieri PM, Laub GA, Masaryk TJ, Modic MT. **Intracranial circulation: pulse-sequence considerations in three-dimensional (volume) MR angiography.** *Radiology* 1989;171:785-791
8. De Coene B, Hajnal JV, Gatehouse P, et al. **MR of the brain using fluid-attenuated inversion recovery (FLAIR) pulse sequences.** *AJNR Am J Neuroradiol* 1992;13:1555-1564
9. Le Bihan D, Breton E, Lallemand D, Grenier P, Cabanis E, Laval-Jeantet M. **MR imaging of intravoxel incoherent motions: application to diffusion and perfusion in neurologic disorders.** *Radiology* 1986;161:401-407
10. Moseley ME, Kucharczyk J, Mintorovitch J, et al. **Diffusion-weighted MR imaging of acute stroke: correlation with T2-weighted and magnetic susceptibility-enhanced MR imaging in cats.** *AJNR Am J Neuroradiol* 1990;11:423-429
11. Chien D, Kwong KK, Gress DR, Buonanno FS, Buxton RB, Rosen BR. **MR diffusion imaging of cerebral infarction in humans.** *AJNR Am J Neuroradiol* 1992;13:1097-1102; discussion 1103-1105
12. Tsuruda JS, Chew WM, Moseley ME, Norman D. **Diffusion-weighted MR imaging of the brain: value of differentiating between extraaxial cysts and epidermoid tumors.** *AJNR Am J Neuroradiol* 1990;11:925-931; discussion 932-934
13. Henkelman R. **Diffusion-weighted MR imaging: a useful adjunct to clinical diagnosis or a scientific curiosity?** *AJNR Am J Neuroradiol* 1990;11:932-934
14. Fisher M, Sotak C. **Diffusion-weighted MR imaging and ischemic stroke.** *AJNR Am J Neuroradiol* 1992;13:1103-1105
15. Yetkin FZ, Papke RA, Mark LP, Daniels DL, Mueller WM, Haughton VM. **Location of the sensorimotor cortex: functional and conventional MR compared.** *AJNR Am J Neuroradiol* 1995;16:2109-2113
16. Roberts TP, Rowley HA. **Mapping of the sensorimotor cortex: functional MR and magnetic source imaging.** *AJNR Am J Neuroradiol* 1997;18:871-880
17. Hedera P, Wu D, Collins S, et al. **Sex and electroencephalographic synchronization after photic stimulation predict signal changes in the visual cortex on functional MR images [see comments].** *AJNR Am J Neuroradiol* 1998;19:853-857
18. Lee D, Port NL, Kruse W, Georgopoulos AP. **Variability and correlated noise in the discharge of neurons in motor and parietal areas of the primate cortex.** *J Neurosci* 1998;18:1161-1170
19. Bryan RN, Kraut M. **Functional magnetic resonance imaging: you get what you (barely) see [editorial; comment].** *AJNR Am J Neuroradiol* 1998;19:991-992
20. Claussen C, Laniado M, Kazner E, Schormer W, Felix R. **Application of contrast agents in CT and MRI (NMR): their potential in imaging of brain tumors.** *Neuroradiology* 1985;27:164-171
21. Graif M, Bydder GM, Steiner RE, Niendorf P, Thomas DG, Young IR. **Contrast-enhanced MR imaging of malignant brain tumors.** *AJNR Am J Neuroradiol* 1985;6:855-862
22. Meyer JS, Hayman LA, Amano T, et al. **Mapping local blood flow of human brain by CT scanning during stable xenon inhalation.** *Stroke* 1981;12:426-436
23. Giller CA, Purdy P, Lindstrom WW. **Effects of inhaled stable xenon on cerebral blood flow velocity.** *AJNR Am J Neuroradiol* 1990;11:177-182
24. Good WE, Gur D. **Xenon-enhanced CT of the brain: effect of flow activation on derived cerebral blood flow measurements [see comments].** *AJNR Am J Neuroradiol* 1991;12:83-85
25. Kashiwagi S, Yamashita T, Nakano S, et al. **The wash-in/wash-out protocol in stable xenon CT cerebral blood flow studies.** *AJNR Am J Neuroradiol* 1992;13:49-53
26. Aoki S, Sasaki Y, Machida T, Ohkubo T, Minami M. **Cerebral aneurysms: detection and delineation using 3-D-CT angiography.** *AJNR Am J Neuroradiol* 1992;13:1115-1120
27. Dix JE, Evans AJ, Kallmes DE, Sobel AH, Phillips CD. **Accuracy and precision of CT angiography in a model of carotid artery bifurcation stenosis.** *AJNR Am J Neuroradiol* 1997;18:409-415
28. Wise SW, Hopper KD, Schwartz TA, Ten Have TR, Kasales CJ. **Technical factors of CT angiography studied with a carotid artery phantom.** *AJNR Am J Neuroradiol* 1997;18:401-408
29. Brant-Zawadzki M, Pereira B, Weinstein P, et al. **MR imaging of acute experimental ischemia in cats.** *AJNR Am J Neuroradiol* 1986;7:7-11
30. Bryan RN, Levy LM, Whitlow WD, Killian JM, Preziosi TJ, Rosario JA. **Diagnosis of acute cerebral infarction: comparison of CT and MR imaging [see comments].** *AJNR Am J Neuroradiol* 1991;12:611-620
31. Yuh WT, Crain MR, Loes DJ, Greene GM, Ryals TJ, Sato Y. **MR imaging of cerebral ischemia: findings in the first 24 hours [see comments].** *AJNR Am J Neuroradiol* 1991;12:621-629
32. Elster AD, Moody DM. **Early cerebral infarction: gadopentetate dimeglumine enhancement [see comments].** *Radiology* 1990;177:627-632
33. Moody DM, Bell MA, Challa VR. **Features of the cerebral vascular pattern that predict vulnerability to perfusion or oxygenation deficiency: an anatomic study.** *AJNR Am J Neuroradiol* 1990;11:431-439
34. Crain MR, Yuh WT, Greene GM, et al. **Cerebral ischemia: evaluation with contrast-enhanced MR imaging.** *AJNR Am J Neuroradiol* 1991;12:631-639
35. Moody DM, Bell MA, Challa VR, Johnston WE, Prough DS. **Brain microemboli during cardiac surgery or aortography [see comments].** *Ann Neurol* 1990;28:477-486
36. Moody DM, Brown WR, Challa VR, Stump DA, Reboussin DM, Legault C. **Brain microemboli associated with cardiopulmonary bypass: a histologic and magnetic resonance imaging study.** *Ann Thorac Surg* 1995;59:1304-1307
37. Steinberg GK, De La Paz R, Mitchell RS, Bell TE, Albers GW. **MR and cerebrospinal fluid enzymes as sensitive indicators of subclinical cerebral injury after open-heart valve replacement surgery.** *AJNR Am J Neuroradiol* 1996;17:205-212; discussion 213-215

38. Moody D. **Is There a Role for MR in the Development of Safer Cardiac Surgery?** *AJNR Am J Neuroradiol* 1996;17:213-215
39. Hachinski VC, Potter P, Merskey H. **Leuko-araiosis.** *Arch Neurol* 1987;44:21-23
40. Braffman BH, Zimmerman RA, Trojanowski JQ, Gonatas NK, Hickey WE, Schlaepfer WW. **Brain MR: pathologic correlation with gross and histopathology. 1. Lacunar infarction and Virchow-Robin spaces.** *AJR. Am J Roentgenol* 1988;151:551-558
41. Braffman BH, Zimmerman RA, Trojanowski JQ, Gonatas NK, Hickey WE, Schlaepfer WW. **Brain MR: pathologic correlation with gross and histopathology. 2. Hyperintense white-matter foci in the elderly correlation with gross and histopathology.** *AJR. Am J Roentgenol* 1988;151:559-566
42. Hendrie HC, Farlow MR, Austrom MG, Edwards MK, Williams MA. **Foci of increased T2 signal intensity on brain MR scans of healthy elderly subjects.** *AJNR Am J Neuroradiol* 1989;10:703-707
43. Kobari M, Meyer JS, Ichijo M, Oravez WT. **Leukoaraiosis: correlation of MR and CT findings with blood flow, atrophy, and cognition.** *AJNR Am J Neuroradiol* 1990;11:273-281
44. Bryan R, Cai J, Burke G, et al. **Prevalence and anatomic characteristics of infarct-like lesions on MR images of middle-aged adults: the atherosclerosis risk in communities study.** *AJNR Am J Neuroradiol* 1999;20:1273-1280
45. Willinsky R, Terbrugge K, Montanera W, Mikulis D, Wallace MC. **Venous congestion: an MR finding in dural arteriovenous malformations with cortical venous drainage.** *AJNR Am J Neuroradiol* 1994;15:1501-1507
46. Moody DM, Brown WR, Challa VR, Anderson RL. **Periventricular venous collagenosis: association with leukoaraiosis.** *Radiology* 1995;194:469-476
47. Hurst RW, Bagley LJ, Galetta S, et al. **Dementia resulting from dural arteriovenous fistulas: the pathologic findings of venous hypertensive encephalopathy [see comments].** *AJNR Am J Neuroradiol* 1998;19:1267-1273
48. Glusker P, Horoupian DS, Lane B. **Familial arteriopathic leukoencephalopathy: imaging and neuropathologic findings.** *AJNR Am J Neuroradiol* 1998;19:469-475
49. Pulpeiro JR, Cortes JA, Macarron J, Perez R, Bal M. **MR findings in primary antiphospholipid syndrome [see comments].** *AJNR Am J Neuroradiol* 1991;12:452-453
50. Loes DJ, Biller J, Yuh WT, et al. **Leukoencephalopathy in cerebral amyloid angiopathy: MR imaging in four cases.** *AJNR Am J Neuroradiol* 1990;11:485-488
51. Tomsick TA, Brott TG, Chambers AA, et al. **Hyperdense middle cerebral artery sign on CT: efficacy in detecting middle cerebral artery thrombosis.** *AJNR Am J Neuroradiol* 1990;11:473-477
52. von Kummer R, Meyding-Lamade U, Forsting M, et al. **Sensitivity and prognostic value of early CT in occlusion of the middle cerebral artery trunk.** *AJNR Am J Neuroradiol* 1994;15:9-15; discussion 16-18
53. Tomsick T, Brott T, Barsan W, Broderick J, Haley EC, Spilker J. **Thrombus localization with emergency cerebral CT.** *AJNR Am J Neuroradiol* 1992;13:257-263
54. Rauch RA, Bazan CD, Larsson EM, Jinkins JR. **Hyperdense middle cerebral arteries identified on CT as a false sign of vascular occlusion.** *AJNR Am J Neuroradiol* 1993;14:669-673
55. Bozzao L, Angeloni U, Bastianello S, Fantozzi LM, Pierallini A, Fieschi C. **Early angiographic and CT findings in patients with hemorrhagic infarction in the distribution of the middle cerebral artery [see comments].** *AJNR Am J Neuroradiol* 1991;12:1115-1121
56. Pessin MS, Teal PA, Caplan LR. **Hemorrhagic infarction: guilt by association? [comment].** *AJNR Am J Neuroradiol* 1991;12:1123-1126
57. Schwartz RB, Mulkern RV, Gudbjartsson H, Jolesz F. **Diffusion-weighted MR imaging in hypertensive encephalopathy: clues to pathogenesis.** *AJNR Am J Neuroradiol* 1998;19:859-862
58. van Gijn J, van Dongen KJ, Vermeulen M, Hijdra A. **Perimesencephalic hemorrhage: a nonaneurysmal and benign form of subarachnoid hemorrhage.** *Neurology* 1985;35:493-497
59. Rinkel GJ, Wijdevicks EF, Vermeulen M, et al. **Nonaneurysmal perimesencephalic subarachnoid hemorrhage: CT and MR patterns that differ from aneurysmal rupture.** *AJNR Am J Neuroradiol* 1991;12:829-834
60. Gomori JM, Grossman RI, Goldberg HI, Zimmerman RA, Bilaniuk LT. **Intracranial hematomas: imaging by high-field MR.** *Radiology* 1985;157:87-93
61. Edelman RR, Johnson K, Buxton R, et al. **MR of hemorrhage: a new approach.** *AJNR Am J Neuroradiol* 1986;7:751-756
62. Zimmerman RD, Heier LA, Snow BR, Liu DP, Kelly AB, Deck MD. **Acute intracranial hemorrhage: intensity changes on sequential MR scans at 0.5 T.** *AJR. Am J Roentgenol* 1988;150:651-661
63. Thulborn KR, Sorensen AG, Kowall NW, et al. **The role of ferritin and hemosiderin in the MR appearance of cerebral hemorrhage: a histopathologic biochemical study in rats [see comments].** *AJNR Am J Neuroradiol* 1990;11:291-297
64. Mittl RL, Grossman RI, Hiehle JF, et al. **Prevalence of MR evidence of diffuse axonal injury in patients with mild head injury and normal head CT findings.** *AJNR Am J Neuroradiol* 1994;15:1583-1589
65. Hayman LA, Taber KH, Ford JJ, et al. **Effect of clot formation and retraction on spin-echo MR images of blood: an in vitro study.** *AJNR Am J Neuroradiol* 1989;10:1155-1158
66. Janick PA, Hackney DB, Grossman RI, Asakura T. **MR imaging of various oxidation states of intracellular and extracellular hemoglobin [see comments].** *AJNR Am J Neuroradiol* 1991;12:891-897
67. Boyko OB, Burger PC, Shelburne JD, Ingram P. **Non-heme mechanisms for T1 shortening: pathologic, CT, and MR elucidation.** *AJNR Am J Neuroradiol* 1992;13:1439-1445
68. Braun IF, Chambers E, Leeds NE, Zimmerman RD. **The value of unenhanced scans in differentiating lesions producing ring enhancement.** *AJNR Am J Neuroradiol* 1982;3:643-647
69. Haimes AB, Zimmerman RD, Morgello S, et al. **MR imaging of brain abscesses.** *AJR. Am J Roentgenol* 1989;152:1073-1085
70. Sandhu V, Handel S, Pinto R, Glass T. **Neuroradiologic diagnosis of subdural empyema and CT limitations.** *AJNR Am J Neuroradiol* 1980;1:39-44
71. Weingarten K, Zimmerman RD, Becker RD, Heier LA, Haimes AB, Deck MD. **Subdural and epidural empyemas: MR imaging.** *AJR. Am J Roentgenol* 1989;152:615-621
72. Levy JA, Hoffman AD, Kramer SM, Landis JA, Shimabukuro JM, Oshiro LS. **Isolation of lymphocytotropic retroviruses from San Francisco patients with AIDS.** *Science* 1984;225:840-842
73. Kelly WM, Brant-Zawadzki M. **Acquired immunodeficiency syndrome: neuroradiologic findings.** *Radiology* 1983;149:485-491
74. Whelan MA, Kricheff II, Handler M, et al. **Acquired immunodeficiency syndrome: cerebral computed tomographic manifestations.** *Radiology* 1983;149:477-484
75. Post MJ, Sheldon JJ, Hensley GT, et al. **Central nervous system disease in acquired immunodeficiency syndrome: prospective correlation using CT, MR imaging, and pathologic studies.** *Radiology* 1986;158:141-148
76. Levy RM, Rosenbloom S, Perrett LV. **Neuroradiologic findings in AIDS: a review of 200 cases.** *AJR. Am J Roentgenol* 1986;147:977-983
77. Post MJ, Tate LG, Quencer RM, et al. **CT, MR, and pathology in HIV encephalitis and meningitis.** *AJR Am J Roentgenol* 1988;151:373-380
78. Flowers CH, Mafee ME, Crowell R, et al. **Encephalopathy in AIDS patients: evaluation with MR imaging.** *AJNR Am J Neuroradiol* 1990;11:1235-1245
79. Cohen WA, Maravilla KR, Gerlach R, et al. **Prospective cerebral MR study of HIV seropositive and seronegative men: correlation of MR findings with neurologic, neuropsychologic, and cerebrospinal fluid analysis.** *AJNR Am J Neuroradiol* 1992;13:1231-1240
80. Wehn SM, Heinz ER, Burger PC, Boyko OB. **Dilated Virchow-Robin spaces in cryptococcal meningitis associated with AIDS: CT and MR findings.** *J Comput Assist Tomogr* 1989;13:756-762
81. Tien RD, Chu PK, Hesselink JR, Duberg A, Wiley C. **Intracranial cryptococcosis in immunocompromised patients: CT and MR findings in 29 cases.** *AJNR Am J Neuroradiol* 1991;12:283-289
82. Mathews VP, Alo PL, Glass JD, Kumar AJ, McArthur JC. **AIDS-related CNS cryptococcosis: radiologic-pathologic correlation.** *AJNR Am J Neuroradiol* 1992;13:1477-1486
83. Wrobel CJ, Meyer S, Johnson RH, Hesselink JR. **MR findings in acute and chronic coccidioidomycosis meningitis.** *AJNR Am J Neuroradiol* 1992;13:1241-1245
84. Dousset V, Armand JP, Lacoste D, et al. **Magnetization transfer study of HIV encephalitis and progressive multifocal leukoencephalopathy.** *Groupe d'Epidemiologie Clinique du SIDA en Aquitaine. AJNR Am J Neuroradiol* 1997;18:895-901

85. Ruiz A, Ganz WI, Post MJ, et al. Use of thallium-201 brain SPECT to differentiate cerebral lymphoma from toxoplasma encephalitis in AIDS patients [see comments]. *AJNR Am J Neuroradiol* 1994;15:1885-1894
86. Kim TK, Chang KH, Kim CJ, Goo JM, Kook MC, Han MH. Intracranial tuberculoma: comparison of MR with pathologic findings. *AJNR Am J Neuroradiol* 1995;16:1903-1908
87. Provenzale JM, Allen NB. Wegener granulomatosis: CT and MR findings. *AJNR Am J Neuroradiol* 1996;17:785-792
88. Rafto SE, Milton WJ, Galetta SL, Grossman RI. Biopsy-confirmed CNS Lyme disease: MR appearance at 1.5 T. *AJNR Am J Neuroradiol* 1990;11:482-484
89. Fernandez RE, Rothberg M, Ferencz G, Wujack D. Lyme disease of the CNS: MR imaging findings in 14 cases. *AJNR Am J Neuroradiol* 1990;11:479-481
90. Greco A, Tannock C, Brostoff J, Costa DC. Brain MR in chronic fatigue syndrome. *AJNR Am J Neuroradiol* 1997;18:1265-1269
91. Agrons GA, Han SS, Husson MA, Simeone F. MR imaging of cerebral gumma. *AJNR Am J Neuroradiol* 1991;12:80-81
92. Brightbill TC, Ihmeidan IH, Post MJ, Berger JR, Katz DA. Neurosyphilis in HIV-positive and HIV-negative patients: neuroimaging findings. *AJNR Am J Neuroradiol* 1995;16:703-711
93. Cox J, Murtagh FR, Wilfong A, Brenner J. Cerebral aspergillosis: MR imaging and histopathologic correlation [see comments]. *AJNR Am J Neuroradiol* 1992;13:1489-1492
94. Schnider P, Trattnig S, Kollegger H, Auff E. MR of cerebral Whipple disease. *AJNR Am J Neuroradiol* 1995;16:1328-1329
95. Cema F, Mehrad B, Luby J, Fleckenstein BD and J. St. Louis encephalitis and the substantia nigra: MR imaging evaluation. *AJNR Am J Neuroradiol* 1999;20:1281-1283
96. Kimura K, Dosaka A, Hashimoto Y, Yasunaga T, Uchino M, Ando M. Single-photon emission CT findings in acute Japanese encephalitis. *AJNR Am J Neuroradiol* 1997;18:465-469
97. Cordoliani YS, Sarrazin JL, Felten D, Caumes E, Leveque C, Fisch A. MR of cerebral malaria. *AJNR Am J Neuroradiol* 1998;19:871-874
98. Bonawitz C, Castillo M, Mukherji SK. Comparison of CT and MR features with clinical outcome inpatients with Rocky Mountain spotted fever. *AJNR Am J Neuroradiol* 1997;18:459-464
99. Horowitz AL, Kaplan RD, Grewe G, White RT, Salberg LM. The ovoid lesion: a new MR observation in patients with multiple sclerosis. *AJNR Am J Neuroradiol* 1989;10:303-305
100. Powell T, Sussman JG, Davies-Jones GA. MR imaging in acute multiple sclerosis: ring like appearance in plaques suggesting the presence of paramagnetic free radicals [see comments]. *AJNR Am J Neuroradiol* 1992;13:1544-1546
101. Drayer B, Burger P, Hurwitz B, Dawson D, Cain J. Reduced signal intensity on MR images of thalamus and putamen in multiple sclerosis: increased iron content? *AJR Am J Roentgenol* 1987;149:357-363
102. Russo C, Smoker WR, Kubal W. Cortical and subcortical T2 shortening in multiple sclerosis. *AJNR Am J Neuroradiol* 1997;18:124-126
103. Miki Y, Grossman RI, Udupa JK, et al. Computer-assisted quantitation of enhancing lesions in multiple sclerosis: correlation with clinical classification. *AJNR Am J Neuroradiol* 1997;18:705-710
104. Filippi M, Rossi P, Campi A, Colombo B, Pereira C, Comi G. Serial contrast-enhanced MR in patients with multiple sclerosis and varying levels of disability. *AJNR Am J Neuroradiol* 1997;18:1549-1556
105. Grossman RI, Lenkinski RE, Ramer KN, Gonzalez-Scarano F, Cohen JA. MR proton spectroscopy in multiple sclerosis. *AJNR Am J Neuroradiol* 1992;13:1535-1543
106. Guttmann CR, Ahn SS, Hsu L, Kikinis R, Jolesz FA. The evolution of multiple sclerosis lesions on serial MR. *AJNR Am J Neuroradiol* 1995;16:1481-1491
107. Lexa FJ, Grossman RI, Rosenquist AC. Dyke Award paper. MR of wallerian degeneration in the feline visual system: characterization by magnetization transfer rate with histopathologic correlation. *AJNR Am J Neuroradiol* 1994;15:201-212
108. Mehta RC, GB Pike GB, Enzmann DR. Magnetization transfer MR of the normal adult brain. *AJNR Am J Neuroradiol* 1995;16:2085-2091
109. Torniak MM, Rosenblum JD, Prager JM, Metz CE. Magnetization transfer: a potential method to determine the age of multiple sclerosis lesions. *AJNR Am J Neuroradiol* 1994;15:1569-1574
110. Leehle JF, Jr, Grossman RI, Ramer KN, Gonzalez-Scarano F, Cohen JA. Magnetization transfer effects in MR-detected multiple sclerosis lesions: comparison with gadolinium-enhanced spin-echo images and nonenhanced T1-weighted images. *AJNR Am J Neuroradiol* 1995;16:69-77
111. Loevner LA, Grossman RI, McGowan JC, Ramer KN, Cohen JA. Characterization of multiple sclerosis plaques with T1-weighted MR and quantitative magnetization transfer. *AJNR Am J Neuroradiol* 1995;16:1473-1479
112. van Buchem MA, Udupa JK, McGowan JC, et al. Global volumetric Estimation of disease burden in multiple sclerosis based on magnetization transfer imaging. *AJNR Am J Neuroradiol* 1997;18:1287-1290
113. Phillips MD, Grossman RI, Miki Y, et al. Comparison of T2 lesion volume and magnetization transfer ratio histogram analysis and of atrophy and measures of lesion burden in patients with multiple sclerosis. *AJNR Am J Neuroradiol* 1998;19:1055-1060
114. Xiong L, Matthes JD, Li J, Jinkins JR. MR imaging of "spray-heads": toluene abuse via aerosol paint inhalation. *AJNR Am J Neuroradiol* 1993;14:1195-1199
115. Rubinstein D, Escott E, Kelly JP. Methanol intoxication with putaminal and white matter necrosis: MR and CT findings. *AJNR Am J Neuroradiol* 1995;16:1492-1494
116. Gaul HP, Wallace CJ, Auer RN, Fong TC. MR findings in methanol intoxication. *AJNR Am J Neuroradiol* 1995;16:1783-1786
117. Zeiss J, Velasco ME, McCannand KM, Coomlis RJ. Cerebral CT of lethal ethylene glycol intoxication with pathologic correlation. *AJNR Am J Neuroradiol* 1989;10:440-442
118. Korogi Y, Takahashi M, Shinzato J, Okalima T. MR findings in seven patients with organic mercury poisoning (Minamata disease). *AJNR Am J Neuroradiol* 1994;15:1575-1578
119. Truwit CL, Denaro CP, Lake JR, DeMarco T. MR imaging of reversible cyclosporin A-induced neurotoxicity. *AJNR Am J Neuroradiol* 1991;12:651-659
120. George AE, de Leon MJ, Gentes CI, et al. Leukoencephalopathy in normal and pathologic aging: 1. CT of brain lucencies. *AJNR Am J Neuroradiol* 1986;7:561-566
121. de Leon MJ, George AE, Reisberg B, et al. Alzheimer's disease: longitudinal CT studies of ventricular change. *AJR Am J Roentgenol* 1989;152:1257-1262
122. Holodny AI, Waxman R, George AE, Rusinek H, Kalnin AJ, de Leon M. MR differential diagnosis of normal-pressure hydrocephalus and Alzheimer disease: significance of perihypocampal features. *AJNR Am J Neuroradiol* 1998;19:813-819
123. Bradley WG, Jr, Whittemore AR, Watanabe AS, Davis SJ, Teresi LM, Homyak M. Association of deep white matter infarction with chronic communicating hydrocephalus: implications regarding the possible origin of normal-pressure 40 hydrocephalus [see comments]. *AJNR Am J Neuroradiol* 1991;12:31-39
124. Roman GC. White matter lesions and normal-pressure hydrocephalus: Binswanger disease or Hakim syndrome? [comment]. *AJNR Am J Neuroradiol* 1991;12:40-41
125. George AE. Chronic communicating hydrocephalus and periventricular white matter disease: a debate with regard to cause and effect [comment]. *AJNR Am J Neuroradiol* 1991;12:42-44
126. Donnal JF, Heinz ER, Burger PC. MR of reversible thalamic lesions in Wernicke syndrome. *AJNR Am J Neuroradiol* 1990;11:893-894; discussion 895-896
127. Gallucci M, Bozzao A, Splendiani A, Masciocchi C, Passariello R. Wernicke encephalopathy: MR findings in five patients. *AJNR Am J Neuroradiol* 1990;11:887-892
128. Kato Y, Matsumura K, Kinoshita Y, Narita Y, Kuzuhara S, Nakagawa T. Detection of pyramidal tract lesions in amyotrophic lateral sclerosis with magnetization-transfer measurements. *AJNR Am J Neuroradiol* 1997;18:1541-1547
129. Brunberg JA, Kanal E, Hirsch W, Van Thiel DH. Chronic acquired hepatic failure: MR imaging of the brain at 1.5 T. *AJNR Am J Neuroradiol* 1991;12:909-914
130. Savoiardo M, Halliday WC, Nardocci N, et al. Hallervorden-Spatz disease: MR and pathologic findings. *AJNR Am J Neuroradiol* 1993;14:155-162
131. Young IR, Bydder GM, Hall AS, et al. Extracerebral collections: recognition by NMR imaging. *AJNR Am J Neuroradiol* 1983;4:833-834
132. Moon KL, Jr, Brant-Zawadzki M, Pitts LH, Mills CM. Nuclear magnetic resonance imaging of CT-isodense subdural hematomas. *AJNR Am J Neuroradiol* 1984;3:319-322

133. Fobben E, Grossman R, Atlas S, et al. **MR Characteristics of Subdural Hematomas and Hygromas at 1.5T.** *AJNR Am J Neuroradiol* 1989;10:687-693
134. Mc Cluney KW, Yeakley JW, Fenstermacher MJ, Baird SH, Bonmati CM. **Subdural hygroma versus atrophy on MR brain scans: "the cortical vein sign".** *AJNR Am J Neuroradiol* 1992;13:1335-1339
135. Orrison WW, Gentry LR, Stimac GK, Tarrel RM, Espinosa MC, Cobb LC. **Blinded comparison of cranial CT and MR in closed head injury evaluation.** *AJNR Am J Neuroradiol* 1994;15:351-356
136. Gentry LR, Thompson B, Godersky JC. **Trauma to the corpus callosum: MR features.** *AJNR Am J Neuroradiol* 1988;9:1129-1138
137. Blatter DD, Bigler ED, Gale SD, et al. **MR-based brain and cerebrospinal fluid measurement after traumatic brain injury: correlation with neuropsychological outcome.** *AJNR Am J Neuroradiol* 1997;18:1-10
138. Bigler ED, Blatter DD, Anderson CV, et al. **Hippocampal volume in normal aging and traumatic brain injury [see comments].** *AJNR Am J Neuroradiol* 1997;18:11-23
139. Roppolo HM, Latchaw RE, Meyer JD, Curtin HD. **Normal pituitary gland: 1. Macroscopic anatomy-CT correlation.** *AJNR Am J Neuroradiol* 1983;4:927-935
140. Roppolo HM, Latchaw RE. **Normal pituitary gland: 2. Microscopic anatomy-CT correlation.** *AJNR Am J Neuroradiol* 1983;4:937-944
141. Peyster RG, Hoover ED, Viscarello RR, Moshang T, Haskin ME. **CT appearance of the adolescent and preadolescent pituitary gland.** *AJNR Am J Neuroradiol* 1983;4:411-414
142. Pojunaas KW, Daniels DL, Williams AL, Haughton VM. **MR imaging of prolactin-secreting microadenomas.** *AJNR Am J Neuroradiol* 1986;7:209-213
143. Davis PC, Hoffman JC, Jr., Malko JA, et al. **Gadolinium-DTPA and MR imaging of pituitary adenoma: a preliminary report.** *AJNR Am J Neuroradiol* 1987;8:817-823
144. Peck WW, Dillon WP, Norman D, Newton TH, Wilson CB. **High-resolution MR imaging of pituitary microadenomas at 1.5 T. experience with Cushing disease.** *AJR Am J Roentgenol* 1989;152:145-151
145. Newton DR, Dillon WP, Norman D, Newton TH, Wilson CB. **Gd-DTPA-enhanced MR imaging of pituitary adenomas.** *AJNR Am J Neuroradiol* 1989;10:949-954
146. Mark LP, Haughton VM, Hendrix LE, et al. **High-intensity signals within the posterior pituitary fossa: a study with fat-suppression MR techniques.** *AJNR Am J Neuroradiol* 1991;12:529-532
147. Fujisawa I, Asato R, Nishimura K, et al. **Anterior and posterior lobes of the pituitary gland: assessment by 1.5 T MR imaging.** *J Comput Assist Tomogr* 1987;11:214-220
148. Fujisawa I, Nishimura K, Asato R, et al. **Posterior lobe of the pituitary in diabetes insipidus: MR findings.** *J Comput Assist Tomogr* 1987;11:221-225
149. Fujisawa I, Kikuchi K, Nishimura K, et al. **Transection of the pituitary stalk: development of an ectopic posterior lobe assessed with MR imaging.** *Radiology* 1987;165:487-489
150. Kucharczyk W, Lenkinski RE, Kucharczyk J, Henkelman RM. **The effect of phospholipid vesicles on the NMR relaxation of water: an explanation for the MR appearance of the neurohypophysis? [see comments].** *AJNR Am J Neuroradiol* 1990;11:693-700
151. Tien R, Kucharczyk J, Kucharczyk W. **MR imaging of the brain in patients with diabetes insipidus [see comments].** *AJNR Am J Neuroradiol* 1991;12:533-542
152. Moses AM, Clayton B, Hochhauser L. **Use of TI-weighted MR imaging to differentiate between primary polydipsia and central diabetes insipidus [comment] [see comments].** *AJNR Am J Neuroradiol* 1992;13:1273-1277
153. Lundin P, Bergstrom K, Nyman R, Lundberg PO, Muhr C. **Macroprolactinomas: serial MR imaging in long-term bromocriptine therapy [see comments].** *AJNR Am J Neuroradiol* 1992;13:1279-1291
154. Sato N, Sze G, Endo K. **Hypophysitis: endocrinologic and dynamic MR findings.** *AJNR Am J Neuroradiol* 1998;19:439-444
155. Sartoretti-Schefer S, Wichmann W, Aguzzi A, Valavanis A. **MR differentiation of adamantinoid and squamous-papillary craniopharyngiomas.** *AJNR Am J Neuroradiol* 1997;18:77-87
156. Surnida M, Uozumi T, Mukada K, Arita K, Kurisu K, Eguchi K. **Rathke cleft cysts: correlation of enhanced MR and surgical findings.** *AJNR Am J Neuroradiol* 1994;15:525-532
157. Sze G, Milano E, Johnson C, Heier L. **Detection of brain metastases: comparison of contrast-enhanced MR with unenhanced MR and enhanced CT.** *AJNR Am J Neuroradiol* 1990;11:785-791
158. Yuh WT, Engelken JD, Muhonen MG, Mayr NA, Fisher DJ, Ehrhardt JC. **Experience with high-dose gadolinium MR imaging in the evaluation of brain metastases.** *AJNR Am J Neuroradiol* 1992;13:335-345
159. Yuh WT, Tali ET, Nguyen HD, Simonson TM, Mayr NA, Fisher DJ. **The effect of contrast dose, imaging time, and lesion size in the MR detection of intracerebral metastasis [published erratum appears in AJNR Am J Neuroradiol 1995 Jun-Jul;16:1384].** *AJNR Am J Neuroradiol* 1995;16:373-380
160. Sze G, Johnson C, Kawamura Y, et al. **Comparison of single- and triple-dose contrast material in the MR screening of brain metastases [see comments].** *AJNR Am J Neuroradiol* 1998;19:821-828
161. Ginsberg LE, Lang FF. **Neuroradiologic screening for brain metastases—can quadruple dose gadolinium be far behind? [comment].** *AJNR Am J Neuroradiol* 1998;19:829-830
162. Elster AD, Chen MY. **Can nonenhancing white matter lesions in cancer patients be disregarded?** *AJNR Am J Neuroradiol* 1992;13:1309-1315; discussion 1316-1318
163. Lee YY, Van Tassel P. **Intracranial oligodendrogliomas: imaging findings in 35 untreated cases.** *AJR Am J Roentgenol* 1989;152:361-369
164. Lipper MH, Eberhard DA, Phillips CD, Vezina LG, Cail WS. **Pleomorphic xanthoastrocytoma, a distinctive astroglial tumor: neuroradiologic and pathologic features.** *AJNR Am J Neuroradiol* 1993;14:1397-1404
165. Poptani H, Gupta RK, Roy R, Pandey R, Jain VK, Chhabra DK. **Characterization of intracranial mass lesions with in vivo proton MR spectroscopy.** *AJNR Am J Neuroradiol* 1995;16:1593-1603
166. Valk PE, Dillon WP. **Radiation injury of the brain.** *AJNR Am J Neuroradiol* 1991;12:45-62
167. Dillon W, Brant-Zawadzki M, Sherry RG. **Transient computed tomographic abnormalities after focal seizures.** *AJNR Am J Neuroradiol* 1984;5:107-109
168. Schwartz RB, Carvalho PA, Alexander Ed, Loeffler JS, Folkert R, Holman BL. **Radiation necrosis vs high-grade recurrent glioma: differentiation by using dual-isotope SPECT with 201 TI and 99mTc-HMPAO.** *AJNR Am J Neuroradiol* 1991;12:1187-1192
169. Ricci PE, Karis JP, Heiserman JE, Fram EK, Bice AN, Drayer BP. **Differentiating recurrent tumor from radiation necrosis: time for re-evaluation of positron emission tomography? [see comments].** *AJNR Am J Neuroradiol* 1998;19:407-413
170. Patronas NJ, Di Chiro G, Brooks RA, et al. **Work in progress: [18F] fluorodeoxyglucose and positron emission tomography in the evaluation of radiation necrosis of the brain.** *Radiology* 1982;144:885-889
171. Andreasen NC. **Neuroradiology and neuropsychiatry: a new alliance [comment].** *AJNR Am J Neuroradiol* 1992;13:841-843
172. Degreaf G, Lantos G, Bogerts B, Ashtari M, Lieberman J. **Abnormalities of the septum pellucidum on MR scans in first-episode schizophrenic patients [see comments].** *AJNR Am J Neuroradiol* 1992;13:835-840
173. Seidenwurm D, Pounds TR, Globus A, Valk PE. **Abnormal temporal lobe metabolism in violent subjects: correlation of imaging and neuropsychiatric findings.** *AJNR Am J Neuroradiol* 1997;18:625-631
174. Gonzalez RG, Guimaraes AR, Sachs GS, Rosenbaum JF, Garwood M, Renshaw PF. **Measurement of human brain lithium in vivo by MR spectroscopy [see comments] published erratum appears in AJNR Am J Neuroradiol 1993 Nov-Dec;14:14501.** *AJNR Am J Neuroradiol* 1993;14:1027-1037
175. Sunshine JL, Lewin JS, Wu DH, et al. **Functional MR to localize sustained visual attention activation in patients with attention deficit hyperactivity disorder— a pilot study.** *AJNR Am J Neuroradiol* 1997;18:633-637
176. Silverstein AM, Alexander JA. **Acute postictal cerebral imaging.** *AJNR Am J Neuroradiol* 1998;19:1485-1488
177. Meiners LC, van Gils A, Jansen GH, et al. **Temporal lobe epilepsy: the various MR appearances of histologically proven**

- mesial temporal sclerosis.** *AJNR Am J Neuroradiol* 1994;15:1547-1555
178. Oppenheim C, Dormont D, Biondi A, et al. **Loss of digitations of the hippocampal head on high-resolution fast spin-echo MR: a sign of mesial temporal sclerosis.** *AJNR Am J Neuroradiol* 1998;19:457-463
179. Cheon JE, Chang KH, Kim HD, et al. **MR of hippocampal sclerosis: comparison of qualitative and quantitative assessments.** *AJNR Am J Neuroradiol* 1998;19:465-468
180. Lee DH, Gao FQ, Rogers JM, et al. **MR in temporal lobe epilepsy: analysis with pathologic confirmation.** *AJNR Am J Neuroradiol* 1998;19:19-27