Generic Contrast Agents Our portfolio is growing to serve you better. Now you have a *choice*.





This information is current as of May 7, 2025.

CNS *Clostridium perfringens* **Infection: A Rare Complication of Preoperative Embolization of Meningioma**

P.-S. Yen, C.-c. Lin, C.-C. Lee, T. Harnod, T.-W. Loh and Y.-H. Hsu

AJNR Am J Neuroradiol 2006, 27 (6) 1355-1356 http://www.ajnr.org/content/27/6/1355

CASE REPORT

P.-S. Yen C.-c. Lin C.-C. Lee T. Harnod T.-W. Loh Y.-H. Hsu

CNS *Clostridium perfringens* Infection: A Rare Complication of Preoperative Embolization of Meningioma

SUMMARY: Gas gangrene is a severe form of gangrene (tissue death) that usually is caused by *Clostridium perfringens.* It generally occurs at the site of trauma or a recent surgical wound. We report the case of a 45-year-old woman with sphenoid-ridged meningioma who received preoperative transarterial embolization with polyvinyl alcohol. The patient later developed an intratumoral *C perfringens* infection and died despite intensive medical care. The case represents an extremely rare complication following transarterial embolization of meningioma.

Preoperative embolization of meningiomas is considered valuable to allow simpler resection by reducing intraoperative blood loss and decreasing surgery time. Though the embolization procedures are quite safe with careful techniques in our general practice,¹ reports of some rare complications such as tumoral and subarachnoid hemorrhage, scalp necrosis, retinal embolus, facial nerve palsies, and occlusion of the vasa nervorum of cranial nerves—have been published. We now report what is, to the best of our knowledge, the first case of gas gangrene due to *Clostridium perfringens* infection after preoperative embolization of a meningioma. We also propose that prophylactic antibiotics should be used in meningioma embolization to avoid this complication.

Case Report

A 45-year-old woman with complaints of intermittent headache, nausea, dizziness, and progressive loss of visual acuity during a 5-year period was admitted to our emergency department because of aggravation of these symptoms. A cerebral MR imaging study revealed a large meningioma, about 5.5 cm \times 5 cm \times 5 cm in size, at the right sphenoid ridge. The tumor was accompanied with perifocal edema and subfalcine herniation. Because of the neovascularity of the tumor and its large size, preoperative embolization was planned. Cerebral angiography showed a large hypervascular tumor mainly fed by branches of the right middle meningeal artery. Thereafter, selective embolization of the right middle meningeal artery using coaxial microcatheter systems and standard technique was performed. The embolization was performed by using one bottle of $50-100-\mu$ polyvinyl alcohol (PVA) particles (Ultra-Drivalon; CathNet-Science, Paris, France) followed by half a bottle of $150-250-\mu$ PVA particles. A postembolization arteriogram showed a significant decrease in tumor vascularity. The patient then returned to the ward in stable condition. Unfortunately, the patient developed a high fever (\leq 39.3°C) about 1.5 days after the embolization procedure. The patient complained of a persistent headache despite analgesics and mannitol infusion. About 48 hours after the embolization procedure, the patient suddenly lost consciousness with ipsilateral pupil dilation. An emergency brain CT study showed tumor necrosis with gas formation

inside and around the tumor (Fig 1). There was associated severe brain edema with subfalcine and right uncal herniation. Surgical decompression with subtotal tumor removal was immediately performed. Histopathology of the tumor showed marked necrosis with residual meningioma and gas bubble formation. Gram-positive organisms were found (Fig 2). Later, both tissue and blood culture yielded *C perfringens*. Because of irreversible brain stem dysfunction, this patient died 16 days after the onset of uncal herniation, despite intensive management.

Review of the patient's history revealed no obvious symptoms or signs of infection in the upper respiratory tract, gastrointestinal tract, or urinary tract before admission.

Discussion

The goal of preoperative meningioma embolization is safer and more successful surgical procedures by reducing operating time and blood loss. Toward this end, the surgeon should plan the procedure to minimize risks. This procedure is safe and effective,¹ with complication rates ranging from $1.6\%^2$ to 3%.³ Anecdotal reports of complications include tumoral and subarachnoid hemorrhage, scalp necrosis, retinal emboli, facial nerve palsies, and occlusion of the vasa nervorum of cranial nerves. To the best of our knowledge, postembolization C perfringens infection has not been reported. Indeed, C perfringens is a rare cause of CNS infection, usually occurring as a complication of penetrating trauma. For example, neurosurgical procedures were reported in 9 cases: 2 were due to intraoperative contamination^{4,5}; 2 had entry portals due to surgical drain and intra-abdominal infection^{6,7}; and 5 had no precise source of infection.⁸⁻¹² Six of the 9 died of meningitis, and one died of brain abscess.

Spontaneous gas gangrene without external wound or injury occurs frequently in patients who have serious underlying conditions, such as vascular disease (atherosclerosis), diabetes, or colorectal or hematologic malignancy. In our patient, infection might have been due to contamination during the procedure or to hematogenous spread from respiratory gastrointestinal or genitourinary tracts. It was not due to PVA particles, because no bacterial growth was found when PVA was put into anaerobic culture.

The course of infection is variable, depending on host immunity and various toxins produced by *C perfringens*, such as epsilon toxin, which is thought to damage the blood-brain barrier. *C perfringens* is a Gram-positive, gas-forming, sporing

Received May 1, 2005; accepted after revision August 14.

From the Departments of Radiology (P.-S.Y., C.-c.L., C.-C.L.), Neurosurgery (T.H., T.-W.L.), and Pathology (Y.-H.H.), Buddhist Tzu Chi General Hospital, Hualien, Taiwan.

Address reprints requests and correspondence to Dr. Chao-chun Lin, Department of Radiology, Buddhist Tzu Chi General Hospital, 707, Section 3 Chung Yang Rd, Hualien, Taiwan.



Fig 1. Unenhanced axial brain CT image taken about 2 days after the preoperative meningioma embolization showed tumor necrosis with gas formation inside and around the tumor.



Fig 2. Gram stain (\times 400) of the specimen from the tumor shows numerous Gram-positive bacilli consistent with *C perfringens*.

commensal bacterium of the colon and vagina. It grows only in mechanically or chemically damaged tissues. In one study,¹³ extended microembolization with 50–150- μ PVA particles caused significant tumor necrosis seen on MR imaging and histopathologic examination after surgery, as compared with larger particle embolization. The authors concluded that extended microembolization with small PVA particles improved surgical treatment of meningiomas. We had performed about 100 cases of meningioma embolization with 150–250 μ PVA particles without major complications. This was the first in which we extended microembolization with small PVA particles. The devascularization of the tumor with subsequent tumoral ischemia and necrosis may have provided an anaerobic environment for the growth of *C perfringens*. Whether embolization with small particles increased the risk of *C perfringens* infection is unclear, because no other cases were reported.

We conclude that this rare complication of preoperative meningioma embolization should be kept in mind in the differential diagnosis of postembolization fever. *C perfringens* infection remains fatal despite frequent, prompt, and aggressive treatment.

References

- Gruber A, Killer M, Mazal P, et al. Pre-operative embolization of intracranial meningiomas: a 17-year single-center experience. *Minim Invasive Neurosurg* 2000;43:8–29
- Lasjaunias P, Berenstein A. Surgical neuroangiography. Vol 2: Endovascular treatment of craniofacial lesions. Berlin: Springer-Verlag;1987
- 3. Bendszus M, Rao G, Burger R, et al. **Is there a benefit of preoperative meningioma embolization**? *Neurosurgery* 2000;47:1306–311; discussion 1311–12
- Cohen JE, Mierez R, Tsai EC. Postcraniotomy gas-containing brain abscess: a neurosurgical emergency: case report. Surg Neurol 1999;51:568–70
- Willis AT, Jacobs SI. Case of meningitis due to Clostridium welchii. J Pathol Bacteriol 1964;88:312–14
- Debast SB, van Rijswijk E, Jira PE, et al. Fatal Clostridium perfringens meningitis associated with insertion of a ventriculoperitoneal shunt. Eur J Clin Microbiol 1993;12:720–21
- 7. Ho K. Clostridium perfringens meningitis. Int Surg 1982;67:271-73
- Alexander JG. A case of fatal Clostridium welchii toxaemia due to ward infection. J Clin Pathol 1969;22:508
- Bornstein DJ, Weinberg AN, Swartz MN, et al. Anaerobic infections: review of current experience. *Medicine* 1964;43:207–32
- Long JG, Preblud SR, Keyserling HL. Clostridium perfringens meningitis in an infant: case report and literature review. Pediatr Infect Dis J 1987;6:752–54
- Kristopaitis T, Jensen R, Gujrati M. Clostridium perfringens: a rare cause of postoperative spinal surgery meningitis. Surg Neurol 1999;51:448–50; discussion 450–51
- Penrose-Stevens A, Ibrahim A, Redfern RM. Localized pneumocephalus caused by Clostridium perfringens meningitis. Br J Neurosurg 1999;13:85–86
- 13. Wakhloo AK, Juengling FD, Van Velthoven V, et al. Extended preoperative polyvinyl alcohol microembolization of intracranial meningiomas: assessment of two embolization techniques. *AJNR Am J Neuroradiol* 1993;14:571–82