

Are your **MRI contrast agents** cost-effective?

Learn more about generic **Gadolinium-Based Contrast Agents**.



**FRESENIUS  
KABI**

caring for life

**AJNR**

## **Closing the Uncertainty Gap in the Diagnosis of Parotid Tumors**

J.A. Castelijns and C. René Leemans

*AJNR Am J Neuroradiol* 2009, 30 (3) 442-443

doi: <https://doi.org/10.3174/ajnr.A1423>

<http://www.ajnr.org/content/30/3/442>

This information is current as of April 19, 2024.

ical journals in the world are published completely or partly in English.

We at *AJNR* do our best to publish author's names correctly, but because of an increasing number of articles originating from outside of the United States, we urge authors to take responsibility and to let us know exactly how they want their name to appear. In this editorial, I have attempted to give our readers an idea about the complexity involved.

## References

1. Sun XL, Zhou J. English versions of Chinese author's names in biomedical journals: observations and recommendations. *Science Editor* 2002;25:3-4
2. Han S. Formats of Korean author's names. *Science Editor* 2005;28:189-90
3. Japanese name. From Wikipedia, the free encyclopedia. Available at: [http://en.wikipedia.org/wiki/Japanese\\_name](http://en.wikipedia.org/wiki/Japanese_name). Accessed September 12, 2008.
4. Black B. Indexing the names of authors from Spanish- and Portuguese-speaking countries. *Science Editor* 2003;26:118-21
5. Ruiz-Perez R, Delgado Lopez-Cozar E, Jimenez-Contreras E. Spanish personal name variations in national and international biomedical databases: implications for information retrieval and bibliometric studies. *J Med Libr Assoc* 2002;90:411-30
6. Kidambi M. Indian names: a guide for science editors. *Science Editor* 2008;31:120-21
7. Notzon B, Nesom G. The Arabic naming system. *Science Editor* 2005;28:20-21

M. Castillo  
Editor-in-Chief

DOI 10.3174/ajnr.A1367

## EDITORIAL

### Closing the Uncertainty Gap in the Diagnosis of Parotid Tumors

In this issue of the *AJNR American Journal of Neuroradiology*, Habermann et al<sup>1</sup> assess the value of diffusion-weighted echo-planar MR imaging in differentiating various types of primary parotid gland tumors in a large group of patients with a lesion of the parotid gland. They claim that, to a certain extent, diffusion-weighted echo-planar MR imaging has the potential for differentiation between various primary tumors of the parotid gland.

Parotid gland tumors are relatively rare and include a large variety of histologic types. Therefore, imaging of those tumors is a major challenge for radiologists.

The management of tumors of the parotid gland requires a detailed understanding of the anatomy and the pathologic processes affecting these glands and surrounding structures. Patients with these diseases are usually referred to a specialist from a head and neck center, where multidisciplinary management is possible. Management by the clinician, usually a surgical specialist for head and neck diseases, is based on history, physical examination, and, always, imaging modalities (most notably, MR imaging) and other investigations if indicated, such as sonography-guided fine-needle aspiration cytology (USgFNAC). In general, FNAC by experienced clinicians or radiologists (under sonography guidance) is able to give certain diagnoses with high, albeit not 100%, accuracy.

The most important division between diagnosis groups lies between benign and malignant (roughly 20%) lesions. Treat-

ment is usually by surgical excision, but the extent is dictated by the pathologic process. The preoperative information on whether a parotid gland tumor is benign or malignant may be helpful to prevent treatment delay in the case of a malignant tumor and in patient counseling as to likely treatment and possible sequels. Also, there are well-established indications for performing an elective neck dissection in the case of high-grade malignant tumors.

All aforementioned methods can help to establish a working diagnosis, but ultimately often only histopathologic assessment after removal of the lesion yields a definite diagnosis on which additional management, if indicated, may be based. Also, within the benign category a distinction between the 2 most frequent tumors (ie, pleomorphic adenoma and Warthin tumor) is valuable because a Warthin tumor, especially in elderly patients or those reluctant to submit to surgery, can be observed with repeated clinical or radiologic measurements.

Therefore, every effort to close the uncertainty gap is highly valuable in patient management, though everyday practice often makes extensive work-up impractical and unnecessary. This is because in many patients, removal is indicated or patients elect to have surgery. Also, if waiting lists for surgery do not dictate otherwise, this is often the most practical and rapid solution. On the other hand, if waiting lists are long, less uncertainty about the diagnosis is extremely helpful to expedite surgery in those in whom high suspicion of a malignant tumor is present.

In contrast to FNAC, a noninvasive examination such as MR imaging might not only be helpful in providing a diagnosis but also with regard to staging by providing information on the exact localization and extent of the lesion. Differentiation between benign and malignant disease is often possible on the basis of morphologic appearance alone.<sup>2</sup> However, even the combined accuracy of the modalities USgFNAC and conventional MR imaging may still be imperfect.

New MR imaging modalities such as proton MR spectroscopy,<sup>3</sup> dynamic contrast-enhanced MR imaging,<sup>4</sup> and diffusion-weighted MR imaging<sup>5</sup> have all shown promising results in the differentiation between benign and malignant salivary disease. Regarding diffusion-MR imaging, one should cope with some major challenges. First, diffusion-weighted MR imaging should be, in principle, independent from the MR imaging machine that is used, but it is not entirely certain if this is really true. Hein et al<sup>6</sup> applied some type of standardization of apparent diffusion coefficient (ADC) values by comparing it with ADC values of healthy tissues. This may be difficult to realize in the head and neck region because it may be difficult to find reliable tissue for reference. Second, there is some specificity of ADC values for certain tumor types, but these reports are not unequivocal. Therefore, more studies with large patient groups need to be performed. It is reported that the choice of b-value might be helpful in the differentiation of benign and malignant tumors.<sup>7</sup> The lower the applied b-values, the higher the influence of perfusion, whereas applying only the ADC by high b-values reflects mainly diffusion.

The value of Habermann's study is that it is the first to report on patients with parotid tumors examined with diffusion-weighted MR imaging. The main limitation of this study

was that the number of cases was not satisfactory for every different tumor. Furthermore, the number of malignant tumors as a whole was very limited.

The differentiation of benign and malignant masses may be improved by combining morphologic features and newer MR imaging techniques, such as diffusion-weighted MR imaging, MR spectroscopy, and dynamic contrast-enhanced MR imaging. Additional multicenter research with larger numbers of cases with an adequate distribution among tumor types with use of these novel MR methods is warranted.

## References

1. Habermann CR, Aradt C, Graessner J. **Diffusion-weighted echo-planar MR imaging of primary parotid gland tumors: Is a prediction of different histologic subtypes possible?** *AJNR Am J Neuroradiol* 2009;30:581–96
2. Prades JM, Oletski A, Faye MB, et al. **Parotid gland masses: diagnostic value of MR imaging with histopathologic correlations.** *Morphologie* 2007;91:44–51

3. King AD, Yeung DK, Ahuja AT, et al. **Salivary gland tumors at in vivo proton MR spectroscopy.** *Radiology* 2005;237:563–69
4. Yabuuchi H, Fukuya T, Tajimia T, et al. **Salivary gland tumors: diagnostic value of gadolinium-enhanced dynamic MR imaging with histopathologic correlation.** *Radiology* 2003;226:345–54
5. Wang J, Takashima S, Takayama F, et al. **Head and neck lesions: characterization with diffusion-weighted echo-planar MR imaging.** *Radiology* 2001;220:621–30
6. Hein PA, Eskey CJ, Dunn JF, et al. **Diffusion-weighted imaging in the follow-up of treated high-grade gliomas: tumor recurrence versus radiation injury.** *AJNR Am J Neuroradiol* 2004;25:201–09
7. Thoeny HC, De Keyzer F, Boesch C, et al. **Diffusion-weighted imaging of the parotid gland: influence of the choice of b-values on the apparent diffusion coefficient value.** *J Magn Reson Imaging* 2004;20:786–90

J.A. Castelijns

Department of Radiology

C. René Leemans

Department of Otolaryngology/Head and Neck Surgery

VU University Medical Center

Amsterdam, the Netherlands

DOI 10.3174/ajnr.A1423