

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



FRESENIUS
KABI

caring for life

AJNR

Reply:

P. Mukherjee, M. Wahl and A.J. Barkovich

AJNR Am J Neuroradiol 2009, 30 (6) e82

doi: <https://doi.org/10.3174/ajnr.A1606>

<http://www.ajnr.org/content/30/6/e82>

This information is current as
of April 19, 2024.

Reply:

Thank you for your insightful comments on our recent diffusion tractography study of partial agenesis of the corpus callosum¹ and for bringing to our attention your prior work on MR imaging of partial callosal agenesis.² Our own structural MR imaging analysis showing significant variation in the size and location of callosal fragments in individuals with partial agenesis is in good agreement with your previous results. Taken together, they provide strong evidence that callosal fragments in individuals with partial callosal agenesis do not represent the incomplete growth of a normal callosum.

Our study supported and extended this hypothesis by using diffusion tensor imaging (DTI) and high angular resolution diffusion imaging to perform 3D tractography, a technique unavailable at the time of your study. We determined the connectivity of axons passing through each callosal fragment and compared the pattern of connectivity with what would be expected based on the location of the fragment alone. We thus demonstrated not only that the positions of callosal fragments are variable (as can be shown with conventional MR imaging) but that the connectivity through a callosal fragment is also variable and cannot be determined on the basis of its anatomic location. These results confirm, in human subjects, what was found in the mouse studies you cite using horseradish peroxidase track tracing.^{3,4}

In reading your article, we found the breakdown of your results based on associated brain abnormalities, such as Chiari II or Dandy-Walker malformations, particularly interesting. Although 10 of all 25 subjects in your study demonstrated a callosal fragment corresponding to the position of a normal callosum, only 2 of 10 subjects without associated abnormalities had a fragment at a normal position. Our cohort was restricted to individuals with either no or minor associated

abnormalities, and all 6 of our subjects had connectivity patterns different from what would be expected based on incomplete callosal growth. These results together suggest that subjects with partial agenesis without associated abnormalities are more likely to develop callosal fragments unexplainable by incomplete callosal growth. Because individuals with associated brain abnormalities likely have different underlying causes of callosal dysgenesis, this result is perhaps not surprising. However, a more comprehensive classification of callosal morphology and connectivity based on etiology could yield significant insight into the developmental mechanisms of callosal dysgenesis. This awaits further investigation in larger cohorts.

References

1. Wahl M, Strominger Z, Jeremy RJ, et al. **Variability of homotopic and heterotopic callosal connectivity in partial agenesis of the corpus callosum: a 3T diffusion tensor imaging and Q-ball tractography study.** *AJNR Am J Neuroradiol* 2009;30:282–89. Epub 2008 Nov 11
2. Rubinstein D, Youngman V, Hise JH, et al. **Partial development of the corpus callosum.** *AJNR Am J Neuroradiol* 1994;15:869–75
3. Ozaki HS, Shimada M. **The fibers which course within the Probst's longitudinal bundle seen in the brain of a congenitally acallosal mouse: a study with the horseradish peroxidase technique.** *Brain Res* 1988;441:5–14
4. Olavarria J, Serra Oller MM, Yee KT, et al. **Topography of interhemispheric connections in neocortex of mice with congenital deficiencies of the callosal commissure.** *J Comp Neurol* 1988;270:575–90

P. Mukherjee

M. Wahl

A.J. Barkovich

Department of Radiology

University of California

San Francisco, Calif

DOI 10.3174/ajnr.A1606