Generic Contrast Agents

FRESENIUS KABI

Our portfolio is growing to serve you better. Now you have a *choice*.



Reply:

E.M. Genden

AJNR Am J Neuroradiol 2010, 31 (2) E31 doi: https://doi.org/10.3174/ajnr.A1998 http://www.ajnr.org/content/31/2/E31

This information is current as of May 5, 2025.

Reply:

The American Thyroid Association guidelines for the management of patients with thyroid nodules¹ outlines the role of sonography and fine-needle aspiration for the evaluation of thyroid nodules. Our work and the work of others²⁻⁴ suggest that fluorodeoxyglucosepositron-emission tomography (FDG-PET) positivity is associated with a significant risk of malignancy, not unlike the cytologic reading of "indeterminate cytology," which, according to the task force (R9, recommendation B), states that a "lobectomy or total thyroidectomy should be considered."1 We support the work of the American Thyroid Association and believe that FDG-PET positivity simply represents an adjunct to sonography and cytology for the risk assessment of a patient with a thyroid nodule. We do not advocate thyroidectomy for patients with benign cytology and retract any implication of such; however, we do inform patients of the data on FDG-PET positivityafter all, the false-negative rate for fine-needle aspiration of thyroid nodules is higher than the false-positive rate. The American Thyroid Association recommends that a lobectomy or total thyroidectomy be considered for indeterminate cytology, which is associated with a 5%-10% risk of malignancy. Do we feel comfortable withholding surgical intervention when FDG-PET positivity is associated with a 25%–50% risk of malignancy?

References

- Cooper DS, Doherty GM, Haugen BR, et al. Management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* 2006;16:109–42
- Choi JY, Lee KS, Kim HJ, et al. Focal thyroid lesions incidentally identified by integrated 18F-FDG PET/CT: clinical significance and improved characterization. J Nucl Med 2006;47:609–15
- Kang KW, Kim SK, Kang HS, et al. Prevalence and risk of cancer of focal thyroid incidentaloma identified by 18F-fluorodeoxyglucose positron emission tomography for metastasis evaluation and cancer screening in healthy subjects. J Clin Endocrinol Metab 2003;88:4100-04
- 4. Chen YK, Ding HJ, Chen KT, et al. Prevalence and risk of cancer of focal thyroid incidentaloma identified by 18F-fluorodeoxyglucose positron emission tomography for cancer screening in healthy subjects. Anticancer Res 2005;25:1421-6

E.M. Genden Chairman—Department of Otolaryngology Mount Sinai Medical Center New York, New York

DOI 10.3174/ajnr.A1998