

Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a choice.



[VIEW CATALOG](#)

ERRATUM

AJNR

AJNR Am J Neuroradiol 2015, 36 (6) E52
doi: <https://doi.org/10.3174/ajnr.A4381>
<http://www.ajnr.org/content/36/6/E52>

This information is current as
of May 6, 2025.

Please note that there is an error in Table 2 of the February 2015 article “Assessment of Intracranial Collaterals on CT Angiography in Anterior Circulation Acute Ischemic Stroke” by Yeo et al (*AJNR Am J Neuroradiol* 2015;36:289–94), originally published on-line on October 16, 2014, doi:10.3174/ajnr.A4117). The population of patients with good collaterals by Miteff scoring with good outcomes has been switched with the population with poor outcomes. The corrected Table 2 is reproduced below with the corrected segment shown in bold. The authors apologize for the error.

<http://dx.doi.org/10.3174/ajnr.A4381>

Table 2: Determinants of functional outcome in patients with thrombolyzed acute ischemic stroke

Variable	mRS 0–1 (n = 93)	mRS 2–6 (n = 107)	P Value
Median age (yr) (range)	58 (33–89)	71 (34–92)	.040
Female sex (%)	20 (21.5)	43 (40.2)	.005
Hypertension (%)	51 (54.8)	80 (74.8)	.034
Diabetes (%)	22 (23.6)	38 (35.8)	.069
Dyslipidemia (%)	45 (48.4)	57 (53.3)	.491
Atrial fibrillation (%)	23 (24.7)	50 (46.9)	.001
Smoker (%)	33 (35.5)	24 (22.4)	.119
Median NIHSS pre-tPA (range)	14 (3–29)	21 (8–33)	.477
Mean SBP pre-tPA (range)	142 (110–209)	154 (110–210)	.777
Median onset-to-treatment time (range)	152 (91–275)	158 (73–265)	.199
ASPECTS <7	36 (38.7)	62 (57.9)	.010
Recanalization of vessel	70 (75.3)	76 (71)	.040
Good collaterals by Tan method	74 (79.6)	49 (45.8)	.001
Good collaterals by Miteff method	66 (61.7)	25 (26.9)	.001
Good collaterals by Maas method	42 (45.2)	23 (21.5)	.005
Median score by ASPECTS methodology (range)	11 (0–18)	8 (1–18)	.021

Note:—SBP indicates systolic blood pressure.