

Are your MRI contrast agents cost-effective?

Learn more about generic Gadolinium-Based Contrast Agents.



FRESENIUS
KABI

caring for life

AJNR

Reply:

J.T. Philip, M.A. Flores, R.D. Beegle, S.C. Dodson, S.A. Messina and J.V. Murray, Jr.

AJNR Am J Neuroradiol 2023, 44 (2) E12

doi: <https://doi.org/10.3174/ajnr.A7770>

<http://www.ajnr.org/content/44/2/E12>

This information is current as
of April 18, 2024.

REPLY:

We appreciate the feedback from Branstetter et al. As with any retrospective study, our study has limitations. One of the main limitations of our study is that the lumbar punctures (LPs) were performed by a variety of radiologists with variable levels of experience, including trainees.¹ While this feature introduces potential confounding variables, we believe it is more indicative of how LPs are performed in most radiology departments; thus, it is a more accurate estimation of post-dural puncture headaches (PDPHs) necessitating an epidural blood patch (EBP). The number of LPs performed in radiology has been increasing.² They are performed by a variety of practitioners with a variety of experience levels, including increasingly by nonphysician providers such as in the study of Rodriguez et al.³ We believe our study is a “real-world” estimation of EBP rates in a heterogeneous patient population with a heterogeneous group of operators. Our study was designed to examine whether there were differences in EBP rates when comparing traumatic with atraumatic needles stratified to the body mass index, not to determine whether there were different EBP rates based on sex or age.

We do not know why our rates of EBP in the Quincke group were higher than those found in the study of Rodriguez et al.³ A different patient cohort certainly could be contributory. At the institution where the data were collected, there was a relatively low threshold to perform an EBP in patients with post-LP headaches, and there was a streamlined process for clinicians to refer patients for EBP. Perhaps the threshold for performing an EBP was different at the two institutions. The rate of post-LP “any headache” was higher than the “severe headache” rate at 2.2% in the study of Rodriguez et al.³ Our rate of EBP in the Quincke group was well below that reported in the study from Hatfield et al.⁴ for fluoroscopically guided LPs with a 22-ga Quincke needle, which was 15.2%.

We acknowledge an error in Table 3 of our article. As stated in our results section, we found no statistically significant difference

in rates of EBP when comparing the 22-ga Quincke with the 20-ga Quincke needle. The confidence interval in Table 3 is correct. The corrected *P* value is .411.

We believe that our study and other studies involving both fluoroscopically guided and non-fluoroscopically guided LPs are compelling in demonstrating an increased rate of PDPHs necessitating an EBP when comparing traumatic-versus-atraumatic spinal needles. However, these studies suffer from the inherent limitations of retrospective studies. We believe a well-designed prospective randomized study would likely end any remaining debate.


REFERENCES


1. Philip JT, Flores MA, Beegle RD, et al. **Rates of epidural blood patch following lumbar puncture comparing atraumatic versus bevel-tip needles stratified for body mass index.** *AJNR Am J Neuroradiol* 2022;43:315–18 [CrossRef Medline](#)
2. Johnson DR, Waid MD, Rula EY, et al. **Comparison of radiologists and other specialists in the performance of lumbar puncture procedures over time.** *AJNR Am J Neuroradiol* 2021;42:1174–81 [CrossRef Medline](#)
3. Rodriguez D, Branstetter BF, Agarwal V, et al. **JOURNAL CLUB: incidence of complications following fluoroscopically guided lumbar punctures and myelograms.** *AJR Am J Roentgenol* 2016;206:20–25 [CrossRef Medline](#)
4. Hatfield MK, Handrich SJ, Willis JA, et al. **Blood patch rates after lumbar puncture with Whitacre versus Quincke 22- and 20-gauge spinal needles.** *AJR Am J Roentgenol* 2008;190:1686–89 [Medline](#)

 J.T. Philip

University of Texas Southwestern Medical Center
Dallas, Texas

 M.A. Flores

 R.D. Beegle

 S.C. Dodson

AdventHealth Medical Group Radiology at Central Florida
Orlando, Florida

 S.A. Messina

Mayo Clinic Rochester
Rochester, Minnesota

 J.V. Murray, Jr.

Mayo Clinic
Jacksonville, Florida

<http://dx.doi.org/10.3174/ajnr.A7770>