

Generic Contrast Agents

Our portfolio is growing to serve you better. Now you have a *choice*.



[VIEW CATALOG](#)

AJNR









This information is current as of May 10, 2025.

Endovascular Management of Acute Stroke in the Elderly: A Systematic Review and Meta-Analysis

C.A. Hilditch, P. Nicholson, M.H. Murad, A. Rabinstein, J. Schaafsma, A. Pikula, T. Krings, V.M. Pereira, R. Agid and W. Brinjikji

AJNR Am J Neuroradiol published online 22 March 2018
<http://www.ajnr.org/content/early/2018/03/22/ajnr.A5598>

Endovascular Management of Acute Stroke in the Elderly: A Systematic Review and Meta-Analysis

 C.A. Hilditch,  P. Nicholson,  M.H. Murad,  A. Rabinstein,  J. Schaafsma,  A. Pikula,  T. Krings,  V.M. Pereira,  R. Agid, and  W. Brinjikji



ABSTRACT

BACKGROUND: Acute ischemic stroke occurs more frequently, presents with more severe symptoms, and has worse outcomes in elderly patients. The safety and efficacy of endovascular therapy for acute stroke in this age group has not been fully established.

PURPOSE: We present the results of a systematic review and meta-analysis examining clinical, procedural, and radiologic outcomes of endovascular therapy for acute stroke in patients older than 80 years of age.

DATA SOURCES: We searched PubMed, MEDLINE, and EMBASE from 1992 to week 35 of 2017 for studies evaluating endovascular therapy for acute stroke in the elderly.

STUDY SELECTION: Two independent reviewers selected studies and abstracted data. The primary end point was good functional outcome at 3 months defined as modified Rankin Scale score of ≤ 2 .

DATA ANALYSIS: Data were analyzed using random-effects meta-analysis.

DATA SYNTHESIS: Seventeen studies reporting on 860 patients were included. The rate of good functional outcome at 3 months was 27% (95% CI, 21%–32%). Mortality at 3 months was 34% (95% CI, 23%–44%). Successful recanalization was achieved in 78% of patients (95% CI, 72%–85%). Procedure-related complications occurred in 11% (95% CI, 4%–17%). The incidence of intracranial hemorrhage was 24% (95% CI, 15%–32%), and for symptomatic intracranial hemorrhage, it was 8% (95% CI, 5%–10%). The mean time to groin was 251 minutes (95% CI, 224–278 minutes). Procedure time was 99 minutes (95% CI, 67–131 minutes).

LIMITATIONS: I^2 values were above 50% for all outcomes, indicating substantial heterogeneity.

CONCLUSIONS: Good functional recovery in octogenarians treated with endovascular therapy for acute stroke can be achieved in a high proportion of patients despite the higher incidence of comorbidity in this cohort. Outcomes are inferior to those reported for younger patients; however, endovascular therapy can allow at least 1 in 4 patients older than 80 years of age to regain independent function at 3 months. More research is required to improve patient selection in the elderly, but age should not be a discriminator when deciding to offer endovascular therapy for patients with acute stroke.

ABBREVIATION: EVT = endovascular therapy

Due to improvement in health care and better standards of living, the proportion of elderly individuals (older than 80 years of age) is expected to at least double in the United States by

the year 2050, and more than double in the European Union by 2080.^{1,2} With aging of the general population, the burden of cardiovascular and cerebrovascular disease will substantially increase. In fact, prior studies have shown that individuals older than 80 years of age have the highest incidence of acute ischemic stroke.³

Endovascular therapy (EVT) has been proven to improve outcomes in patients with stroke from large-vessel occlusion.⁴ However, 3 of the 6 major trials of EVT for acute stroke excluded older patients.^{5–7} Trials that included octogenarians clearly showed better outcomes among patients treated with EVT in this age sub-

Received November 13, 2017; accepted after revision January 24, 2018.

From the Division of Neuroradiology (C.A.H., P.N., T.K., V.M.P., R.A., W.B.), Joint Division of Medical Imaging, and Division of Neurosurgery (T.K., V.M.P.), Department of Surgery, University Health Network, University of Toronto, Toronto, Ontario, Canada; Evidence-Based Practice Center (M.H.M.) and Department of Neurology (A.R.), Mayo Clinic, Rochester, Minnesota; and Division of Neurology (J.S., A.P.), Toronto Western Hospital, Toronto, Ontario, Canada.

Data could be made available by contacting the corresponding author via e-mail.

Please address correspondence to Christopher Alan Hilditch, MD, Toronto Western Hospital, Department of Neuroradiology, 399 Bathurst St, Toronto, ON M5T 2S8 Canada; e-mail: christopheralan.hilditch@uhn.ca; @hilditchca



Indicates article with supplemental on-line table.

<http://dx.doi.org/10.3174/ajnr.A5598>

group compared with intravenous thrombolysis alone, with odds ratios even more favorable than among younger patients.⁸ Yet, octogenarians were relatively underrepresented even in those trials that did not exclude them, suggesting a possible selection bias (ie, only the octogenarians with greater chances of recovery may have been preferentially enrolled). Meanwhile, there have been several prospective and retrospective observational studies reporting the outcomes of endovascular revascularization in acute stroke in octogenarians. However, many of these studies are limited by small sample sizes or restricted to single centers, thus limiting the generalizability of the results.⁹⁻²⁴ To better understand the outcomes after EVT in octogenarians, we performed a systematic review and meta-analysis of published observational studies⁹⁻²⁴ that examined clinical and radiologic outcomes in these patients; we also included data from the Highly Effective Reperfusion evaluated in Multiple Endovascular Stroke trials (HERMES) meta-analysis of randomized controlled trials of EVT in acute stroke.⁸

MATERIALS AND METHODS

Literature Search

A comprehensive literature search of the databases PubMed, Ovid MEDLINE, and Ovid EMBASE was designed and conducted by an experienced librarian with input from the authors. The key words “stroke,” “mechanical thrombectomy,” “elderly,” “octogenarian,” “nonagenarian,” and “outcomes” were used in “AND” and “OR” combinations. The search was limited to articles published from 1992 to week 35 of 2017. All studies reporting outcomes of mechanical thrombectomy in patients older than 80 years of age were included. Inclusion criteria were the following: 1) English language; 2) studies reporting patients with acute stroke treated endovascularly in a patient population 80 years of age or older; and 3) studies reporting at least 3 months of clinical follow-up. Studies including patients treated with older endovascular modalities (ie, Merci retriever [Concentric Medical, Mountain View, California], intra-arterial thrombolysis alone, and so forth) were excluded. Two of the authors selected the included studies (C.A.H., P.N.). Differences were resolved by a senior author with 5 years of experience in meta-analysis (W.B.).

Outcomes and Data Extraction

For each study, we extracted the following baseline information: number of patients, mean or median age, imaging selection criteria, clinical selection criteria, exclusion criteria, mean or median NIHSS score, mean or median ASPECTS, endovascular technique used, type of study (prospective or retrospective), number of patients achieving a good functional outcome (defined as a modified Rankin Scale score of ≤ 2 at 3 months), number of patients in whom successful recanalization was achieved (defined as TICI 3 or 2b), procedure-related complications (eg, vessel perforation and groin complications), number of patients with intracranial hemorrhage and whether this was symptomatic, time from symptom onset to intervention (defined as groin puncture), and procedure time.

The primary end point of this study was good functional outcome at 3 months. Secondary outcomes included the following: mortality at 3 months, rate of successful revascularization, rate of procedure-related complications, rate of intracranial hemorrhage including symptomatic intracranial hemorrhage, time from symptom onset to groin puncture, and procedure time.

METHODOLOGIC QUALITY

We modified the Newcastle-Ottawa Quality Assessment Scale to assess the methodologic quality of the studies included in this meta-analysis. This tool is designed for use in comparative studies; however, because the studies did not include a control group, we assessed study risk of bias based on selected items from the tool, focusing on the following questions: 1) Did the study include all patients or consecutive patients versus a selected sample? 2) Was the study retrospective or prospective? 3) Was clinical follow-up satisfactory, thus allowing ascertainment of all outcomes? 4) Were outcomes clearly reported?

Statistical Analysis

We estimated from each cohort the cumulative incidence (event rate) and 95% confidence interval for each outcome. Confidence intervals were symmetric and 2-sided. Event rates for each intervention were pooled in a meta-analysis using the random-effects model.²⁵ Anticipating heterogeneity between studies, we chose this model a priori because it incorporates within-study variance and between-study variance. Heterogeneity of treatment effect across studies was evaluated using the I^2 statistic, in which $I^2 > 50\%$ suggests substantial heterogeneity.²⁶ A sensitivity analysis was conducted to compare the primary end point (good functional outcome at 90 days) among the included studies, depending on whether they reported an exclusion criterion of a pre-morbid mRS of ≥ 2 . Statistical analysis was conducted using STATA software, Version 14 (StataCorp, College Station, Texas).

RESULTS

Study Selection and Characteristics

The initial literature search yielded 1072 articles. On review of the abstracts and titles, we excluded 1008 articles. Sixty-four articles were selected for full-text screening. Of these, 24 were excluded (case reports, conference abstracts, review articles, and 1 article written in Russian). In all, 40 potentially relevant articles were identified. Of these, a further 23 were excluded for the following reasons: Some studies did not separate patients by age, the definition of elderly was too young, some studies reported outdated endovascular techniques (primarily those that included patients treated with the first-generation Merci device), and 1 study only provided the mRS score at 30 days rather than at 90 days.

In total, 17 articles including 860 patients aged 80 years or older who underwent EVT for acute stroke were included. In all studies, either a stent retriever alone was used as the primary method of recanalization or a combination of stent retriever plus or minus aspiration. One study included patients who had undergone intra-arterial thrombolysis in addition to stent-retriever and/or aspiration.

Five studies were prospective, and 12 were retrospective. Overall, 3 were considered to have a low risk of bias (using a modified Newcastle-Ottawa Quality Assessment Scale), whereas 14 had limitations. Clinical and imaging inclusion criteria for EVT varied considerably in the studies (On-line Table). Eight of the 17 included studies only performed EVT in patients with acute stroke with a pre-morbid mRS of ≤ 2 . All except 5 studies clearly defined their inclusion criteria. All included studies reported mRS at 3 months. The smallest study included 16 patients, and the largest study included 111 patients. A summary of included stud-

ies is provided in On-line Table. A flow chart for study selection is provided in Fig 1. The mean/median presentation NIHSS score was 17.8 (available for 16 studies). Seven studies included presentation ASPECTS with their results, the mean of which was 8.5.

Clinical Outcomes

Overall good functional outcome (mRS ≤ 2) was achieved in 27% of patients (95% CI, 21%–32%). A meta-analysis forest plot for good functional outcome at 3 months is shown in Fig 2. Overall mortality (reported by 14 studies) was 34% (95% CI, 23%–44%). Successful recanalization (reported by 15 studies) was achieved in 78% of patients (95% CI, 72%–85%). Procedure-related complication rates (reported by 4 studies) were 11% (95% CI, 4%–17%). The incidence of intracranial hemorrhage (reported in 10 studies) was 24% (95% CI, 15%–32%), and for symptomatic intracranial hemorrhage (reported by 14 studies), the incidence was 8% (95% CI, 5%–10%). The overall reported mean time to groin (documented in 13 studies) was 251 minutes (95% CI, 224–278 minutes). Procedural time in these elderly patients was 99 minutes

(95% CI, 67–131 minutes). A summary of the results is shown in the Table.

Sensitivity Analysis

Eight of the 17 included studies only performed EVT in patients with a premorbid mRS of ≤ 2 . A statistically significantly higher proportion of patients achieved a good functional outcome at 90 days (mRS ≤ 2) in those studies that only included pre-morbidly independent patients compared with those studies that did not have this prerequisite ($P = .007$). The mean rate of good functional outcome at 90 days in the studies with this inclusion criterion was 31.7% (95% CI, 18.4%–41.1%) compared with 20.4% (95% CI, 16.1%–24.7%) in those studies that did not use this criterion.

Heterogeneity

Heterogeneity of treatment effect across studies was evaluated using the I^2 statistic, in which $I^2 > 50\%$ suggests substantial heterogeneity.²⁶ Substantial heterogeneity was noted for all outcomes. The calculated heterogeneity of treatment effect for good neurologic outcome at 3 months was $I^2 = 70\%$.




FIG 1. Flow chart for study selection for studies included.




FIG 2. Meta-analysis forest plot for good functional outcomes at 3 months.

DISCUSSION

This systematic review and meta-analysis of 16 observational studies^{9–24} and data from the HERMES study,⁸ including 860 patients 80 years of age or older who underwent mechanical thrombectomy for acute ischemic stroke, has demonstrated a number of compelling findings. First more than one-quarter (27%) of patients achieved good functional outcome (functionally independent) at 90 days, while one-third (34%) of patients died. Excellent rates of TICI 2b/3 recanalization were achieved, close to 80%. Symptomatic intracranial hemorrhage rates were 8%, and procedural complication rates were approximately 11%. These findings suggest that mechanical thrombectomy provides substantial benefit to a sizeable proportion of octogenarians, despite the associated comorbidities in this cohort of patients. More

Summary of clinical outcomes

Outcome		Heterogeneity (I ²)
Good functional outcome at 3 mo	27% (95% CI, 21%–32%)	70%
Mortality within 3 mo	34% (95% CI, 23%–44%)	91%
Successful revascularization	78% (95% CI, 72%–85%)	83%
Procedure-related complication	11% (95% CI, 4%–17%)	56%
Any ICH	24% (95% CI, 15%–32%)	83%
Symptomatic ICH	8% (95% CI, 5%–10%)	61%
Time to groin (min)	251 (95% CI, 224–278)	88%
Procedure time (min)	99 (95% CI, 67–131)	98%

Note:—ICH indicates intracranial hemorrhage.

work is, however, still needed to further improve stroke outcomes in this patient population. These findings are particularly important, given that this demographic group will likely represent a growing proportion of mechanical thrombectomy candidates as the life expectancy of the general population continues to increase.

Ischemic stroke has increasing incidence with age and tends to present with more severe symptoms and has worse outcomes in the elderly compared with younger patients. Thus, it is of interest to determine the safety and efficacy of EVT in this older age group. The randomized trials that have established the remarkable effectiveness of EVT for patients with stroke with large-vessel occlusions do not conclusively answer the question of the value of EVT for octogenarians in daily practice. Exclusion of patients older than 80 years in some trials^{5–7} and the relatively low number of these patients compared with other age groups in those trials that did not consider age among the inclusion criteria^{27–29} are limiting factors that must be taken into consideration when interpreting the data from these randomized controlled studies.

The HERMES meta-analysis⁸ of 5 randomized controlled trials of EVT ($n = 1287$ including 634 assigned to mechanical thrombectomy; mean age, 68 years) reported good functional outcomes in 46% of patients and a mortality rate of 15%. However, among patients 80 years of age and older in the EVT arms, the rates were 29.8% for good 90-day functional outcome and 28% for mortality. These results are comparable with those that we obtained through our meta-analysis of observational studies (which also included data for octogenarians from the HERMES meta-analysis). Our meta-analysis did demonstrate a higher rate of symptomatic intracranial hemorrhage than the randomized trials (8% compared with 2%–5% across all age groups in the trials^{8,30}) as well as high rates of procedure-related complications (11%).

Meta-analysis of trial data clearly demonstrated that octogenarians treated with EVT had markedly better chances of achieving a good functional outcome compared with patients not treated with EVT⁸ (common odds ratio = 3.68; 95% CI, 1.95–6.92), in great part because of the poor outcomes observed in the latter group. Thus, EVT should be offered to patients aged 80 years and older. The challenge remains how to identify those older patients who will not benefit from recanalization to avoid futile and potentially harmful interventions. Eight of the 17 studies, including the HERMES data, included only octogenarians who were premorbidly functionally independent ($mRS \leq 2$). The mean rate of good functional outcome

($mRS < 2$) at 90 days was higher in studies that included only premorbidly independent patients compared with those that did not (20% compared with 32%). This finding highlights the need for careful patient selection for EVT in older patients. An inclusion criterion of only treating patients with a premorbid mRS of ≤ 2 may still exclude a large proportion of patients with stroke (of any age) who could benefit from EVT.

An mRS of 3 may convey a good quality of life in a patient older than 80 years of age, and the potential “shift effect” of EVT toward a lower mRS may still be regarded as a very favorable outcome.

Limitations

While data in this meta-analysis were derived from a strictly defined patient population with at least 3 months of follow-up, our study has limitations. In our assessment of study risk of bias, most of the studies had a medium risk of bias. Publication bias is also possible because centers with better results in treating older patients with stroke may have been more prone to report their results (however, if present, this bias would overestimate the chances of recovery after EVT). I^2 values were $>50\%$ for all outcomes, indicating substantial heterogeneity. Marked variability in patient imaging and clinical inclusion and exclusion criteria across studies may have contributed to this heterogeneity. The 2 studies that reported the worst outcomes included only patients older than 90 years. If these studies were excluded from the meta-analysis, the overall proportion of patients with good neurologic outcome would have slightly increased to 28%. Few studies provided detailed and quantifiable descriptions of the premorbid condition of the patients they treated (On-line Table). Overall, the quality of evidence (confidence in estimates) is low because of imprecision, heterogeneity, and methodologic limitations of the included studies.^{31,32} Due to lack of availability of data from the individual studies, it was not possible to stratify outcomes on the basis of stroke onset-to-recanalization time, concomitant use of thrombolytic therapy, type of anesthesia used, or patient collateral or perfusion mismatch status. It was not possible to perform an mRS shift analysis from the data collected from the included studies, and this issue may have underestimated the benefit of EVT for acute stroke in elderly patients.

CONCLUSIONS

A large proportion of patients 80 years of age or older with severe stroke from large-vessel occlusion can benefit from EVT, and good recanalization rates can be achieved. Our findings suggest that age should not be a discriminator of whether EVT should be attempted in acute stroke. The outcomes in older patients are, however, poorer than in younger patients. When one counsels patients and families, they should be informed that recovery of independent function can be expected in approximately 1 in 4 cases. Further research is necessary to refine selection criteria for EVT in this age group.

REFERENCES

- United States Census Bureau. 2014 National Population Projections Tables. <https://www.census.gov/data/tables/2014/demo/popproj/2014-summary-tables.html>. Accessed August 29, 2017
- Eurostat. Statistics Explained. People in the EU—population projections. http://ec.europa.eu/eurostat/statistics-explained/index.php/People_in_the_EU_-_population_projections#Europop2013_.E2.80.94_population_projections. June 2015. Accessed August 29, 2017
- Writing Group Members, Mozaffarian D, Benjamin E, Go A, et al; American Heart Association Statistics Committee; Stroke Statistics Subcommittee. **Heart Disease and Stroke Statistics: 2015 Update—a report from the American Heart Association.** *Circulation* 2016;133:e38–360 CrossRef Medline
- Saver JL, Goyal M, van der Lugt A, et al; HERMES Collaborators. **Time to treatment with endovascular thrombectomy and outcomes from ischemic stroke: a meta-analysis.** *JAMA* 2016;316:1279–88 CrossRef Medline
- Jovin TG, Chamorro A, Cobo E, et al; REVASCAT Trial Investigators. **Thrombectomy within 8 hours after symptom onset in ischemic stroke.** *N Engl J Med* 2015;372:2296–306 CrossRef Medline
- Saver JL, Goyal M, Bonafe A, et al; SWIFT PRIME Investigators. **Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke.** *N Engl J Med* 2015;372:2285–95 CrossRef Medline
- Bracard S, Ducrocq X, Mas JL, et al; THRACE investigators. **Mechanical thrombectomy after intravenous alteplase versus alteplase alone after stroke (THRACE): a randomised controlled trial.** *Lancet Neurol* 2016;15:1138–47 CrossRef Medline
- Goyal M, Menon BK, van Zwam WH, et al; HERMES collaborators. **Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials.** *Lancet* 2016;387:1723–31 CrossRef Medline
- Broussalis E, Weymayr F, Hitzl W, et al. **Endovascular mechanical recanalization of acute ischaemic stroke in octogenarians.** *Eur Radiol* 2016;26:1742–50 CrossRef Medline
- Kurre W, Aguilar-Pérez M, Niehaus L, et al. **Predictors of outcome after mechanical thrombectomy for anterior circulation large vessel occlusion in patients aged ≥ 80 years.** *Cerebrovasc Dis* 2013;36:430–36 CrossRef Medline
- Ribo M, Flores A, Mansilla E, et al. **Age-adjusted infarct volume threshold for good outcome after endovascular treatment.** *J Neurointerv Surg* 2014;6:418–22 CrossRef Medline
- Azkune Calle I, Bocos Portillo J, Anton-Ladislao A, et al. **Clinical outcome of mechanical thrombectomy for stroke in the elderly.** *J Stroke Cerebrovasc Dis* 2017;26:582–88 CrossRef Medline
- To CY, Rajamand S, Mehra R, et al. **Outcome of mechanical thrombectomy in the very elderly for the treatment of acute ischemic stroke: the real world experience.** *Acta Radiol Open* 2015;4:2058460115599423 CrossRef Medline
- Cohen JE, Gomori JM, Leker RR. **Stent retriever-based thrombectomy in octogenarians.** *Interv Neurol* 2016;5:111–17 CrossRef Medline
- Khan MA, Baird GL, Miller D, et al. **Endovascular treatment of acute ischemic stroke in nonagenarians compared with younger patients in a multicenter cohort.** *J Neurointerv Surg* 2016;9:727–31 CrossRef Medline
- Kleine JF, Boeckh-Behrens T, Prothmann S, et al. **Discrepancy between early neurological course and mid-term outcome in older stroke patients after mechanical thrombectomy.** *J Neurointerv Surg* 2016;8:671–76 CrossRef Medline
- Möhlenbruch M, Pfaff J, Schönenberger S, et al. **Endovascular stroke treatment of nonagenarians.** *AJNR Am J Neuroradiol* 2017;38:299–303 CrossRef Medline
- Sallustio F, Koch G, Motta C, et al. **Efficacy and safety of mechanical thrombectomy in older adults with acute ischemic stroke.** *J Am Geriatr Soc* 2017;65:1816–20 CrossRef Medline
- Son S, Kang DH, Hwang YH, et al. **Efficacy, safety, and clinical outcome of modern mechanical thrombectomy in elderly patients with acute ischemic stroke.** *Acta Neurochir (Wien)* 2017 Jul 2. [Epub ahead of print] CrossRef Medline
- Zacharatos H, Hassan AE, Vazquez G, et al. **Comparison of acute nonthrombolytic and thrombolytic treatments in ischemic stroke patients 80 years or older.** *Am J Emerg Med* 2012;30:158–64 CrossRef Medline
- Lima A, Haussen DC, Rebello LC, et al. **Endovascular therapy for large vessel stroke in the elderly: hope in the new stroke era.** *Cerebrovasc Dis* 2016;42:421–27 CrossRef Medline
- Castonguay A, Zaidat O, Novakovic R, et al. **Influence of age on clinical and revascularization outcomes in the North American Solitaire Stent-Retriever Acute Stroke Registry.** *Stroke* 2014;45:3631–36 CrossRef Medline
- Parrilla G, Carreón E, Zamarro J, et al. **Recanalization and mortality rates of thrombectomy with stent-retrievers in octogenarian patients with acute ischemic stroke.** *Cardiovasc Intervent Radiol* 2015;38:288–94 CrossRef Medline
- Figueiredo S, Carvalho A, Rodrigues M, et al. **Endovascular stroke treatment of patients over 80 years old: cumulative evidence from the “real world.”** *J Stroke Cerebrovasc Dis* 2017;26:2949–53 CrossRef Medline
- DerSimonian R, Laird N. **Meta-analysis in clinical trials.** *Control Clin Trials* 1986;7:177–88 CrossRef Medline
- Higgins JP, Thompson SG, Deeks JJ, et al. **Measuring inconsistency in meta-analyses.** *BMJ* 2003;327:557–60 CrossRef Medline
- Campbell BCV, Mitchell PJ, Kleinig TJ, et al; EXTEND-IA Investigators. **Endovascular therapy for ischemic stroke with perfusion-imaging selection.** *N Engl J Med* 2015;372:1009–18 CrossRef Medline
- Goyal M, Demchuk AM, Menon BK, et al; ESCAPE Trial Investigators. **Randomized assessment of rapid endovascular treatment of ischemic stroke.** *N Engl J Med* 2015;372:1019–30 CrossRef Medline
- Berkhemer OA, Fransen PS, Beumer D, et al. **A randomized trial of intraarterial treatment for acute ischemic stroke.** *N Engl J Med* 2015;372:11–20 CrossRef Medline
- Campbell B, Hill M, Rubiera M, et al. **Safety and efficacy of Solitaire stent thrombectomy: individual patient data meta-analysis of randomized trials.** *Stroke* 2016;47:798–806 CrossRef Medline
- Guyatt GH, Oxman AD, Kunz R, et al. **GRADE guidelines, 6: rating the quality of evidence—imprecision.** *J Clin Epidemiol* 2011;64:1283–93 CrossRef Medline
- Balshem H, Helfand M, Schünemann HJ, et al. **GRADE guidelines, 3: rating the quality of evidence.** *J Clin Epidemiol* 2011;64:401–06 CrossRef Medline