

On-line Table 1: Characteristics of the patient cohort and different subgroups

	All Ruptured Aneurysms	Ruptured Aneurysms with 6-Month Angiographic Follow-Up	Adequately Occluded Aneurysms at 6 Months	Incompletely Occluded Aneurysms at 6 Months
Patients/aneurysms	901	713	589	124
Men	290 (32%)	226 (32%)	180 (31%)	46 (37%)
Mean age (yr)	53.5 (median, 52; range, 11–89)	52.4 (median, 52; range, 11–78)	52.2 (median, 51; range, 11–78)	53.0 (median, 52; range, 25–75)
Mean aneurysm size (mm)	7.7 (median, 6; range, 2–35)	7.7 (median, 6; range, 2–35)	6.9 (median, 6; range, 2–26)	11.4 (median, 10; range, 2–35)
Small (0–5 mm)	365 (41%)	291 (41%)	267 (45%)	24 (19%)
Medium (6–10 mm)	344 (38%)	270 (38%)	226 (38%)	44 (35%)
Large (11–24 mm)	182 (20%)	143 (20%)	95 (16%)	48 (39%)
Giant (≥ 25 mm)	10 (1%)	9 (1%)	1 (0%)	8 (6%)
Anterior circulation	688 (76%)	537 (75%)	451 (77%)	86 (69%)
AcomA	339	260	229 (39%)	31 (25%)
PcomA	164	126	105	21
MCA	86	66	51	15
Pericallosal	36	34	31	3
Carotid tip	26	20	14	6
Ophthalmic	20	18	9	9
Ant Choroidal	15	11	11	0
Hypophyseal	1	1	0	1
Cavernous carotid	1	1	1	0
Posterior circulation	213 (24%)	176 (25%)	138 (23%)	38 (31%)
Basilar tip	122	107	78 (13%)	29 (23%)
PICA	35	24	19	5
SCA	27	22	22	0
Vertebral artery	12	8	8	0
Basilar trunk	9	7	5	2
PCA	7	7	5	2
AICA	1	1	1	0